

<p>U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) WASHINGTON, D.C. 20460</p> <p><b>NOTICE OF PESTICIDE:</b> <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i></p>	<p>EPA REGISTRATION NO. <i>Transferred to 8660-108</i> <del>ATOS 800</del></p> <p>DATE OF JANCE MAR 30 1983</p> <p>TERM OF ISSUANCE Until re-registration</p> <p>NAME OF PESTICIDE PRODUCT Dursban DE™ 1.25 Galf Insecticide</p>
<p>NAME AND ADDRESS OF REGISTRANT (Include ZIP code)</p> <p>Patterson Green-Up Co. Div. of Curry Cartwright, Inc. 1331 Union Avenue Kansas City, MO 64101</p> <p style="text-align: right;"><i>161/89297</i> <i>18/1</i></p>	
<p><b>NOTE:</b> Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p>	
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p><i>See side panel for additional precautions</i></p>	
<p><input checked="" type="checkbox"/> ATTACHMENT IS APPLICABLE</p> <p>SIGNATURE OF APPROVING OFFICIAL <i>[Signature]</i> DATE <i>[Date]</i></p>	

**BEST AVAILABLE COPY**

c. On page 4 of 4 insert the following sentence in the peach tree borer directions (immediately after the statement "This treatment is for ornamentals only"): "Do not use on trees with edible fruit."

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca  
Product Manager (15)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767)

Enclosures

**BEST AVAILABLE COPY**

3 of 6

DURSBAN-DDVP  
1.25  
TURF INSECTICIDE

Control of Ants, Chinch Bugs, Sod Webworms  
and Numerous Other Pests on Turf and  
Ornamental Plants. For Area Control of  
Ticks and Chiggers

ACCEPTED  
with COMMENTS  
to EPA Letter Dated:

MAR 30 1983

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
~~2169-284~~

For Commercial/Industrial Use Only

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro 2-pyridyl) phosphorothioate] .....	12.70%
2,2-Dichlorovinyl dimethyl phosphate .....	3.19%
Related Compounds .....	0.24%
Aromatic Petroleum Solvent .....	77.75%
INERT INGREDIENTS: .....	6.12%
Total	100.00%

Transferred to  
8660 108

Contains 1 lb. Chlorpyrifos & .25 lb. Dichlorvos (DDVP) per Gallon  
EPA REG. NO. 2169-284 EPA Est. No. 2169-KS-1

KEEP OUT OF REACH OF CHILDREN <

WARNING <

See Side Panel For Additional Precautionary Statements

-10 pt type  
14 pt type

STATEMENTS OF PRACTICAL TREATMENTS

If Swallowed - Call a physician or Poison Control Center immediately.

If possible vomiting should be induced under medical supervision.

Have victim drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.

If On Skin - Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes - Flush eyes with plenty of water. Call a physician immediately.

Note To Physician - Chlorpyrifos and DDVP are cholinesterase inhibitors. Treat symptomatically. Atropine only by injection is an Antidote.

Net Contents \_\_\_\_\_ One Gallon

PATTERSON GREEN-UP COMPANY  
DIV CURRY CARTWRIGHT INC.  
1331 UNION AVE, BOX 4029  
KANSAS CITY, MISSOURI 64101

**BEST AVAILABLE COPY**

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

**WARNING:** May be fatal if swallowed. May be absorbed through the skin. May be injurious to eyes and skin. Keep children and pest off treated areas until spray has dried. Do not take internally. Do not get in eyes or on skin. Do not smoke while using. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before re-use. Flush contaminated eyes with plenty of water and get medical attention. Avoid breathing spray mist or vapors. Do not spray vegetable gardens or other plants used for food purposes.

**BEST AVAILABLE COPY**

PHYSICAL AND CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and wildlife. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated areas. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

STORAGE & DISPOSAL - BEST COPY AVAILABLE 10/27/2020

Do not contaminate water, food or feed by storage or disposal.  
**Pesticide Disposal** - Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.  
**Container Disposal** - Triple rinse ( or equivalent ). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

**Notice** - Buyer assumes all risk and liability of use, storage and/or handling of material not in accordance with the terms of this label.

PESTS OUT-OF-DOORS

**General Information-** Use this product for control of certain pests on outside surfaces of buildings and on lawns and ornamental plants, and for area control of Chiggers. Consult the following directions for dosage and application details to control the pests listed.

**PESTS ON OUTSIDE SURFACES OF BUILDINGS:** Use this product at the rate of 5 1/2 ounces per gallon of water (0.5% Chlorpyrifos emulsion) to control Ants, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Mosquitoes, Sowbugs, Spiders, and Ticks by application as a residual spray to outside surfaces of buildings including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Do not treat or allow the spray to contact plants (lawns, flowering plants, shrubs, trees, evergreens, etc.) because of possible injury. Repeat treatment as needed to maintain effectiveness.

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE (cont.)

**NOTE-**To help prevent infestations of buildings by Ants, Crickets, Earwigs, Millepedes, and Sowbugs (Pillbugs) treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also buildings foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 32 ounces (1 quart) in 100 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1,000 sq.ft. to thoroughly and uniformly wet the band area.

**PESTS OF LAWNS, OTHER ORNAMENTAL AND RECREATIONAL TURF AREAS AND PERENNIAL TURF GRASSES GROWN FOR SEED:** Use this product to control pests listed in the following table by application at the recommended dosages and in accordance with the directions given below. Dilute in enough water to obtain complete and uniform coverage of pest infested areas and apply as a coarse, low pressure spray using suitable spray equipment. Thoroughly water after treatment to wash the insecticide into the turf, except as noted. For best results, the lawn should be moist at time of treatment.

DOSAGES AND DIRECTIONS FOR USE ON TURF **BEST EQUIPMENT AVAILABLE**

PEST	DOSAGE per 1,000 sq.ft.	REMARKS
Ants, Armyworms, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Ticks, Sod Webworms.	*3 Fl. Ounces  Use in a minimum of 2 1/2 to 3 Gallons water per 1,000 sq.ft.	Spray when pests first appear; retreat when needed. For Sod Webworms delay watering or mowing the treated area for 12 to 24 hours after treatment.
Turfgrass Hyperodes	6 Fl. Ounces	Spray suspected problem areas in Mid-April and in Mid-May or as recommended by Local Agricultural Extension Service Specialist.
White Grubs of European Chafer and Japanese Beetle only.	6 to 12 Ounces	Spray when Grubs are actively feeding near the soil surface in late July and August or as recommended by Extension Specialist. Water treated area after treatment with 1/2 to 1 inch of water to wash insecticide into soil.
Bluegrass Billbug	6 to 12 Ounces	Spray early in the season when adults first appear. Retreat when needed.

\*3 Fluid Ounces/1,000 sq.ft = 1 gallon per acre.

**AREA CONTROL OF TICKS AND CHIGGERS-** Use for noncrop areas such as roadsides, footpaths, and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply 1 quart per acre in sufficient water to mist all areas including low underbrush, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage usually 40 to 100 gallons of water per acre.

**ATTENTION-** Do not allow public use of treated areas during application or until spray has dried.

**BEST AVAILABLE COPY**

DIRECTIONS FOR USE (Cont.)

**PESTS OF ORNAMENTALS;** Use this product to treat flowers, shrubs, vines, and flowering trees, and evergreens found to be infested with pests listed in the following table. Dilute with water according to directions given in the table and apply using suitable hand or power spray equipment in a manner to provide complete coverage. For best results apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid over spraying. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. **CAUTION:** Do not use on oleas, Camellias, poinsettias, rose bushes, or variegated Ivy because of possible injury to foliage.

**RATES AND DILUTIONS FOR USE ON ORNAMENTAL PLANTS**

Pests	Amount of Concentrate to use in water to Make		
	3 Gals.	10 Gals.	100 Gals.
Leafhoppers, Mites, Bagworms, Angestriped Oakworm, Terr. Tent Caterpillar, Applebug, Fall Webworm, Beetles, Grasshoppers, Downy Necked Caterpillar	1 fl.oz.	3.2 fl.oz.	2 pints
Leafhoppers, Aphids, Mimosa Webworm, Redhumped Caterpillar, Aphids (Exposed stages)	2 fl.oz.	6.4 fl.oz.	2 quarts
White Borer, Lilac Borer, Elm Bark Borer, Scale Insects, Oak Lecanium, Pear Thaw, Tea, and Peach Scale.	4 fl. oz.	12.8 fl.oz.	1 gallon
White Tree Borer	11.6 fl.oz.	38.5 fl.oz.	3 Gallon

**ELM BARK BORER** - Mix 1 gallon in 100 gallons water (1 quart in 25 gals.) and spray trunks and lower limbs of trees when adults begin to emerge. Consult your local Agricultural Extension Service for the proper timing.

**WHITE TREE BORER**- Dilute 11.5 fl.oz. in 3 gallons of water. Apply to flowering trees and shrubs of the Genus PRUNUS. Thoroughly spray trunk at ground level up to the first limbs. Apply as coarse wet spray 3 times May 1st; June 1st; and July 1st. Do not treat the fruit or upper portions of the tree. This treatment is for ornamentals only.

**ADDITIONAL INFORMATION**- MOSQUITOES coming to rest on areas treated for control of Turf Ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

**BEST AVAILABLE COPY**