

LaRoche Industries Inc.

CONTAINER PRINT SPECIFICATIONS

Subject	Vertagreen Diazinon Pre-Weed Plus	Account: VG50078
		Page: 1 of 5
		Date: 11/5/86/86

Copy for Front

This product must not be used on golf courses and sod farms

ACCEPTED

APR 27 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 3442-827

VERTAGREEN
 DIAZINON PRE-WEED PLUS
 CONTAINS
 INSECTICIDE PLUS BALAN™ PRE-EMERGENT HERBICIDE

Transferred to
8660-115

Plus 30-5-5 Fertilizer

For Lawns and Ornamental Turf

Pre-Emergence Control of Crabgrass, Foxtail, Watergrass and
Annual Blue Grass

LAWN INSECT CONTROL

For Control of Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Armyworms, Billbugs, Brown Dog Ticks, Crickets, Cutworms, Earwigs, Fleas, Sowbugs (Pillbugs), Springtails, Bermudagrass Mites, Chiggers, Clover Mites, Leafhoppers, & Millipedes Infesting Home Lawns and Other Ornamental and Recreational Turf Grass Areas.

Keep Out of Reach of Children
WARNING;

Read Complete Cautions on Back Panel

NOTE: Regarding above copy, minimum type size is:
18 Pt. for **WARNING**
12 Pt. for **KEEP OUT OF REACH OF CHILDREN**

ACTIVE INGREDIENTS:	By Weight
Diazinon [0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....	5.00%
N-Butyl-N-Ethyl-alpha-alpha-alpha Trifluoro-2,6-Dinitro-p-Toluidine.....	2.06%
INERT INGREDIENTS:	92.94%
Total	100.00%

Diazinon a trademark of CIBA GEIGY for Insecticide
U. S. Patent No. 2,754,243 and other patents pending
BALAN The Trademark of BENEFIN Elanco Product Company
a division of Eli Lilly & Company

Subject	Vertagreen Diazinon Pre-Weed Plus	Account: VG50078 Page 2 of 5 Date: 11/5/86
----------------	-----------------------------------	--

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED -- Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED -- Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN -- Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES -- Flush eyes with plenty of water. Call a physician if irritation persists.

Net Weight 40 lbs. - Covers 20,000 Sq. Ft. of Turf
 EPA Reg. No. 3442-827
 EPA Est. No. 3442-FL-2

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

This product is designed for a commercial or home-owner annual Lawn Service Program which features 1 to 2 granule insecticide + pre-emergent herbicide applications per year. Insect control will be effective when applied anytime during insect activity periods. The weed control benefit will only be effective when applied before weed seeds germinate. Apply this product when the grass is dry and water in lightly after application.

INSECTS AND WEEDS -- For the control of Ants, Armyworms, Bermudagrass Mites, Billbugs, Brown Dog Ticks, Chiggers, Chinch Bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Leafhoppers, Millipedes, Sod Webworms, Sowbugs, and Springtails. Pre-emergent weed control of Annual Bluegrass (Poa Annua), Smooth & Hairy Crabgrass, Goosegrass (Silver Crabgrass or Crowfoot), Watergrass (Barnyardgrass) and Yellow & Green Foxtail.

FIRST APPLICATION -- Apply 40 lbs. per 20,000 sq. ft. The first application should be made 10 to 20 days before your local "Frost Free" date in the spring. This will control existing insects (as listed on this label) along with pre-emergence control of Crabgrass, goosegrass, foxtail, water grass and certain broadleaf weeds; namely carpetweed, chickweed, lambsquarter, purslane and pigweed.


Subject	Vertagreen Diazinon Pre-Weed Plus	Account: VG50078
		Page: 3 of 5
		Date: 11/5/86

SECOND APPLICATION -- Apply 40 lbs. per 20,000 sq. ft. Turf that has been power raked, aerated, or verticut will benefit from this application since spring rains encourage additional weed seed germination. Apply in late May or early June after the first treatment. Insect control will be effective as in the first application, however only the ungerminated weed seeds will be controlled. Established weeds will not be effected.

ANNUAL BLUEGRASS (Poa Annua) -- Apply 40 lbs. per 20,000 sq. ft. Treatment of heavy infestations will result in "thin" turf. Observe good cultural practices such as fertilization and watering to fill in these areas. Repeat treatment the following year for best control.

NORTHERN COOL SEASON GRASS should be treated in the late summer to early fall (late-July to late-September) or in early spring prior to annual bluegrass seed germination. It is difficult to get 100% control the first year in heavy turf growth.

WEED CONTROL FOR ANNUAL GRASSES



Approximate weed germination times (1) Crabgrass, Foxtail, Watergrass, Goosegrass. -- (2) Annual Bluegrass (Poa Annua).

INSECT CONTROL THROUGH THE YEAR -- If subsequent generations of insects appear in the turf after these applications, continue throughout the season with 5% Diazinon Crawling Insect Control Granules. EPA Reg. No. 3442-811 according to the directions on the label.

RESEEDING -- Delay reseeding for a least 6 weeks after application of this product.

WHERE TO USE -- When applied as directed the following grasses are tolerant -- perennial bluegrass, perennial ryegrass, centipedegrass, fescue, zoysiagrass, bermudagrass, St. Augustine grass and Bahia grass. Apply only to established stands of these grasses.

Subject	Vertagreen Diazinon Pre-Weed Plus	Account: VG50078
		Page 4 of 5
		Date: 11/5/86

STORAGE AND DISPOSAL

(Note: Heading must be minimum 12 Pt. type size)

Do not contaminate water, food, feed by storage or disposal.

Storage: Keep away from food, feedstuffs, and domestic water supplies. Keep container closed.

Disposal: Do not reuse empty container. Wrap and put in trash collection.

GUARANTEED FERTILIZER ANALYSIS

Nitrogen (N).....	30%
Available Phosphoric Acid (P ₂ O ₅).....	5%
Soluble Potash (K ₂ O).....	5%

Sources of Plant Nutrients: Urea, Diammonium Phosphate, Potassium Sulfate
Supplies .6 lb. N per 1,000 sq. ft. of turf when applied as directed.

SPREADER SETTING -- Scott-4, Cyclone-3-3/4.

Walking speed and spreader condition may require setting adjustments

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals

WARNING
KEEP OUT OF REACH OF CHILDREN

May be absorbed through the skin. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry. Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling and before smoking or eating.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Subject	Vertagreen Diazinon Pre-Weed Plus	Account: VG50078
		Page 5 of 5
		Date: 11/5/86

NOTICE: LaRoche Industries Inc. can accept no responsibility for injury to persons or animals or for loss or damage to turf, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this label.

Manufactured and Registered by
LaRoche Industries Inc.
1100 Johnson Ferry Road NE - Atlanta, GA 30342
Vertagreen is a Registered Trademark

VTG
UPC Code
25704-50078
Proof of Purchase

STORE IN A DRY PLACE

Below for Internal Use Only

APPROVALS - MARKETING	DATE	REGULATORY SRVS.	DATE
BY <i>Allen J. Foster</i>	<i>11/12/86</i>	<i>C. D. Anderson</i>	<i>11-12-86</i>