


8660-133
1 of 5

(FRONT PANEL)

GREENUP®

SEVIN® 11.7

LIQUID FLOWABLE



For Use on Flowers, Vegetables, Fruits, Lawns, Shrubs and Ornamentals

For Control of Aphids, Bagworms, Fleas, Gypsy Moth Larvae,
Japanese Beetle, Mosquitoes, Tent Caterpillars,
Tomato Hornworm, Sod Webworms, and Other Insects As Listed.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 11.7%

INERT INGREDIENTS: 88.3%

TOTAL 100.0%

Contains 1 lb. Carbaryl per gallon.

Sevin® is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

U.S. Patent No. 2,903,478 and 3,009,855


KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET CONTENTS _____

BEST AVAILABLE COPY



295

8660-133
2 of 5

(BACK PANEL)

NET CONTENTS _____

GREENUP®

SEVIN® 11.7

LIQUID FLOWABLE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or inhaled. Avoid breathing of spray mist. Do not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face thoroughly after handling and before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Give victim 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.


IF IN EYES: Flush eyes with plenty of water. If irritation persists get medical attention.

IF ON SKIN: Remove contaminated clothing and wash skin with soap and water.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. **BEE CAUTION:** This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area



BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For all uses, mix with water as directed. Keep spray mixture well agitated. Repeat applications at 7 to 10 day intervals for continued control. Do not rotate to any crop not listed on this label or other Carbaryl labels.

ORNAMENTAL INSECT CONTROL: Use 2-1/2 tbsp. per gal. of water (equal to 1 quart in 25 gals. of water) and apply when insects or their damage first appears. Spray thoroughly to cover both sides of foliage, fruit and stems. Repeat weekly or as required to control Aphids, Bagworms, Birch Leafminer, Blister Beetles, Flea Beetles, Gypsy Moth Larvae, Japanese Beetles, June Beetles, Lace Bugs, Leafhoppers, Leaf Rollers, Mealybugs, Mimosa Webworm, Oak Leaf Miner, Periodical Cicadae (17-year Locust), Plant Bugs, Psyllid, Rose Aphids, Tort Caterpillar, Thrips (exposed stages), and Willow Leaf Beetle on flowers, shrubs, ornamental and forest trees. Do not apply to Boston Ivy.

VEGETABLE INSECT CONTROL: Precautions in using: Apply when insects or their damage is first seen. Repeat at 7 to 10 day intervals, or as necessary. Do not apply within 1 day of harvest on all vegetable crops listed, except 3 days on Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi, Head Lettuce, Garden Beets (root), Horseradish, Radish, Parsnips, Rutabaga, Salsify and Turnips; and 14 days of harvest for Collards, Dandelions, Endive, Garden Beets (tops), Spinach, Swiss Chard, and Turnip Tops, Kale, Leaf Lettuce, Mustard Greens, Parsley and Salsify Tops.

To avoid possible injury to Lettuce and vining crops, do not apply when foliage is wet or when rain or excessive humidity is expected within the next 2 days.
NOTE: Application of this product to the tassel region of Corn during the pollen shed period will seriously reduce bee population.

EUROPEAN CORN BORER: Apply thoroughly into whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge.

CORN EARWORM: Apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Re-apply immediately after a rain.

HOW TO MIX

Use 2-1/2 tbsp. per gal. of water on young plants and 5 tbsp. per gal. of water on mature plants to control Corn Earworms, European Corn Borer, Fall Armyworms, Flea Beetles, Japanese Beetle, Leafhoppers, Velvetbean Caterpillar, and Western Bean Cutworm on Beans (Green, Lima, Snap, Cowpeas, and Black-eyed Peas); Flea Beetle, Harlequin Bug, Leafhopper, Armyworm, Imported Cabbage Worm, Corn Bagworm, Tarnished Plant Bug and Stink Bugs on Carrots, Chinese Cabbage, Collards, Dandelion, Endive, Garden Beets, Horseradish, Kale, Lettuce, Mustard Greens, Parsley, Parsnips, Rutabagas, Salsify, Spinach, Swiss Chard and Turnips.

BEST AVAILABLE COPY

Use only the low rate, therefore, 2-1/2 tbsp. per gal. (1 quart per 25 gals. water) for Pickleworms, Melonworms, Cucumber Beetle, Flea Beetle, Leafhoppers and Squash Bugs on Cucumbers, Melons, Pumpkins and Squash.


For Tomatoes, Potatoes, Eggplant and Peppers for the control of Colorado Potato Beetle, Flea Beetles, Leafhoppers, Tomato Fruit Worm, Tomato Hornworm, Tomato Pinworm and European Corn Borer Larvae, use 2-1/2 tbsp per gal. of water. May be applied up to day of harvest.

FRUIT TREE INSECT CONTROL FOR THE HOMEOWNER - Apple and Pear trees: Use 2-1/2 tbsp. per gal. of water. Apply to non-bearing trees 7 to 10 days after full leaf stage and repeat at 2 to 3 week intervals. On bearing trees, to avoid thinning of fruit, delay use for at least 30 days after bloom. Use at 10 to 14 day intervals to control Green Apple Aphid, Fruit Tree and Red Banded Leaf Roller, Bagworms, Plum Curculio and Codling Moth. Do not apply later than 1 day before fruit harvest.

PEACHES AND PLUM TREES - Use at the same rate, the same schedule and observe the same precautions for bearing trees. Controls Plum Curculio, Lesser Peach Tree Borer, Oriental Fruit Moth, Twig Borers, Fruit Tree Leaf Roller, and June Beetles. Do not apply later than 1 day before harvest.

FRUIT REDUCTION ON ORNAMENTAL TREES: When fruit production is not desired on Apple trees or Crataegus Sp. (Crabapples), spray 5 tbsp. per gal. of water directed particularly to the blooms at a time between 10 and 15 days after full bloom. One treatment is prescribed and complete fruit drop should not be expected.

LAWN INSECT CONTROL: Use one pint in 10 to 15 gals. of water to cover 500 sq. ft. of lawn surface. Use a suitable pressure type sprayer for control of Sod Webworm, Chinch Bugs, Ants, Earwigs, Fall Armyworms, Fleas, Leafhoppers, Millipedes, Mosquito, and Brown Dog Tick. For best results mow lawn first, remove thatch, and apply after irrigation. Repeat in 3 to 4 weeks. For Sod Webworms, apply May 15th and repeat at 30-day intervals through August.



595

8660-133
5 of 5

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.


WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. **FOLLOW DIRECTIONS CAREFULLY.** Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

**PARKER FERTILIZER CO.
201 4TH STREET
SYLACAUGA, AL 35150**



Date Printed: 06/23/92
Date Approved:
Supersedes:
Label Change: