

1 of 3

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED

**BEST AVAILABLE COPY**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

OCT 29 1984

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the purpose of  
registered under FIFRA Section 24(b).

2169-331

Transferred to  
8660-140

#### Homeowner Protective Clothing

Homeowner applicators must wear the following protective clothing during the application process: long-sleeved shirt, long pants, waterproof gloves, full foot coverings, and a hat.

**CAUTION:** Do not allow children to handle this product. Children and pets should not be allowed in treated areas until treated surfaces are dry.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING** — Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. May be fatal if swallowed or absorbed through the skin. Do not breathe vapor or spray mist.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Do not use on humans or household pets. Keep children and animals away from treated areas until spray has dried. Food utensils such as a teaspoon or measuring cup should not be used for food purposes after use with pesticides.

##### ENVIRONMENTAL HAZARDS

This product is toxic to fish, bees, and other wildlife. Do not apply directly to lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop cover or bees are blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

##### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Storage — Store in a cool dry place.

Pesticide Disposal — Pesticide, empty container, or free water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures.

Container Disposal — Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose in a sanitary landfill, or by other approved state and local procedures.



## LINDANE 20% INSECTICIDE

**INSECT CONTROL FOR ORNAMENTALS — FRUIT TREES — FLOWERS  
GRASS & VEGETABLES**

| ACTIVE INGREDIENTS                             |         |
|------------------------------------------------|---------|
| Lindane (Gamma Isomer of Benzene Hexachloride) | 20.00%  |
| Xylene                                         | 80.00%  |
| INERT INGREDIENTS                              |         |
|                                                | 13.15%  |
| TOTAL                                          |         |
|                                                | 100.00% |

**12 FT. WARNING: KEEP OUT OF REACH OF CHILDREN 4-8 FT.**  
See Side Panel For Additional Precautionary Statements

NET CONTENTS: 1 PINT

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See attached supplemental labeling for complete directions on ornamentals, fruit trees, flowers, bulbs, vegetables, livestock.

##### STATEMENT OF PRACTICAL TREATMENT

**If Swallowed** — Call a physician or Poison Control Center. Induce vomiting. Gastric lavage is indicated if more than 1 hour has passed. DO NOT INDUCE VOMITING vomiting may cause pneumonia.

**If Inhaled** — Remove victim to fresh air. Apply artificial respiration if indicated.

**If On Skin** — Remove contaminated clothing and areas with soap and water.

**If In Eyes** — Flush eyes with plenty of water. Get medical attention persists.

For additional precautions and complete directions product refer to attached brochure.

Notice: Seller makes no warranty, expressed or implied, in the use of this product other than indicated on the label. Assumes all risk of use and or handling of this material and or handling is contrary to label instructions.

EPA Reg No 2169-

EPA Est No 2169-

NET CONTENTS: 1 PINT

**BEST AVAILABLE**

P.O. Box 4029

patterson GREEN UP company

Kansas City, Missouri 64101

Division Curry-Carrwright, Inc.

#### STORAGE & DISPOSAL — 8 FT.

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE** — Store product in original container in a cool, dry, locked place out of reach of children.

**DISPOSAL** — Do not re-use empty container. Wrap container and put in trash.

| PLANTS                                         | INSECTS                                          | DOSAGE RATE IN FLUID MEASURE                                                   | COMMENTS                                                                                                                                                                                            |
|------------------------------------------------|--------------------------------------------------|--------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Elm, Birch, Locust, Holly                      | Leaf Miners                                      | 1 tsp./gal. water                                                              | Apply when adult miners appear in April & mid-May. Treat when smaller miners appear in July.                                                                                                        |
| Pines                                          | Pine Bark Beetles                                | 3 Fluid Ounces per gal. water                                                  | Apply to trunks up as high as sprayer will reach. Spray to point of run-off in April or May.                                                                                                        |
| Pine, Spruce                                   | White Pine Weevil                                | 1 tbsp./gal. water                                                             | Thoroughly spray entire tree in late April when buds swell.                                                                                                                                         |
| Tasuo, Hemlock                                 | Tasuo Weevil                                     | 1 tbsp./gal. water                                                             | Apply in May or early June to the trunks & around tree base. Soak the bark and soil surface.                                                                                                        |
| Dogwood, Lilac, Rhododendron                   | Dogwood Borer, Lilac Borer, Rhodo. Borer         | 1 tbsp./gal. water                                                             | Make 3 applications. May 1st; June 1st; July 1st. Treat limbs, twigs and base of trunk.                                                                                                             |
| Apple, Peach, Cherry                           | Bark Beetles Flat-headed & Round-headed Borer    | 1 tbsp./gal. water                                                             | Apply to bark of tree and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to runoff. Do not apply to fruit or foliage. Do not apply within 60 days of harvest.                   |
| Peaches                                        | Peach Tree Borer                                 | 1 tbsp./gal. water                                                             | Apply to bark of tree and large limbs in spring. Apply to trunk, crotch to ground. Soak bark to runoff. Do not apply within 60 days of harvest.                                                     |
| Maple, Aspen, Poplar, Ash, Willow, Elm, Locust | Aphids, Bark Beetles, Borers                     | 2 tbsp./gal. water                                                             | Apply in spring when buds begin to break. Repeat in 10-14 days. For Aphids, apply when present in summer & fall.                                                                                    |
| Birch                                          | Bronze Birch Borer                               | 1 tbsp./gal. water                                                             | Make 3 treatments. May 1st, June 1st; July 1st. Soak bark.                                                                                                                                          |
| As (Tulip, Iris, Glads, Lilies)                | Aphids, Bulb Fly, Iris Borer, Thrips             | 1 tbsp./gal. water                                                             | Treat when insects appear. For Iris borer soak soil around numerous roots May & June.                                                                                                               |
| Vegetable Sets - Cabbage                       | White Grubs, Cutworms, Wireworms, Soil Treatment | 1 tbsp./2 gal. water. Use 2 gal. of mixture to treat 250 sq. ft. of soil area. | Do not apply after thinning. Do not treat plant foliage. Do not use in soil to be used for root crops. Apply around base of plant within 24 hours after sets are placed in soil. Repeat in 14 days. |
| Malars, Peppers, Okra, Tomatoes                | Same for insects as for Cabbage                  | Same as for cabbage                                                            | Do not apply after edible parts start to form.                                                                                                                                                      |
| Flowers, Roses, Petunias, Carnations, etc.     | Aphids, Thrips, White Flies, Leafhoppers         | 1 tbsp./2 1/2 gal.                                                             | Apply as insect appear. Repeat sprays as needed for control. Thoroughly spray.                                                                                                                      |

**BEST AVAILABLE COPY**