

18 JUN 1992

Mr. Daniel Paradiso, Jr.
c/o Parker Fertilizer
201 West-Forth Street
Sylacauga, AL. 35150

Dear Mr. Paradiso:

Subject: Vertagreen Thiram 75
EPA Registration No. 8660-25
Vertagard MCPP Clover - Chickweed Killer
EPA Registration No. 8660-24
Vertagreen Weed And Feed
EPA Registration No. 8660-34
Protect 3 Way
EPA Registration No. 8660-104
Verta-Green St. Augustine Weed & Feed
EPA Registration No. 8650-147 ✓
Re: Label Amendment
Your Submission Dated May 22, 1992

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide act, as amended, is acceptable, provided that under the Environmental Hazard Section of the label replace the statement "Do not apply directly to water or wetlands" to read "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark." Please submit five (5) copies of your final printed labels incorporating this amendment. A stamped copy of labeling is enclosed for your records.

Sincerely

Joanne L. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (H-7505C)

CONCURRENCES

SYMBOL	H-7505C enclosure					
SURNAME	Melton					
DATE	6/17/92					

BEST AVAILABLE COPY

8660-147
1 of 4

(FRONT PANEL)

VERTAGREEN®

ST. AUGUSTINE

WEED & FEED

25-3-3

CONTAINS TRIMEC® STARBURST

ACCEPTED
with COMMENTS
in EPA Letter Dated

10 JUN 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

8660-147

For Use On St. Augustine, Bahia, Bermuda, Centipede and Fescue Grasses

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	0.159%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.508%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o- anisic acid	0.062%
INERT INGREDIENTS:	99.271%

Total 100.000%

Used at 20 lbs. on 5,000 sq. ft. this product contains:

*2.64 lbs. or 0.132% of 2,4-dichlorophenoxyacetic acid per ton,
or 0.23 lb. per acre
**8.40 lbs. or 0.420% of 2-(2-methyl-4-chlorophenoxy) propionic acid,
or 0.73 lb. per acre
***1.04 lbs. or 0.052% of 3,6-dichloro-o-anisic acid, or 0.09 lb. per acre

TRIMEC® is a Trademark of PBI Gordon Corporation
U.S. Patent No. 3,284,186

KEEP OUT OF THE REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with soap and water.
IF IN EYES: Immediately flush with plenty of water.
If irritation develops, get medical attention.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 20 LBS. (9.1 kg)

EPA Reg. No. 8660-147

EPA Est. 9198-OH-1

(BACK PANEL)

NET WEIGHT 20 LBS. (9.1 kg)

VERTAGREEN®

ST. AUGUSTINE

WEED & FEED

25-3-3

CONTAINS TRIMEC® STARBURST

GUARANTEED ANALYSIS

Total Nitrogen (N)	25.00%
1.1% Ammoniacal Nitrogen	
23.9% Urea Nitrogen	
Available Phosphoric Acid (P_2O_5)	3.00%
Soluble Potash (K_2O)	3.00%

Sources of Plant Nutrients: Ammonium Phosphate, Muriate of Potash, Sulfur Coated Urea, and Urea

Chlorine (Cl), Maximum	3.00%
----------------------------------	-------

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes, or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

This product can reach groundwater from improper handling. To minimize contamination from spills during loading and cleaning of equipment, take the following steps:

Handling: When handling this product, wear chemical resistant gloves. Wash nondisposable gloves thoroughly with soap and water before removing. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Cleaning of Equipment: When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CONTROLS SUCH WEEDS AS: Black Medic, Chickweed, Clover, Cocklebur, Creeping Charlie, Dandelion, Dock, Florida Pusley, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida, Spurge, Wild Onion, and Wild Lettuce.

TIME TO APPLY:

Apply anytime broadleaf weeds are present, except to avoid treating St. Augustinegrass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or water on weak foliage at time of application aids control. Do not water for 24 to 48 hours AFTER application. Do NOT use on Dichondra or Lippia ground cover or similar covers. Do NOT apply in areas underlaid by roots of desirable trees and shrubs. Do NOT use on newly seeded lawns.

HOW TO APPLY:

Use 20 lbs. on 5,000 sq. ft. for Bahia, Bermuda, Centipede, Fescue and St. Augustine grasses. Overapplication may cause temporary discoloration.

SUGGESTED SPREADER SETTINGS

<u>MAKE OF SPREADER</u>	<u>SPREADER SETTINGS</u>
Kmart	7 - 8
Ortho 3000.	6 - 7
Scotts C-1.	K - L
Scotts 35	6 - 8
Cyclone	4.75 - 5.25
Central	7
Precision	6

NOTE: The above suggested spreader settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application. Close spreader when filling, stopping, or turning. DO NOT HAND SPREAD.

Avoid getting on concrete, stucco or asbestos siding, as the composition of certain fertilizer elements can cause permanent stain.

STORAGE AND DISPOSAL**FOR HOMEOWNER USE ONLY:**

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured By:

PARKER FERTILIZER CO.
201 4TH STREET
SYLACAUGA, AL 35150

Date Printed: 05/08/92

Date Approved:

Supersedes:

Label Change: