P	123	8660		/
US ENVIRONMENTAL PROTECTION AGENCY	EPA REGISTRATION	NO.	01 5 00 1099	3Z
OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767)	DOGU-154		7000	
WASHINGTON, DC 20460	1		: July 31, 1994	
NOTICE OF PESTICIDE: REGISTRATION	NAME OF PESTICIDE			
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	Parker rort:	ılızer +	0.176 Elmension	8
ME AND ADDRESS OF REGISTRANT (Include ZIP code)				
Г	٦			j
Parker rertilizer Co.				1
201 West Fourth Street				
P.O. Box 540				1
Sylacauga, Alabama 35150				
L				
OTE: Changes in labeling formula differing in substance fr	om that accepted in c	onnection wi	h this registration mus	t be
abmitted to and accepted by the Registration Division prior oduct always refer to the above U.S. EPA registration numbers.	to use of the label in per.	commerce. I	n any correspondence o	on this
n the basis of information furnished by the registrant, the a ne Federal Insecticide, Fungicide, and Rodenticide Act.	bove named pesticide	is hereby R	egistered/Reregistered	under
copy of the labeling accepted in connection with this Reg	istration/Reregistrati	on is returne	d herewith.	ĺ
egistration is in no way to be construed as an indorsement calth and the environment, the Administrator, on his motion ide in accordance with the Act. The acceptance of any narct is not to be construed as giving the registrant a right to y others.	, may at any time sus ne in connection with	pend or canc the registrat	el the registration of a ion of a product under	pest- this
ints product is conditionally resection 3(c)(7)(C) provided that you	_	CCULúah C 9	with FIFRA	
l. Add the phrase "EFA Registra	ition 60. d660~	154" to 1	ine labei	i
pefore you release the produ				ŀ
 Submit five (5) copies of the you release the product for 	e final printe		x Letore	
and rounded that woncento Concent	the gangractur	er of to	e active	ł
and provided that monsanto Company, ingredient of this product, submits				
the registration of dithickly as lie	•			1
If these conditions are not comp	liec With, the	renstr	ation will be	
subject to cancellation in accordance				ıse
for snippent of the product constitu]
A stamped copy of the laber is ended	_			
	V/11	/		
4	7. 11111	16.		
	Inne I. Milley	. 537		•
	oduct Hanager (ngicide-Heruici		n	
	gicide-nervici gistration bivi			[
	DECT			
ATTACHMENT IS APPLICABLE	BEST AVA	AILAB	LE COPY	
GNATURE OF APPROVING OFFICIAL			DATE .	
1/1/10/			10/15/92	
Form 8570-6 (Rev. 5-76) PREVIOUS EDITION	MAY BE USED UNTIL	SUPPLY IS E		<u>i</u>
<i>f</i>		_		

(

1110-1000. E1 Engen W. J. 000. 118

Parker Fertilizer + 0.17% Dimension

FOR PROFESSIONAL TURF APPLICATOR USE ONLY

Parker Fertilizer + 0.17% Dimension is a selective herbicide that provides preemergence control of listed annual grasses and broadleaf weeds in established cool and warm-season turfgrasses in lawns and ornamental turf.

EPA Reg. No. 8660-XXX

Dimension is a registered trademark of Monsanto Company.

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION:

CAUSES EYE IRRITATION.

HARMFUL IF INHALED.

Avoid contact with eyes or clothing.

Avoid breathing dust.

Wash thoroughly with soap and water after handling.

Remove contaminated clothing and wash before reuse.

Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

BEST AVAILABLE COPY

CONTRACTORS

15 OCT 1992

Their the Birlinst Indebution.
Only by a load Codecard again.

as and with for the perfect.

= 18

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists.

IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

This pesticide is toxic to fish. Drift or runoff may adversely affect aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

BIRD CAUTION:

While avian toxicology studies have shown low acute toxicity to birds, the potential for avian reproduction effects is being evaluated. To reduce potential exposure to birds, do not apply this product in split applications; make only one application per season; do not apply more than 294 pounds of this product per acre $(0.5\ lb\ ai/A)$. Avoid known bird nesting sites and areas where birds are actively feeding when applying this product.

ACTIVE INGREDIENTS:

*This product contains .085 pound of the active ingredient, dlthlopyr, per

INBRT INGREDIENTS:..... 99.83%

50 pound bag.

Product protected by U.S. Patent No. 4,692,184. Other patents pending.

DIRECTIONS FOR USE					
It is a violation of Federal law to use this product in any manner inconsistent with its labeling.					
For information on the use of this product call 1-800-633-6560 ext 818					
Storage and Disposal					
Do not contaminate water, foodstuffs, feed or seed by storage or disposal.					
STORAGE: Store this product in original container only. In case of spill, sweep up material and dispose of material according to label directions.					
DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.					
Do not reuse emptied container.					
Completely empty bag into application equipment. Dispose of emptied bag(s) in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.					
GENERAL INFORMATION					

This product is a selective herbicide that provides preemergence control of crabgrass and other susceptible annual grasses and broadleat weeds in established lawns and ornamental turf. This product will not harm nearby established ornamentals when used according to label directions:

PRECAUTIONS

For best results, cultural practices that disturb the soil, such as core aerification or verticutting, should be done before application of this product. Turfgrass should be allowed to recover from these cultural practices prior to application of this product. Irrigation or rainfall

501Y

•

within 7 days after application may improve the performance of this product.

Reseeding or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrass. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed to soil contact.

This product may be used on seeded, sodded, or sprigged turfgrass that are well-established. Allow the turf to develop a good root system and uniform stand before application.

Use of this product on golf greens is not recommended.

Use of this product on turfgrass not well-established, weakened by weather, or mechanical related stress may increase the potential for injury.

DO NOT apply over-the-top of desirable flowers, vegetables, strubs or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees.

DO NOT allow domestic animals to feed or forage on the treated turf.

TURFGRASSES

When applied as directed under the conditions described, the following established turgrasses are tolerant to this product:

Cool-Season Grasses

Bentgrass, creeping
Agrostis stolonifera

Bluegrass, Kentucky Poa pratensis

Fescue, fine*
Festuca rubra

Fescue, tall
Festuca arundinacea
Ryegrass, perennial
Lolium perenne

Warm-Season Grasses

Bahiagrass Paspalum notatum Bermudagrass
Cynodon dactylon

Buffalograss
Buchloe dactyloides

Carpetgrass
Axomopus affinis

Centipedegrass

Bremcohloa ophiuroides

Kikuyugrass Pennisetum clandestinum

St. Augustinegrass
Stenotaphrum secundatum

Zoysiagrass
Zoysia japonica

*The following fine fescue varieties have been found to be sensitive to this product: 'Atlants', 'Banner', 'Beauty', 'Bilgart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Koket', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tatjana', 'Walforf' and 'Waldina'.

APPLICATION DIRECTIONS

This product provides preemergence control of listed annual grasses and broadleaf weeds when applied prior to their germination in established turfgrass.

Control of Annual Grasses and/or Annual Broadleaf Weeds

This product should be applied prior to the germination of annual grasses and/or broadleaf weeds. Use the higher recommended use rates where weed infestations are dense and/or applications are made early, and longer residual control is desirable.

For preemergence control of annual bluegrass and listed winter ankidal broadleaf weeds, apply this product in the fall prior to their germination and emergence. Use the higher recommended use rates where weed infestations are dense.

Refer to the following table for recommended use rates of this product for the preemergence control of annual grasses and/or broadleaf weeds.

Weeds Suggested Use Rate (1b/A product) 0.17% a.i.				
	North			
Crabgrass, Large Smooth Foxtail, Green Yellow	74 to 147*	150 to 225*		
Barnyardgrasses Crowfootgrass Kikuyugrass Lespedeza, common Oxalis, Buttercup Oxalis Creeping Woodsorrel Yellow Woodsorrel Purslane, common Speedwell, corn Smutgrass	150	225		
Bluegrass, annual Bittercress Chickweed Geranium, Carolina Goosegass Henbit Parsley-piert Pineappleweed Ryegrass, Annual Perennial Shepherdspurse Spurge Prostrate Spotted		225 to 294*		

 $^{^{\}pm}$ Higher use rates are recommended where weed infestations are heavy and/or dense.

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Calibrate application equipment prior to usage and apply this product uniformly over the turfgrass area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAPEFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

PARKER FERTILIZER CO. P.O. BOX 540 201 W. 4TH STREET. SYLACAUGA, AL 35150