	EPA REGISTRATION NO.	DATE OF ISSUANCE		
US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS	8660-157	DATE OF ISSUANCE		
REGISTRATION DIVISION (7S-767)	TERM OF ISSUANCE			
WASHINGTON, DC 20460	Conditional, Expires: 1	December 31, 1997		
NOTICE OF PESTICIDE: REGISTRATION	E OF DESTICIDE REGISTRATION NAME OF PESTICIDE PRODUCT			
	Dimension [®] 270-G A Granule Preemergence			
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	Turf Herbicide			
ME AND ADDRESS OF REGISTRANT (Include ZIP code)	······································			
г.	L_			
Purceli Industries, Inc.				
F.C. EOX 540 Sylacausa, AL 35150				
Syrucausa, nr. 55190				
OTE: Changes in labeling formula differing in substance ubmitted to and accepted by the Registration Division pri- roduct always refer to the above U.S. EPA registration num	or to use of the label in commerce, mber.	In any correspondence on this		
n the basis of information furnished by the registrant, the ne Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is hereby F	egistered/Reregistered under		
copy of the labeling accepted in connection with this Re	egistration/Reregistration is returne	d herewith.		
egistration is in no way to be construed as an indorsemer ealth and the environment, the Administrator, on his motion of a necordance with the Act. The acceptance of any n act is not to be construed as giving the registrant a right to y others.	on, may at any time suspend or can ame in connection with the registra to exclusive use of the name or to i	cel the registration of a pest- tion of a product under this ts use if it has been covered		
Enis (court to conditionally t section (C)(7)(c) provided that yes		and Flee		
 Add the phrase "anA hogistra percre you release the produce Add: "In the State of New You what,", directly relievant, Secondaries the Anarodout "BO No bestgrass (Aurostis famuis) is this paragraph because it 	of for shipment. The violation statement. We apply this product to Varieties." from the other	oury and Suiloik Colonau E statementa		
Cuntinued on Page 2 of this boti	<u>.C</u> e			
Pro Enclosure rur	anne I. Hiller oduct Manager (23) ogicide-merbicide Branch gistration Division (75050)		
Pro Enclosure rur	oduct Manager (23) ngicide-rerbicide granch	2)		
Pro Enclosure rur	oduct Manager (23) ngicide-rerbicide granch	DATE OCT - 3 1995		

Page 2 of Notice of Registration

294

Dimension[®] 270-G A Granule Preemergence Turf Herbicide

EPA Registration No. 8660-157

- 4. Correct the claim "50 lb. Bag treats up to 21,750 Sq.Ft. for Crabgrass Control" to read: "50 lb. Bag treats up to 21,780 Sq.Ft. for Crabgrass Control".
- 5. Submit a revised confidential statement of formula to reflect correct upper and lower limits of the active ingredient in this product. Compliance with PR Notice 92-1 is required.
- 6. Submit five (5) copies of the final printed label before you release the product for shipment.

and provided that Rohm and Hass Company, the manufacturer of the active ingredient of this product, submits <u>all</u> the required data to continue the registration of dithiopyr as listed in the EPA Fact Sheet No. 223.


If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Nour release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Dimension® 270-G is a selective herbicide that provides preemergence control of listed annual grasses, such as crabgrass, and broadleaf weeds in established cool and warm-season turfgrass in lawns and ornamental turf. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. ACTIVE INGREDIENT:

Dithlopyr, 3,5-pyridine-ticarbothioic acid, 2-(difluoromethyl)-4-(2-methyloropyl)-8-(trifluoromethyl)- S,S-dimethyl ester INERT INGREDIENTS: 0.27%

99.73% Total 100.00%

*This product contains 0.135 pounds of the active ingredient, dithiopyr, per 50 pound bag. Product protected by U.S. Patent No, 4,692,184. Other patents pending. EPA REG NO. 8660-_____EPA Est. No. 8660-AL-1

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.



50 lb, Bag Treats up to 21, 750 Sq.Ft.

for Crabgrass Control

PRECAUTIONS Use of this product on golf greens is not recommended. DO NOT apply this prod-uct to Colonial bentgrass. For best results, cultural practices that disturb the soil, h as core, spike, hyro- serification, or verticutting, and top dressing should be e before application of this product. This product should not be applied until grasses have recovered from stresses related to these or other mechanical operations. Irrigation or rainfall within 7 days after application will improve the per-formance of this product.

Reseeding or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turgrass. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment ((silf seeders) designed to give good seed-to-soil contact. This product may be used on seeded, sodded, or sprigged turgrass that are well-established. Allow the turf to develop a good root system and uniform stand before application. uniform stand before application

Use of this product on turfgrass not well-established, weakened by lack of mois-ture, chemical injury, insects, disease, or temperature extremes that cause stress will increase the potential for injury. Allow turfgrass weakened by these stresses to recover before application of this product, DO NOT apply to flowers, vegetables, shrubs or trees. DO NOT use clippings from treated turf for mulching around veg-etables or fruit trees. DO NOT allow domestic animals to feed or forage on the treated turf. treated turf

This product can be applied only once per year at a maximum use rate of 185 pounds per acre (one-half pound active ingredient per acre). DO NOT make re-beat or split applications of this product during a single growing season.

TURFGRASS

When applied as directed under the conditions described, the following established turigrasses are tolerant to this product: _____

Cool-Season Grasses

Bentgrass, creeping* Bluegrass, Kentucky	Agrostis stolonifera – Pos prateosis
Fescue, fine** Fescue, tall	Festuce rubm
Rys grass, perenniat	Festuce arundinacea Lollum parenne

Warm-Season Grasses

Paspalum notatum
Cynodon dactylon
Buchice dactyloides
Axonopus affinis
Eremochica ophiuroides
Pennisetum clandestinum
Stenotaphrum secundatum
Zoysia japonica

* DO NOT apply this product to Colonial bentgrass. Use of this product on certain varieties of creeping bentgrass, such as "Carmen", "Cohansey", or "Seaside" will result in undestrable turf injury. Not all varieties of creeping bentgrass have been

It is a violation of Federal law to use this product in any manner inconsistent with its theling
GENERAL INFORMATION
This product is a selective herbicide that provides preemergence control of rabin grasses and other susceptible annual grasses and broadleaf weeds in established inavns and ornamental tur? This product will not harm nearby established ornamentals when used according to label directions.
PRECAUTIONS
Use of this product on golf greens is not recommended. DO NOT apply this product provides preemergence control of a science, spike, hyro, serification, or verticuting, and top dressing should be abole applied until has core, spike, hyro, serification, or verticuting, and top dressing should be applied until.
AppLICATION EQUIPMENT AND INSTRUCTIONS
Apply this product with drop or rotary-type spreaders designed to apply granular the there is the previous and then applying the remaining hait at a selection application. Check equipment frequently to ensure equipment is functioning property and applying uniform distribution of granules.
AppLiCATION EQUIPMENT AND INSTRUCTIONS
Apply this product with drop or rotary-type spreaders designed to apply granular the trequired amount over the area and then applying the remaining hait at a selections.
PRECAUTIONS
Use of this product on golf greens is not recommended. DO NOT apply this product provides preemergence control of fisted annual grasses and broadleaf weeds when applied prior to their germination in established furgrass.
This product provides preemergence control of this product. This product should not be applied until

Page 5 of 18

weeds when applied prior to their germination in established turigrass. Control of Annual Grasses and/or Annual Broadleaf Weeds This product should be applied prior to the germination of annual grasses and/or broadleaf weeds. Use the higher recommended rates where weed infestations are dense and/or applications are made early and longer residual control is desirable. For premergence control and/or suppression of annual bluegrass and listed whiter annual broadleaf weeds, apply this product prior to their germination and emergence Use the higher recommended use rates where weed infestations are dense.

Refer to the following table for recommended use rates of this product for the preemergence control of annual grasses and/or broadleaf weeds.

Weeds Controlled and Use Rate Recommendations

<u>*Use 100 to 150 Pounds/Acre = 2,3 to 3,4 Pounds/1000 sq. ft.</u> FOR: Crabgrase; (Large, Smooth, Southein) Fostall; (Green, Yellow) FOR:

"Use 150 to 185 Pounds/Acre = 3.4 to 4.2 Pounds/"000 sq. ft.

Dichondra, Carolina Yellow Woodsorrei Speurge, Pineappieweed Prusitare Spottod	OR:	Barnyardgrass Chickweed Clover, hop Crabgrass, tropical Crowfoolgrass Dichondra, Carolina	Kikuyugrasa Oxalis, Buttercup Creeping Woodsorrei Yellow Woodsorrei	Spurge, Prostate
--	-----	--	---	---------------------

"Use 185 Pounds/Acre = 4,2 Pounds/1000 33. ft.

OR:	Bluegrass, annual Bittercress Geranium, Carolina Chickweed Goosegrass	Henbit Parsley-plert Pinasppleweed Purslane, common Sheperdspurse
-----	---	---

*The higher use rates are recommended where weed infestations are heavy and/or dense. **Use of this product at this rate will provide only partial control and/or sup-pression of listed weeds.

Spreader Settings						
Pounds / 1.000 so.ft.	1.15 be		ation	236	71HE ADDE	4 2 lbs
Cyclone B-1 Scott Drop	xx	XX	xx	×x .	XX.	XX
Scoll Drop	<u>xx</u>	XX	XX	1 101		XX

CONDITIONS OF SALE AND WARRANTY Pursell industries, he, warrants that the product conforms to fits chemical description and its reasonably fit for the purpose stated on the labet only when used in accordance with tabul directions under normal conditions of use, Pursell industries, inc. NAXC3 NO OTIETEX PRESS OF MPLIED WARRANTES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are be cond the control of Pursell industries, inc. and Seller. Risks such as crop injuly, indifficitive, no.; a c other unintended consequences resulting from, but not fimited to, vegether us soft conditions, presence of other malerists, disease, peal, drill to niner crops or property or failures to follow is bed directions will be assumed by the Buyer or User NNO CASE WILL PURSELL # INTY TRIES, INC, OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDI-RECT DAMAGES RESULTING FROM THE HANDLING, STORAGE CA, USE OF THIS PROD-UCT. (awardg.endors 17 55) UCT. (dmt270g.)

