

PM23

8660-160

7/23/99

1 of 14

JUL 23 1999

Mr. Sanford K. Simon
Director, Technical Services
Pursell Industries, Inc.
P.O. Box 540
Sylacauga, AL 35150

Dear Mr. Simon:

subject: Parker Turf Fertilizer plus Dimension® 140 Crabgrass Preventer
(Alternate Brand Name "Crab-Ex Crabgrass Preventer and Fertilizer")
EPA Registration No. 8660-160
Application Dated May 24, 1999, Request To
Amend the Labeling: To Use "First Aid" instead
of "Statements of Practical Treatment", Added a Toll-free
Telephone Number for Product Information, Use of Common
Name of Chemical Rather Than Chemical Name, Addition of
Chemical Abstracts Number for Chemical, Use of "Other"
Instead of "Inert" in the Ingredient Statement, Added
a Description of Product Formulation, Placing Ingredient
Statement on Back Panel, Added the Primary Hazard Directly
Under the Signal Word, Use of New Format To Communicate
Consumer Information, Storage and Disposal Statement
To Discourage Disposal of Unused Product, Revised
Application Rates That Do Not Exceed 0.22 lbs. A.I.
per Acre, and To Add List of Advertising Claims

The proposed labeling to amend the subject registration
has been reviewed and is acceptable under the Federal Insecticide,
fungicide and Rodenticide Act (FIFRA) as amended, provided
that you:

1. Delete your document identification information where it occurs in the proposed labeling.
2. Add the EPA Worker Protection Standard exclusionary statement as required for this product (EPA PR Notice 93-11) and that reads: "not for use on plants being grown for sale or other commercial use, or for seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or lawns and grounds, in a residential setting".
3. The instructions indicate that 3 different rates may be used. Revise the instructions for each use-pattern to tell the using public when to use each of these rates, or list only one rate under the "How to Use" section.

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

2 of 14

-2-

- 4. Submit one (1) copy of the final printed label before releasing the product for shipment bearing the subject labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Joanne I. Miller
 Product Manager (23)
 Herbicide Branch
 Registration Division (7505C)

Enclosure

E.Wilson:Diskette:Dithiopyr:07-23-99

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

3014

FRONT PANEL

Crab-Ex™

Crabgrass Preventer and Fertilizer

For Outdoor Residential Use
Only

Pre and Early Post Emergence Control of Crabgrass

- A selective herbicide plus Fertilizer that provides control of crabgrass and other weeds on established lawns
- Contains Polyon® timed-release nitrogen for a Beautiful, Green, Vibrant Lawn
- Contains the active ingredient found in Dimension® Turf Herbicide

XX - XX - XX

**STOP - Read This Entire Label Before Use
KEEP OUT OF REACH OF CHILDREN**

CAUTION

Causes Eye Irritation, If in Eyes

See back panel for additional precautionary statements

Net Contents: 16, 18 or 20 lbs.


TREATS [or COVERS] 5,000 SQ FT.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JUL 23 1999

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

8660-160



BACK PANEL

Ingredients

ACTIVE INGREDIENT:	
Dithiopyr*	0.14%
OTHER INGREDIENTS**	99.86%
TOTAL	100.00%

*CAS # 97886-45-8

** Other ingredients in Crab-Ex are Premium Sta-Green Fertilizer, including Polyon® slow-release granules and dipropylene glycol to adhere the active ingredient on the fertilizer.

EPA Reg. No. 8660-160	EPA Est. No. 8660-AL-1	8660-OH-1
First letter of run code on bag matches letter above EPA Est. No.		

GUARANTEED ANALYSIS

Total Nitrogen (N)	XX.00%
X.X% Ammoniacal Nitrogen	
X.X% Urea Nitrogen*	
Available Phosphate (P ₂ O ₅)	XX.00%
Soluble Potash(K ₂ O)	XX.00%

Derived From: Polymer-coated urea, urea, diammonium phosphate, and muriate of potash

*X.X% Slowly Available Urea Nitrogen From Polymer-coated urea.

F1061

PRECAUTIONARY STATEMENTS
Hazards to Humans and
Domestic Animals

CAUTION:

- Causes eye irritation, if in eyes.
- Harmful if inhaled.
- Avoid breathing dust.
- Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling.
- If product gets on clothes, remove contaminated cloth and wash before reuse.
- Prolonged or repeated skin contact may cause allergic skin reactions in some individuals.

First Aid

The table below describes the first aid procedures for incidents involving CrabEx.

In the case of accidental...	Immediately...
Contact with eyes	<ul style="list-style-type: none"> • Open eyelids and rinse eye slowly and gently with water. • Continue to rinse eye for 10-15 minutes. • Call a doctor if irritation persists.
Inhalation of dust	<ul style="list-style-type: none"> • Move person to fresh air • Call a doctor or poison control center if breathing difficulty develops.
<p>NOTE: When calling a poison control center, have Crab-Ex label accessible. If advised to seek treatment at an emergency room or doctor's office, bring this product label to show medical personnel.</p> <p>Active Ingredient: Dithiopyr, CAS # 97886-45-8</p>	

Environmental Hazards

- This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas.
- Do not contaminate water when cleaning equipment or disposing of equipment washwater.

Directions for Use
IMPORTANT: Follow label directions

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For outdoor residential use only.

Using these directions

These "Directions for Use" are designed to help you determine the appropriate application procedure for Crab-Ex.

Before you Use

Before using Crab-Ex, make sure that you read the entire "Directions for Use" section of this label.

Read and follow these guidelines when you use Crab-Ex:

- Apply only to lawns and ornamental turf that are well established.
- Wait at least 12 weeks after using Crab-Ex before reseeding, overseeding or sprigging of grass into areas that have been treated.
- Use of product on new lawns or grass that has been weakened or stressed by drought, excessive temperatures or pests, increases chance of injury to grass.
- When applying around ponds or other bodies of water, do not apply in windy conditions or where product can run-off into water.
- Keep people and pets off treated areas until dust has settled.
- Do not use grass clippings as mulch around vegetables or fruit trees.
- Crab-Ex™ Crabgrass Preventer and Fertilizer may be applied to flowers listed on this label.
- Do not make more than 1 application to the same area in the same year.

How to use

[Only one application rate will be given on each end use package. The rate in pounds of product per 1,000 sq. ft. is shown below for a package weight of either a 16, 18, or 20 pound container.]

1. Apply Crab-Ex at the rate of:
 - 3.2 lbs. per 1,000 sq. ft. for 16 lb. bag to treat 5,000 sq. ft., or
 - 3.6 lbs. per 1,000 sq. ft. for 18 lb. bag to treat 5,000 sq. ft., or
 - 4.0 lbs. per 1,000 sq. ft. for 20 lb. bag to treat 5,000 sq. ft.
2. Calibrate your spreader according to the spreader manufacturer's directions. Adjust settings as needed to apply the recommended rate of product.
3. Using a drop-type or rotary spreader, apply granules evenly over grass, avoiding streaking, skips or overlaps. Avoid using spreaders that apply product in narrow or concentrated bands.
4. Check calibration of spreader frequently to insure spreader is functioning properly and applying product evenly.

Spreader Settings

This 16, 18, or 20 pound bag covers 5,000 sq. ft. Use the settings below to apply at a rate of [(3.2 pounds for 16 pound bag), or (3.6 pounds for 18 pound bag), or (4.0 pounds for 20 pound bag)] lbs. per 1,000 sq. ft.

Rotary Spreaders	Drop Spreaders
X	X
X	X
X	X
X	X
X	X

Where to use

This product can be used on the following types of grass when applied as directed:

Cool Season Grasses	Warm Season Grasses
Agrostis Stolonifera	Axonopus affinis
/ Bentgrass, creeping	/ Bahia
/ Bluegrass, Kentucky	/ Bermuda
/ Fescue fine	Buchloe dactyloides
/ Fescue tall	/ Buffalograss
Festuca arundinacea	/ Carpetgrass
Festuca rubra	/ Centipede
Lolium, perenne	Cynodon dactylon
Poa pratensis	Eremochloa ophiuroide
/ Rye grass, perennial	Kikuyugrass
	Paspalum notatum
	Pennisetum clandestinum
	St. Augustine
	Stenotaphrum secundatum
	Zoysia japonica
	/ Zoysiagrass

When to apply

- Product may be applied either in the Spring or Fall for the control of weeds.
- Product will provide up to 4 months of weed control when used as directed, although length of control can be affected by weather, grass conditions and weed pressures.
- To control listed weeds other than crabgrass, applications must be made before weeds emerge.
- Crab-Ex™ will control crabgrass if applied before it emerges and up to 4 weeks after it emerges.

Weeds Controlled

Pre- and Early Post Emergent

- Crabgrass

Pre-Emergent (Preventative)

Grass Weeds	Broadleaf Weeds
Annual Bluegrass	Bittercress
Annual Ryegrass	Buttercup Oxalis
Barnyardgrass	Carolina Geranium
Crabgrass	Chickweed
Crowfootgrass	Common Lespedeza
Goosegrass	Creeping Oxalis
Green Foxtail	Henbit
Kikuyugrass (seedling)	Parsley-piert
Perennial Ryegrass	Pineappleweed
Smutgrass (seedling)	Prostrate Spurge
Yellow Foxtail	Spotted Spurge
	Yellow Woodsorrel
	Corn Speedwell

TOLERANT TREES, SHRUBS, AND FLOWERS**

When applied as directed under the conditions on this label, trees, shrubs, and flowers listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Follow directions given above to determine plant tolerance under your growing conditions. A limited area involving only a few plants should be treated and observed for signs of yellowing, wilting, or lack of growth prior to large-scale application.

Common Name	Botanical Name	Tolerant Cultivars
Abella, Dwarf	Abella X grandiflora	Nana
Ajuga	Ajuga reptans	Bronze
	Ajuga genevensis	Bronze Beauty
Almond, Flowering	Prunus glandulosa	
Apple*	Malus pumila	
Arborvitae	Thuja occidentalis	Nigra; Pyramidalis; Smaragd Techny; Woodwardii.

* Ornamental Species only. DO NOT USE ON FOOD-PRODUCING TREES AND PLANTS.

**Full list of Tolerant Trees, Shrubs, and Flowers listed as an addendum to this label.

Storage and Disposal

Take the following precautions:

IMPORTANT: Precautions during storage and disposal

- Keep out of reach of children
- Do not reuse empty product bags.

Storage conditions

Store Crab-Ex® in a cool and dry location. The storage area should be locked and inaccessible to children.

Disposal

Securely wrap the empty product bag in several layers of newspaper, and place bag in trash.

Important: If all of product can not be used according to label directions, make sure bag is securely sealed or wrapped up before disposing.

Conditions of Sale and Warranty

NOTICE: Buyer and user assumes all risk and liability of use, storage, and/or handling of this product not in accordance with the terms of this label.

Product Information: 800-874-8892

Manufactured By:
Pursell Industries, Inc.
P.O. Box 540
Sylacauga, AL 35150

STA-Green® and CRAB-EX® are registered trademarks of Pursell Industries, Inc. POLYON® is a registered trademark of RLC Technologies, L.L.C. used by permission. The POLYON® design and the GREEN color of the POLYON® capsule are trademarks of RLC Technologies, L.L.C., used by permission. POLYON® Patent Nos. 4,711,659; 4,804,403; 4,969,947; 5,374,292. Dimension® is a registered trademark of Rohm & Haas Company.

THE STA-GREEN® PROMISE - YOUR SATISFACTION GUARANTEED OR YOUR MONEY BACK

If you apply STA-GREEN® CRAB-EX® plus Lawn Fertilizer with POLYON® timed-release nitrogen as directed and are not completely satisfied, Just send a proof of purchase from this package and a cash register receipt to the address below. We will promptly refund your money.

ADDENDUM**CrabEx Crabgrass Preventer and Fertilizer
EPA Reg. No. 8660-160****TOLERANT TREES, SHRUBS, AND FLOWERS**

When applied as directed under the conditions on this label, trees, shrubs, and flowers listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Follow directions given above to determine plant tolerance under your growing conditions. A limited area involving only a few plants should be treated and observed for signs of yellowing, wilting, or lack of growth prior to large-scale application.

COMMON NAME	BOTANICAL NAME	TOLERANT CULTIVARS
Abelia, Dwarf	<i>Abelia X grandiflora</i>	Nana
Ajuga	<i>Ajuga reptans</i> <i>Ajuga genevensis</i>	Bronze Bronze Beauty
Almond, Flowering	<i>Prunus glandulosa</i>	
Apple*	<i>Malus pumila</i>	
Arborvitae	<i>Thuja occidentalis</i>	Nigra Pyramidalis Smaragh Techny Woodwardii
Arborvitae, Golden	<i>Thuja orientalis</i>	
Aster, Chinese	<i>Callistephus chinensis</i>	Dwarf Queen
Ash, Green	<i>Fraxinus pennsylvanica</i>	
Ash, Mountain	<i>Sorbus aucuparia</i>	
Ash, Purple	<i>Fraxinus americana</i>	
Azalea	<i>Rhododendron</i> spp.	Brilliant Buccaneer Carror Chimes (Belgian) Exbury Fashion Hardijzer Beauty Hershey Red Higasa Hinocrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm
Azalea, Flame	<i>Rhododendron calendulaceum</i>	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	<i>Berberis thunbergii</i>	Aurea Dwarf Pigmy Green Kobold Pygmy Red Rose Glow
Barberry, Purple		Atropurpurea
Bearberry (common)	<i>Arctostaphylos uva-ursi</i>	Massachusetts
Birch, River	<i>Betula nigra</i>	
Blueberry*	<i>Vaccinium</i> spp.	Bluecrop

		Blue Jay Jersey North Blue Northland
Bottlebrush	<i>Callistemon citrinus</i>	
Boxwood, Japanese		Japonica
Boxwood, Weller	<i>Buxus sempervirens</i>	
Broom	<i>Cytisus spp.</i>	Moonlight
Broom	<i>Genista pilosa</i>	Vancouver Gold
Bugle Carpet		
Camellia	<i>Camellia japonica</i> <i>Camellia sasangua</i>	Debutante Mathotiana Supreme Chansonette
Carex, Variegated	<i>Carex</i>	
Cedar, Red	<i>Juniperus virginiana</i>	
Cockscomb, Plumosa	<i>Celosia cristata</i>	Red Kewpie
Coleus	<i>Coleus blumei</i>	
Copper leaf	<i>Acalypha wilkesiana</i>	
Cotoneaster	<i>Cotoneaster apiculatus</i>	
Coyotebrush	<i>Baccharis petularis</i>	
Cycads	<i>Cycads revoluta</i>	
Cypress, Bald	<i>Taxodium distichum</i>	
Cypress, Italian	<i>Cupressus sempervirens</i>	Glaucia
Cypress, Japanese False	<i>Chamaecyparis obtusa</i>	Gracilis
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	
Daffodil	<i>Narcissus spp.</i>	King Alfred
Daylily	<i>Hemerocallis spp.</i>	Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilson's Yellow
Dogwood	<i>Cornus florida</i>	
Dogwood, American	<i>Cornus sericea</i>	Flavarimaea
Douglas Fir	<i>Pseudotsuga menziesii</i>	
Dusty Miller	<i>Senecio cineraria</i>	
Elm, Drake	<i>Ulmus parvifolia</i>	
Euonymus	<i>Euonymus fortunei</i>	Argenteo-variegata Auereo-marginata Colorata Emerald Gaiety Emerald 'N Gold Gold Edge Gold Princess Silver King Tricolor Vegetus
Fan Palm, European	<i>Chamaerops humilis</i>	
Fan Palm, Mexican	<i>Washingtonia robusta</i>	
Fescue	<i>Festuca glauca</i>	
Fetterbush	<i>Leucothoe fontiana</i>	Rainbow
Ficus	<i>Ficus retusa</i>	Nitidia
Fir Fraser	<i>Abies fraseri</i>	
Forsythia	<i>Forsythia X intermedia</i> <i>Spring Glory</i>	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark Weeping
Gardenia	<i>Gardenia jasminoides</i>	Mystery

		Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf San Jose Sargent Blue Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan
	<i>Juniperus chinensis</i>	Tamariscifolia
	<i>Juniperus sabina</i>	Weberi Youngstown Yukon Belle
King Palm	<i>Archontophoenix cunninghamiana</i>	
Laurel, Australian	<i>Pittosporum tobira</i>	
Laurel, Mountain	<i>Kalmia latifolia</i>	
Leucothoe	<i>Leucothoe fontanesiana</i>	
Ligustrum, Japanese	<i>Ligustrum japonicum</i>	
Lily, Africian	<i>Agapanthus africanus</i>	Albus Peter Pan
Lily, Africian Blue		
Lily of the Valley	<i>Pieris japonica</i>	Mt. Fire
Lilyturf	<i>Liriope muscari</i>	Evergreen Giant Lilac Beauty Majestic Monroe White Variegata
Liriope, Green	<i>Liriope spicata</i>	
Maple, Japanese	<i>Acer japonicum</i>	
Maple, Norway	<i>Acer platanoides</i>	
Maple, Red	<i>Acer rubrum</i>	
Maple, Silver	<i>Acer saccharinum</i>	
Maple Sugar	<i>Acer saccharum</i>	
Marigold	<i>Tagetes patula</i>	Honeycomb Variegata Wheeler's Dwarf
Mock Orange*	<i>Philadelphus spp</i>	Golden Snowflake Double White
Mopndo Grass	<i>Ophiopogon japonicus</i>	
Moss Rose	<i>Portulaca grandiflora</i>	Sunnyside
Myrtle, Crape	<i>Lagerstoemia indica</i>	Faurei Langer Muskogee Standard Pink
Myrtle, Wax	<i>Myrica cerifer</i>	
Nadina	<i>Nadina domestica</i>	Compacta Nana
Oak, Laurel	<i>Quercus laurifolia</i>	
Oak, Pin	<i>Quercus palustris</i>	
Oak, Red	<i>Quercus rubra</i>	
Oak, Southern L	<i>Quercus virginiana</i>	
Oak, Willow	<i>Quercus phellos</i>	
Oleander	<i>Nerium oleander</i>	Hardy Red

		Petite Pink Sister Agnes
Oteospermum	<i>Osteospermum fruticosum</i>	Wirligig
Pachysandra	<i>Pachysandra terminalis</i>	
Palm, Bangalow		
Pampas Grass	<i>Cortadaria selloana</i>	
Paper Flower	<i>Bougainvillea glabra</i>	Barbara Karst
Peach*	<i>Prunus persica</i>	
Periwinkle, Dwarf	<i>Vinca minor</i>	
Petunia	<i>Petunia X hybrida</i>	Picoti
Photinia, Red Tip	<i>Photinia X fraseri</i>	
Pieris	<i>Pieris japonica</i>	
Pine, Australian	<i>Pinus nigra</i>	
Pine, Japanese Black	<i>Pinus thumbergiana</i>	
Pine, Loblolly	<i>Pinus teada</i>	
Pine, Longleaf	<i>Pinus palustris</i>	
Pine, Mugho	<i>Pinus mugho</i>	
Pine, Scotch	<i>Pinus sylvestris</i>	
Pine, Slash	<i>Pinus elliottii</i>	
Pine, Swiss Mt.	<i>Pinus mugo</i>	
Pine, Virginia	<i>Pinus virginiana</i>	
Pine, White	<i>Pinus strobus</i>	
Pineapple, Guava	<i>Feijoa fellowaiana</i>	
Pittosporum, Japan		
Pontentilla	<i>Pontentilla fruticosa</i>	Abbotswood
Privet	<i>Ligustrum japonicum</i>	Golden Vicary Regal Texanum Wax Yellow Tipped
Privet, Glossy	<i>Ligustrum lucidum</i>	
Pyracantha	<i>Pyracantha koidzumii</i>	Gnome Lalandei Victory
Queen Palm	<i>Arecastrum rammanzoffianum</i>	
Quince, Japanese*		
Rhododendron	<i>Rhododendron spp</i>	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	<i>Rhododendron carolinianum</i>	
Rhododendron, Catawba	<i>Rhododendron catawbiense</i>	
Rodie Max (Rosebay)	<i>Rhododendron maximum</i>	
Rose	<i>Rosa banksiaae</i>	Luta
Rosemary*	<i>Rosmarinus officinalis</i>	
Rosemary, Bog	<i>Andromeda polifolia</i>	Nana
Salvia	<i>Salvia farinacea</i>	Rhea
Sedum	<i>Sedum spurium</i>	Dragon Blood Red Red Carpet Yellow
Sourwood	<i>Oxydendrum arboreum</i>	
Spiraea	<i>Astilbe X arendsii</i>	Fanall

Spiraea	<i>Spiraea spp.</i>	Anthony Waterer Red Dolchica Froebeli Pink Goldenflame Red Snowmound White Van Houtte White
Spiraea, Garland	<i>Spiraea spp.</i>	
Spruce, Black Hills	<i>Picea glauca</i>	
Spruce, Colorado Blue	<i>Picea pungens</i>	
Spruce, Norway	<i>Picea abies</i>	
Spruce, White	<i>Picea glauca</i>	Conica
Sweetflag, Grassyy-Le		
Sweetgum	<i>Liquidambar styraciflua</i>	
Sycamore	<i>Platanus occidentalis</i>	
Tree Fern (Tiki Fern)	<i>Asparagus virgatus</i>	
Trumpet Flower, Evening	<i>Gelsemium sempervirens</i>	
Tulip	<i>Tulip, spp</i>	Apeldoorn
Verbana, Shrub	<i>Lantana sellowiana</i>	
Vervain	<i>Verbena spp.</i>	St. Paul
Viburnum	<i>Viburnum spp.</i>	American Cranberry Bush Arrowood Common Snowball European Cranberry Bush Linden Mohican Wright
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yaupon	<i>Ilex vomitoria</i>	Dwarf
Yew	<i>Taxus cuspidatra</i> <i>Taxus X media</i>	Denisiformis

***Ornamental species only. DO NOT USE ON FOOD PRODUCING TREES, SHRUBS, AND FLOWERS.**

AD CLAIMS

[May be used on some product labels.]

1. Contains POLYON® Timed-Release Fertilizer, used by professional turf managers.
2. It's [POLYON] the green inside.
3. Time-Release Fertilizer – POLYON® – If it's not green ...it's not POLYON™
4. Contains TriKote® Timed-Release Fertilizer.
5. Pre and early Post Emergence Control of Crabgrass
6. Covers XXXX sq. ft.
7. Money-Back Guarantee.
8. Look for other quality STA-GREEN® lawn and garden products. For more information about POLYON® and STA-GREEN products visit us on the Internet at www.fertilizer.com or call our Consumer Service Hotline at 1-800-874-8892.
9. Never apply by hand.
10. Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 sq. ft. per growing season. When used as directed, this product provides x.xx lbs. of nitrogen per 1,000 sq. ft. Consult your local County Extension Service for questions regarding a regular fertilizer program suitable for your geographical region.
11. Will not harm listed ornamentals or shrubs.
12. Non-Staining
13. Provides up to 4 months control of crabgrass
14. Use Before or After germination of crabgrass for preemergent and early postemergent control
15. Controls Crabgrass for up to 120 days
16. Also controls other grasses and broadleaf weeds such as annual bluegrass, goosegrass, yellow foxtail, spotted spurge, chickweed, oxalis, henbit and other weeds listed on this label.
17. Feeds for up to 2 months at 72° F.
18. If you apply Crab-Ex™ crabgrass preventer plus lawn fertilizer with POLYON timed-release nitrogen as directed and are not completely satisfied, send proof of purchase from this package & the cash register receipt to the address below. We will promptly refund your money.
19. Sta-Green® products with POLYON timed-release nitrogen work differently from other slow-release products. POLYON timed-release nitrogen releases in response to soil temperature rather than moisture level. Other slow release products release according to the amount of moisture present in the soil...the more moisture the quicker the nutrient release...putting your grass at risk from burning. Because Sta-Green® products contain POLYON timed-release nitrogen, your grass is fed for up to 2 months at 72° F
20. IMPORTANT – It is important to follow directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to calibrate spreader before beginning, adjusting if necessary. Consult Owner's Manual.
21. NEVER APPLY BY HAND
22. [Used when staining fertilizer nutrients such as ferrous sulfate are used in formula] – This product may [or will] stain sidewalks and driveways. Sweep [vacuum] off any product which may have drifted onto these areas.
23. What the numbers mean:

XX	XX	XX
Nitrogen (N)	Phosphorous (P)	Potassium (K)
Contains Timed-Release Nitrogen for growing a thick green lawn	Improves Root Growth	Improves Plant Hardiness & Drought Resistance