

Mr. Sanford R. Simon
Director, Technical Services
Pursell Industries, Inc.
P.O. Box 540
Sylacauga, AL 35150

JAN - 5 2000

Dear Mr. Simon:

Subject: Dimension® 6-25G
Alternate Brand Names "Vigoro" Crabgrass Preventer and
STA-GREEN® CRAB-EX x® Crabgrass Preventer)
EPA Registration No. 8660-167
Application Dated December 6, 1999, Request To
Amend the Labeling: To Allow Split Applications for
Use in Weed Control in Southern Lawns

The proposed amendment of the registration of the subject pesticide product has been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) as amended, provided that you:

1. Delete your document identification information where it occurs in the proposed labeling.
2. Submit one label and respective Confidential Statement of Formula that reflects each percent active ingredient (a.i.) between 0.06% and 0.25% a.i. for each product shipped under EPA registration no. 8660-167.
3. Submit one (1) copy of the final printed label before releasing the product for shipment bearing the subject labeling.
4. If a range of dosages is given in the directions of use section of any labeling submitted under this range registration, you must tell the using public when to use the various possible dosages.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

CONCURRENCES Miller

SYMBOL				Product Manager (25)			
SURNAME				Herbicide Branch			
DATE				Registration Division (7505C)			

enclosure

FRONT PANEL
Homeowner use label

Dimension® 6-25G

Granular Crabgrass Preventer for Home Lawns

For Outdoor Residential Use Only

Pre and Early Post Emergence Control of Crabgrass

- A selective herbicide that provides control of crabgrass and other weeds on established lawns and ornamental landscapes.
- Provides early post emergence control of crabgrass when applied within four weeks of germination. [Claim for rates 0.22 – 0.25 pounds of AI per acre]
- Contains the active ingredient found in Dimension® Turf Herbicide

**STOP - Read This Entire Label Before Use
KEEP OUT OF REACH OF CHILDREN**

CAUTION

Causes Eye Irritation, If in Eyes

See back panel for additional precautionary statements

Ingredients

ACTIVE INGREDIENT:	
Dithiopyr*	0.06% - 0.250%
OTHER INGREDIENTS**	99.94% - 99.750%
TOTAL	100.00% 100.000%

*CAS # 97886-45-8

** Other ingredients in this product are clay granules used as a carrier and dipropylene glycol to adhere the active ingredient to the clay.

	A	H
EPA Reg. No. 8660-167	EPA Est. No. 8660-AL-1	8660-OH-1
First letter of run code on bag matches letter above EPA Est. No.		

Net Contents: XX.X lbs. TREATS [or COVERS] XXXX sq. ft.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

JAN - 5 2000

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

8660-167

BACK PANEL

[Product to be formulated such that application rate will not exceed 0.25 pounds of AI per acre]

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION:

- Causes eye irritation, if in eyes.
- Harmful if inhaled.
- Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling.
- If product gets on clothes, wash before reuse.
- Prolonged or repeated skin contact may cause allergic skin reactions in some individuals.

First Aid

The table below describes the first aid procedures for incidents involving Dimension® 6-25 G.

In the case of accidental...	Immediately...
Contact with eyes	<ul style="list-style-type: none"> • Open eyelids and rinse eye slowly and gently with water. • Continue to rinse eye for 10-15 minutes. • Call a doctor if irritation persists.
Inhalation of dust	<ul style="list-style-type: none"> • Move person to fresh air • Call a doctor or poison control center if breathing difficulty develops.
<p>NOTE: When calling a poison control center, have this product label accessible. If advised to seek treatment at an emergency room or doctor's office, bring this product label to show medical personal.</p> <p>Active Ingredient: Dithiopyr, CAS # 97886-45-8</p>	

Environmental Hazards

- This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas.
- Do not contaminate water when cleaning equipment or disposing of equipment washwater.

Directions for Use
IMPORTANT: Follow label directions

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For outdoor residential use only.

[To be placed on "Southern Labels"]
Not for Sale in the State of New York.

Using these directions

These "Directions for Use" are designed to help you determine the appropriate application procedure for this product.

Before using this product, make sure that you read the entire "Directions for Use" section of this label.

Before you Use

Read and follow these use precautions when you use this product:

- Apply only to lawns and ornamental turf that are well established.
- Wait at least 12 weeks after using this product before reseeding, overseeding or sprigging of grass into areas that have been treated.
- Do not use this product on new lawns or grass that has been weakened or stressed by drought, excessive temperatures or pests.
- When applying around ponds or other bodies of water, do not apply in windy conditions or where product can be moved into water by wind and washing rains.
- Keep people and pets off treated areas until dust has settled.
- Do not use grass clippings as mulch around vegetables or fruit trees.
- This product may be applied to flowers listed on this label.
- Do not make more than 1 application to the same area in the same year.

[NOTE: The following statement may be used on product labels sold only in the southern region]

- Do not make more than 2 applications to the same area in the same year.

How to use

1. Apply this product at the rate of: X.X lbs per 1000 sq. ft. (1 bag treats X,XXX sq. ft.)
2. Calibrate your spreader according to the spreader manufacturer's directions. Adjust settings as needed to apply the recommended rate of product.
3. Using a drop-type or rotary spreader, apply granules evenly over grass, avoiding streaking, skips or overlaps. Avoid using spreaders that apply product in narrow or concentrated bands.
4. Check calibration of spreader frequently to insure spreader is functioning properly and applying product evenly.

Spreader Settings

This XX pound bag covers X,XXX sq. ft. Use the settings below to apply at a rate of X.X lbs per 1,000 sq. ft.

Rotary Spreaders	Drop Spreaders
X	X
X	X
X	X
X	X
X	X

Where to use

This product can be used on the following types of grass when applied as directed:

Cool Season Grasses

- Agrostis Stolonifera
- Bentgrass, creeping
- Bluegrass, Kentucky
- Fescue fine
- Fescue tall
- Festuca arundinacea
- Festuca rubra
- Lolium, perenne
- Poa pratensis
- Rye grass, perennial

Warm Season Grasses

- Axonopus affinis
- Bahia
- Bermuda
- Buchloe dactyloides
- Buffalograss
- Carpetgrass
- Centipede
- Cynodon dactylon
- Eremochloa ophiuroide
- Kikuyugrass
- Paspalum notatum
- Pennisetum clandestinum
- St. Augustine
- Stenotaphrum secundatum
- Zoysia japonica
- Zoysiagrass

When to apply

- Product may be applied either in the Spring or Fall for the control of weeds.

[NOTE: The following statement may be used on product labels sold only in the southern region]

- Product may be applied in the early Spring and late Fall for the control of weeds. In no instance should applications be made less than 30 days apart and no more than 2 applications per year.

- [Claim for rates 0.18 – 0.25 pounds of active ingredient per acre] Product will provide up to 4 months of weed control when used as directed.
- [Claim for rates 0.15 – 0.17 pounds of active ingredient per acre] Product will provide up to 3 months of weed control when used as directed.
- To get control of listed weeds other than crabgrass, applications must be made before weeds emerge.
- Dimension® 6-25G crabgrass preventer for home lawns will control crabgrass if applied before it emerges.
- [Claim for rates 0.22 – 0.25 pounds of AI per acre] Dimension® 6-25G crabgrass preventer for home lawns will provide early post emergence control of crabgrass when applied within 4 weeks of germination.

Weeds Controlled

Pre- and Early Post Emergent

- Crabgrass

Pre-Emergent (Preventative)

Grass Weeds

- Annual Bluegrass
- Annual Ryegrass
- Barnyardgrass
- Crabgrass
- Crowfootgrass
- Goosegrass
- Green Foxtail
- Kikuyugrass (seedling)
- Perennial Ryegrass
- Smutgrass (seedling)
- Yellow Foxtail

Broadleaf Weeds

- Bittercress
- Buttercup Oxalis
- Carolina Geranium
- Chickweed
- Common Lespedeza
- Common Purslane
- Corn Speedwell
- Creeping Oxalis
- Henbit
- Parcloy-piert
- Pineappleweed
- Prostrate Spurge
- Spotted Spurge
- Yellow Woodsorrel

TOLERANT TREES, SHRUBS, AND FLOWERS

(A full list of Tolerant Trees, Shrubs, and Flowers is given as an addendum to this label.)

When applied as directed under the conditions on this label, trees, shrubs, and flowers listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Follow the use directions given above to determine plant tolerance under your growing conditions. A limited area involving only a few plants should be treated and observed for signs of yellowing, wilting, or lack of growth prior to large-scale application.

Common Name	Botanical Name	Tolerant Cultivars
Abella, Dwarf	Abella X grandiflora	Nana
Ajuga	Ajuga reptans	Bronze
	Ajuga genevensis	Bronze Beauty
Almond, Flowering	Prunus glandulosa	
Apple*	Malus pumila	
Arborvitae	Thuja occidentalis	Nigra; Pyramidalis; Smaragh Techny; Woodwardii.

* Ornamental Species only. DO NOT USE ON FOOD-PRODUCING TREES, SHRUBS, and FLOWERS.

Storage and Disposal

Take the following precautions:

IMPORTANT: Precautions during storage and disposal

- Keep out of reach of children
- Do not contaminate water, foodstuffs, feed or seed by storage or disposal
- Do not reuse empty product bags.

Storage conditions

Store Dimension® 6-25G in original container in a cool and dry location. The storage area should be locked and inaccessible to children.

Disposal

Sweep up any spilled material and use according to label directions.

Securely wrap the empty product bag in several layers of newspaper, and place in trash.

Important: If all of product can not be used according to label directions, make sure bag is securely sealed or wrapped up before disposing.

Conditions of Sale and Warranty

NOTICE: Buyer and user assumes all risk and liability of use, storage, and/or handling of this product not in accordance with the terms of this label.

-TOLL FREE-

Product Information: 1-800-874-8892

Manufactured By:
Pursell Industries, Inc.
P.O. Box 540
Sylacauga, AL 35150

Dimension® is a registered trademark of Rohm & Haas Company. POLYON® Patent Nos. 4,711,659; 4,804,403; 4,969,947; 5,374,292

THE PURSELL® PROMISE
YOUR SATISFACTION GUARANTEED OR YOUR MONEY BACK
If you apply this product as directed and are not completely satisfied, just send a proof of purchase from this package and a cash register receipt to the address below. We will promptly refund your money.

ADDENDUM

**Dimension® 6-25G
GRANULAR CRABGRASS PREVENTER FOR HOME LAWNS
EPA Reg. No. 8660-167**

TOLERANT TREES, SHRUBS, AND FLOWERS

When applied as directed under the conditions on this label, trees, shrubs, and flowers listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Follow directions given above to determine plant tolerance under your growing conditions. A limited area involving only a few plants should be treated and observed for signs of yellowing, wilting, or lack of growth prior to large-scale application.

Common Name	Botanical Name	Tolerant Cultivars
Abella, Dwarf	Abella X grandiflora	Nana
Ajuga	Ajuga reptans	Bronze
	Ajuga genevensis	Bronze Beauty
Almond, Flowering	Prunus glandulosa	
Apple*	Malus pumila	
Arborvitae	Thuja occidentalis	Nigra; Pyramidalis; Smaragh Techny; Woodwardii.
Arborvitae, Golden	Thuja orientalis	
Aster, Chinese	Callistephus chinensis	Dwarf Queen
Ash Green	Fraxinus pennsylvanica	
Ash, Mountain	Sorbus aucuparia	
Ash, Purple	Fraxinus americana	
Azalea	Rhododendron spp.	Brilliant; Buccaneer; Carror; Chimes (Belgian); Exbury; Fashion; Hardijzer Beauty; Hershey Red; Higasa; Hinoocrimson; Holland (Hybrid); Marion Lee; Northern Lights; Orange Cup; Orchid Lights; Snow; Southern Charm.
Azalea, Flame	Rhododendron calendulaceum	
Azalea, Kirishima		
Bamboo, Heavenly		
Barberry	Berberis thunbergii	Aurca; Dwarf Pigmy; Green; Kobold; Pygmy Red; Rose Glow
Barberry, Purple		Atropurpurea
Bearberry(common)	Arctostaphylos uva-ursti	Massachusetts
Birch, River	Betula nigra	
Blueberry*	Vaccinium spp.	Bluecrop; Blue Jay; Jersey; North Blue; Northland
Bottlebrush	Callistemon citrinus	
Boxwood, Japanese		Japonica
Boxwood, Weller	Buxus semperirens	
Broom	Cytisus spp.	Moonlight
Broom	Genista pilosa	Vancouver Gold
Bugle Carpet		
Camellia	Camellia japonica	Chansonette; Debutante; Mathotiana Supreme
Camellia	Camellia sasangua	
Carex, Variegated	Carex	
Cedar, Red	Juniperus virginiana	
Cockscomb, Plumosa	Celosia cristata	Red Kewpie
Coleus	Coleus blumei	
Copper leaf	Acalypha wilkesiana	
Cotoneaster	Cotoneaster apiculatus	
Coyotebrush	Baccharis pelularis	
Cycads	Cycads revoluta	
Cypress, Bald	Taxodium distichum	
Cypress, Italian	Cupressus sempervirens	Glauca
Cypress, Japanese False	Chamaecyparis obtus	Crysil's
Cypress, Leyland	Cupressocyparis leylandii	
Daffodil	Narcissus spp.	King Alfred
Daylily	Heemerocallis spp.	Aztec Gold; Bright Yellow (Hybrid); Single Gold (Evergreen); Wilson's Yellow
Dogwood	Cornus florida	
Dogwood, American	Cornus sericea	Flavirima

Douglas Fir	<i>Pseudotsuga menziesii</i>	
Dusty Miller	<i>Senecio cineraria</i>	
Elm, Drake	<i>Ulmus parvifolia</i>	
Euonymus	<i>Euonymus fortunei</i>	Argento-variegata; Auereo-marginata; Colorata; Emerald Gaiety; Emerald 'N Gold; Gold Edge; Gold Princess; Silver King; Tricolor; Vegetus.
Fan Palm, European	<i>Chamaerops humilis</i>	
Fan Palm, Mexican	<i>Washingtonia robusta</i>	
Fescue	<i>Festuca glauca</i>	
Fetterbush	<i>Leucothoe fontiana</i>	Rainbow
Ficus	<i>Ficus retusa</i>	Nitida
Forsythia	<i>Forsythia X intermedia</i>	Arnold Dwarf; Bronxensis Dwarf; Lynwood gold; Meadowlark; Spring Glory; Weeping.
Fir Fraser	<i>Abies fraseri</i>	
Gardenia	<i>Gardenia jasminoides</i>	Mystery; Radicans
Geranium	<i>Pelargonium X hortorum</i>	
Gum	<i>Eucalyptus citriodora</i>	
Hawthorn	<i>Crataegus spp.</i>	Cockspur white; Crimson Cloud; Enchantress; Jack Evans; Washington White
Heather, Twisted	<i>Erica cinerea</i>	Mediterranean Pink
Hibiscus	<i>Hibiscus spp.</i>	Blue Bird; Brilliant; hula Girl
Hemlock, Canada	<i>Tsuga canadensis</i>	
Holly	<i>Ilex spp.</i>	Blue Boy; Blue Girl;
	<i>Ilex X meserveae</i>	Burfodii; China Girl; Compacta; Forsteri; Hellerie; Japanese Northern Beauty; Needlepoint; nellie R. Stevens; Savannah
Holly	<i>Ilex X attenuata</i>	
Holly, Chinese	<i>Ilex comuta</i>	
Holly, Japanese	<i>Ilex crenata</i>	
Holly, Yaupon	<i>Ilex vomitoria</i>	
Honeysuckle	<i>Lonicera japonica</i>	Clavey's Dwarf; Halliana; Tatarian Canadian White; Zebelli Red
Hosta	<i>Hosta spp.</i>	Hosta
Hosta	<i>Hosta lancifolia</i>	Albo Marginata
Ice Plant	<i>Carpobrotus edulis</i>	
Impatiens	<i>Impatiens wallerana</i>	
Iris	<i>Iris spp.</i>	Dwarf Blue; Wedgewood
Ivy, English	<i>Hedera helix</i>	Bulgaria; Thorndale
Jasmine, Asian	<i>Trachelopermum asiaticum</i>	
Juniper	<i>Juniperus spp.</i>	Arcadia; Armstrong; Bar Harbor
Juniper	<i>Juniperus horizontalis</i>	Blue Chip; Blue Pacific; Blue Rug; Blue Star; Broadmoor; Buffalo; Calgary Carpet; Emerald Sea; Emerald Spreader; Endora Compacta; Fruitlandi; Green; Gold Tip; Hetzi; Hughes; Manhattan Blue; Parsoni; Pfitzeriana; Plumosa; Prince of Wales; Procumbens Dwarf; San Jose; Sargent Blue; Sargent Green; Scandia; scopulorum Moonglow; Scopulorum skyrocker; Spartan
Juniper	<i>Juniperus chinensis</i>	Tamariscifolia
Juniper	<i>Juniperus sabina</i>	Weberi; Youngstown; Yukon Belle
King Palm	<i>Archontophoenix cunninghamiana</i>	
Laurel, Australian	<i>Pittosporum tobina</i>	
Laurel, Mountain	<i>Kalmia latifolia</i>	
Leucothoe	<i>Leucothoe fontanesiana</i>	
Lily, African	<i>Agapanthus africanus</i>	Albus; Peter pan
Lily, African blue		
Lily of the Valley	<i>Pieris japonica</i>	Mt. Fire
Lilyturf	<i>Liriope muscari</i>	Evergreen Giant; Lilac Beauty; majestic; monroe White; Variegata
Ligustrum, Japanese	<i>Ligustrum japonicum</i>	
Liriope, Green	<i>Liriope spicata</i>	
Maple, Japanese	<i>Acer japonicum</i>	
Maple, Norway	<i>Acer platanoides</i>	
Maple, Red	<i>Acer rubrum</i>	
Maple, Silver	<i>Acer saccharinum</i>	
Maple, Sugar	<i>Acer saccharum</i>	
Marigold	<i>Tagetes patula</i>	Honeycomb; Variegata; Wheeler's Dwarf
Mock Orange*	<i>Philadelphus spp.</i>	Golden; Snowflake; Double White
Mondo Grass	<i>Ophiopogon japonicus</i>	
Moss Rose	<i>Portulaca Grandiflora</i>	Sunyside
Myrtle, Crape	<i>Lagerstoemia indica</i>	Faurei; Langer; Muskogee; Standard Pink
Myrtle, Wax	<i>Myrica cerifer</i>	
Nadina	<i>Nadina domestica</i>	Compacta; Nana
Oak, Laurel	<i>Quercus laurifolia</i>	
Oak, Pin	<i>Quercus palustris</i>	

Oak, Red	Quercus rubra	
Oak, Southern L.	Quercus virginiana	
Oak, Willow	Quercus phellos	
Oleander	Nerium oleander	Hardy Red; Petite Pink; Sister Agnes
Oteospermum	Osteospermum fruticosum	Wirligig
Palm, Bangalow		
Pampas Grass	Cortadaria selloana	
Paper Flower	Bougainvillea glabra	Barbara Karst
Pachysandra	Pachysandra terminalis	
Peach*	Prunus persica	
Periwinkle, Dwarf	Vinca minor	
Petunia	Petunia X hybrida	Picoti
Photinia, Red Tip	Photinia X fraseri	
Pieris	Pieris japonica	
Pine, Australian	Pinus nigra	
Pine, Japanese Black	Pinus thumbergiana	
Pine, Loblolly	Pinus taeda	
Pine, Longleaf	Pinus palustris	
Pine, Mugho	Pinus mugho	
Pine, Scotch	Pinus sylvestris	
Pine, Slash	Pinus elliotti	
Pine, Swiss Mt.	Pinus mugo	
Pine, Virginia	Pinus virginiana	
Pine, White	Pinus strobus	
Pineapple, Guava	Feijoa fellowaiana	
Pittosporum, Japan		
Potentilla	Potentilla fruticosa	Abbotswood
Privet	Ligustrum japonicum	Golden Vicary; Regal; Texanum; Wax; Yellow Tipped
Privet, Glossy	Ligustrum lucidum	
Pyracantha	Pyracantha koidzumii	Gnome; Lalandei; Victory
Queen Palm	Arecastrum rammanzoffanum	
Quince, Japanese*		
Rhododendron	Rhododendron spp.	Album; Cunningham White; Fashion; Hardy; PJM; Purple Gem; Silvery Pink
Rhododendron, Carolina	Rhododendron carolinianum	
Rhododendron, Catawba	Rhododendron catawbiense	
Rodie Max (Rosebay)	Rhododendron maximum	
Rose	Rosa banksiaae	Luta
Rosemary*	Rosmarinus officinalis	
Rosemary, Bog	Andromeda polifolia	Nana
Salvia	Salvia farinacea	Rhea
Sedum	Sedum spurium	Dragon Blood Red; Red Carpet; Yellow
Spiraea	Astilbe X arendsii	Fanall
Spiraea	Spiraea spp.	Anthony Waterer Red; Dolchica; Froebeli Pink; Goldenflame Red; Snowmound White; Van Houtte White
Spiraea, Garland	Spiraea spp.	
Spruce, Black Hills	Picea glauca	
Spruce, Colorado Blue	Picea pungens	
Spruce, norway	Picea ables	
Spruce, White	Picea glauca	Conica
Sourwood	Oxydendrum arboreum	
Sweetflag, Grassy-Le		
Sweetgum	Liquidambar styraciflua	
Sycamore	Plantanus accidentails	
Tree Fern (Tiki Fern)	Asparagus virgatus	
Trumpet Flower, evening	Gelsemium sempervirens	
Tulip	Tulip spp.	Apeldoorn
Verbana, Shrub	Lantana sellowiana	
Vervain	Verbena spp.	St Paul
Viburnum	Viburnum spp.	American Cranberry Bush; Arrowwood; Common Snowball; European Cranberry Bush; Linden; Mohican; Wright
Windmill Palm	Trachycarpus fortunei	
Xylosma	Xylosma congestum	
Yaupon	Ilex vomitoria	Dwarf
Yew	Taxus cuspidata; Taxus X media	Denise, mis

* Ornamental Species only. DO NOT USE ON FOOD-PRODUCING TREES, SHRUBS, AND FLOWERS

ADVERTISING CLAIMS

[May be used on some product labels.]

1. Preemergence and early Post Emergence Control of Crabgrass
2. Covers 5,000 sq. ft.
3. Money-Back Guarantee.
4. Look for other quality PURSELL lawn and garden products. For more information about this product visit us on the Internet at www.fertilizer.com or call our Consumer Service Hotline at 1-800-874-8892.
5. Non-Staining
6. Provides up to 4 months control of crabgrass
7. Control Crabgrass for up to 120 days
8. IMPORTANT – It is important to follow directions as indicated on this quality lawn care product as over application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to calibrate spreader before beginning and check calibration during application, adjusting if necessary. Consult Owner's Manual.