



Stage 2-The Caterpillar

This Health Chem SPRAY N KILL Gypsy Moth Caterpillar Biological Spray is your weapon against the ravenous gypsy moth caterpillar. Caterpillars which emerge from the egg mass deposited by the female gypsy moth the previous summer can easily spread to surrounding trees and plants starting in early spring. As the caterpillars feed on leaves they grow to become fat hairy caterpillars about 2" long. Control is best if it begins early before ravenous gypsy moth caterpillars have stripped trees and valuable ornamental plants of their leaves and while gypsy moth caterpillars are still small.

SPRAY N KILL Gypsy Moth Caterpillar Biological Spray is a highly selective biological insecticide for use against gypsy moth caterpillars, tent caterpillars, bagworms, spring and fall cankerworms, and fall webworms.

SPRAY N KILL Gypsy Moth Caterpillar Biological Spray does not harm honey bees foraging treated areas. It does not interrupt activities of beneficial insects.

PRECAUTIONARY STATEMENTS

CAUTION: Avoid contact with skin, eyes or clothing. In case of contact immediately flush eye or skin with plenty of water. Get medical attention if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL

- 1. Treat infested plants first when caterpillars are small.
- 2. Coverage of both the upper and under surfaces of leaves is essential since SPRAY N KILL is most effective when it reaches the caterpillars. After eating SPRAY N KILL caterpillars stop feeding within the hour and within a few days.

- 3. Dying caterpillars move slowly, discolor, then shrivel, blacken and die.
- 4. Repeat applications of SPRAY N KILL as needed to maintain control, usually 3 to 14 days depending on plant growth, rainfall, and infestation level.

MIXING:

SPRAY N KILL may be used with both pressure and hose-attached garden sprayers. Always shake or stir SPRAY N KILL thoroughly before use. Pour recommended amount of SPRAY N KILL onto surface of water in a nearly filled container. Maintain agitation. Do not allow dilute sprays to stand for more than 12 hours.

INSECT PEST	APPLICATION RATE
Gypsy moth caterpillar	Use 4 oz. (8 tablespoons) in 20 gals. water
Bagworm	
Tent caterpillar	Use 2 oz. (4 tablespoons) in 20 gals. of water
Spring and fall cankerworms	
Fall webworms	

Use of a spreader sticker in the spray helps keep the SPRAY N KILL on the foliage, thus maintaining its effectiveness where heavy dew or rains occur.

STORAGE AND DISPOSAL


Storage: Reclose containers of unused product. Store in a cool place, out of direct sunlight and away from heat.

Disposal: Do not reuse container. Discard empty container according to local trash disposal regulations.

NOTICE: Buyer assumes all responsibility for safety. Use in accordance with directions.

List No. 6032
EPA Registration No. 8730-26
EPA Establishment No. 8730 PA 01
Manufactured by
Hercules Inc.
HEALTH-CHEM CORPORATION
107 Broadway, New York, New York 10037

100% HEALTHY ORGAN
SPRAY N KILL
AN INSECTICIDE FOR PEST MANAGEMENT PURPOSES



GYPSY MOTH CATERPILLAR BIOLOGICAL SPRAY

FOR CONTROL OF CATERPILLARS
DOES NOT KILL BEES

ACTIVE INGREDIENT
Bacillus thuringiensis Berliner, 8 800 International Units of Potency per mg. (32 billion International Units per gallon)

NET CONTENTS: 8 fl. oz. (236 ml)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

8730 26