

USES

SEEDS AND SOILS: EZY-WEED provides excellent weed control for lawns, flower beds, vegetable gardens, and other areas where weeds are a problem. It is especially effective against annual weeds that germinate in late fall and early spring. For best results, EZY-WEED should be made in late fall and early spring. A second application can be applied in the spring for early fall germinating weeds. Do not apply after first bloom.


ROSES and PERENNIALS: Chrysanthemums, Shasta Daisies, etc.
Newly Prepared Beds: After your rose or perennial bed has been planted, water the area around the plants to settle soil firmly. Be sure to enter area to be treated free of actively growing weeds since EZY-WEED only kills germinating weed seed and does not kill actively growing plants. The soil to be treated should also be well raked to remove any large clods of soil. Using the shaker can, apply EZY-WEED at a rate of 1 pound per 200 square feet. For large areas, use the larger area. After application of EZY-WEED, thoroughly water the treated area. This will release the chemical weeds from the soil granules and move it into the soil.

Established Flower Beds: In early spring, before weeds begin to grow, cultivate existing beds, removing all litter and debris. Remove all large clods of soil by raking. Apply EZY-WEED, as directed, see above, from shaker can and water the treated area thoroughly to release the chemical weed killer from the granules and to move it into the soil.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

CAUTION!

DO NOT TAKE INTERNALLY.
 AVOID CONTACT WITH EYES.
 WASH AFTER HANDLING OR USING.



Net Contents
1 Pound

EZY-WEED

A pre-emergence herbicide for the control of crabgrass and other annual grasses and broad-leaf weeds such as green foxtail, yellow foxtail, Johnsongrass from seed, Dove grass, witchgrass, carpetweed, common chickweed, lambquarters, purslane and Florida pusley in strawberry, rose, and flower gardens. Contents of this can will treat an area of 200 square feet.

CAUTION — Keep out of reach of children

See other cautions on side panel.

Active ingredient:
 Dimethyl ester of Tetrachloroterephthalic acid 50%
 Inert ingredients 950%
 Total 1000% (100 lbs)

U.S. Patent No. 2,923,034
 U.S.D.A. Reg. No. 5702-1

Packed for

Jackson & Perkins Co.
 World's Largest Rice Growers and Manufacturers
 Medford, Oregon 97501

DIRECTIONS

Do Not Use on the following plants:
 Asparagus
 Butter Beans
 Corn
 Cucumbers
 Eggplant
 Green Beans
 Green Peas
 Green Potatoes
 Green Squash
 Green Tomatoes
 Green Zucchini
 Kidney Beans
 Lima Beans
 Melons
 Okra
 Peas
 Potatoes
 Pumpkins
 Radishes
 Snap Beans
 Squash
 Tomatoes
 Zucchini

APPLICATION

Apply EZY-WEED using the shaker can at the following rates indicated on the can for each type of growing medium and plant. The table below indicates the amount of EZY-WEED needed to treat specific areas or plant beds.

100 square feet	Apply 1/2 lb.
200 square feet	Apply 1 lb.
400 square feet	Apply 2 lbs.

When To Apply: EZY-WEED may be applied at any of the times indicated below.

In the early spring prior to weed emergence.
 Immediately after raking, or right cultivation or weeding.
 Immediately after transplanting, or at least 10 days after plants have been weeded or cultivated.

For Best Results: Apply EZY-WEED to a uniform weed-free soil surface.

Measure area to be treated and then apply the prescribed amount of EZY-WEED.

Water the treated area thoroughly after application of EZY-WEED.

Do not disturb soil after EZY-WEED application.

Shake or wash EZY-WEED can thoroughly.

CAUTION

DO NOT STORE NEAR FEED OR FEEDING EQUIPMENT, LIZERS, OR OTHER PESTICIDES OR CONTAMINANTS.

STORE IN ORIGINAL CONTAINER.

