

DIRECTIONS FOR USE

WHEN AND HOW TO APPLY.

Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

CORN (Post-emergence): For control of annual broadleaf weeds, use 1 pint per acre in adult corn. For control of grasses, use 1 quart per acre in adult corn. For control of weeds in corn, use 1 quart per acre in adult corn.

SMALL GRAINS (Barley, Oats, Rye and Wheat): Annual should be sprayed after seeding (pre-emergence) or at the 1st leaf stage. Do not apply from boot to milk or maturity stage. Do not use in grain intersplanted with a legume, particularly alfalfa or clover cover, unless an interval of 2 weeks has elapsed between the grazing or feeding of treated hay within 2 weeks after treatment.

Annual and Biennial Weeds: Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

Perennial Weeds: Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

GRASS SEED CROPS (Fescue, Kentucky bluegrass, Brome, Bluegrass, Orchard grass, Ryegrass): Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

NON-CROPPED LAND: Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

Annual and Biennial Weeds: Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

Deep Rooted Perennials and Woody Plants: Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

NOTICE: Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.



BUTYL ESTER 6

2,4-D

WEED KILLER

EMULSIFIABLE

CONTAINS 6 POUNDS 2,4-D ACID PER GALLON

ACCEPTED
7/13/72
REGISTERED
SUBJECT
ATTACHED

ACTIVE INGREDIENTS:

Butyl Ester of 2,4-Dichlorophenoxyacetic acid 78.00%

INERT INGREDIENTS

TOLUENE 100.00%

2,4-Dichlorophenoxyacetic acid

NET CONTENTS _____ GALLONS

CAUTION

May cause skin irritation. Avoid contact with eyes, skin, and clothing.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

DO NOT USE EMPTY CONTAINER — PERFORATE, CRUSH, BURY — DO NOT BURN

"HARMFUL IF SWALLOWED"

Protect from Temperatures Below 0 F.

WARNING

Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.

RECOMMENDATIONS

Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth. Apply to the soil surface in the early stages of weed growth.