




(Front)



ArthrobanTM

LAWN



INSECT KILLER

CONTROL THESE INSECTS IN TURF

EUROPEAN CHAFER AND
JAPANESE BEETLE GRUBHYPERODES
WEEVIL

SOD WEBWORM



CHINCH BUG

TREAT AT RATE OF 5 LBS. PER 1000 SQ. FT.
(PACKAGE TREATS 5000 SQ. FT. 50'x100')TREAT AT RATE OF 2 1/2 LBS. PER 1000 SQ. FT.
(PACKAGE TREATS 10,000 SQ. FT. 100'x100')USE WHEN INSECTS ARE ACTIVE
NET WEIGHT 25 LBS.

ACTIVE INGREDIENTS:

Chlorpyrifos* 0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate 0.92%

INERT INGREDIENTS: 99.08%

*U.S. Patent No. 3,244,586.

E.P.A. Reg. No. 8780-34
E.P.A. Est. No. 8780-NY-1

MANUFACTURED IN NEW YORK STATE FOR USE IN NEW YORK STATE


CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER SIDE FOR ADDITIONAL CAUTIONS AND DIRECTIONS

Manufactured by
HIGH POINT MILLS, INC.
1225 Lehigh Station Road,
Hempstead, N.Y. 11467

Arthroban

Arthroban

2 92



TM

ACCEPTED
JUN 11 1976
 Under the Federal Insecticide, Fungicide, and Rodenticide Act
 as amended
 registered under
 EPA Reg. No. 8780-34

ArthrobanTM

LAWN



INSECT KILLER

APPLICATION DIRECTIONS

Treat when insects are active or as recommended by State Agricultural Experiment Station or State Extension Service Specialists.
 Apply Arthroban at the rate of 5 lbs. per 1000 sq. ft. for European Chafer Grubs, Japanese Beetle Grubs and Hyperodes Weevil.
 Apply Arthroban at the rate of 2 1/2 lbs. per 1000 sq. ft. for Sod Webworms and Chinch Bugs.
 Immediately after spreading water lightly using 30 gals. per 1000 sq. ft.
 After 2 to 3 days water again; apply about an inch of water. Rainfall during this period may be counted, but should be measured.

SPREADER SETTINGS

Due to variations in individual machines, these settings are intended to be a guide. It is important to make certain that your machine is spreading the correct amount of material at the setting used.

MAKE OF SPREADER	COVERAGE
5000 sq. ft.	10,000 sq. ft.
Turf Line Rotary	3.75"
Cyclone Rotary	3.75"
Scott Conventional	5.0

*Based on a spreading width of 6 feet.

CAUTION! KEEP OUT OF REACH OF CHILDREN AND ANIMALS MAY BE HARMFUL IF SWALLOWED

Avoid eye contact. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from food, feedstuffs and domestic water supplies.
 NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.
 Keep children and pets off treated areas until grass is dry after initial watering with 30 gals. per 1000 sq. ft.
 This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.
 Use only on home lawns, golf courses and other turf areas not used for pasture or hay.
 Do not reuse container. Bury with waste in a safe place. Do not burn.

FBUS201

Freight Shipping Box
 Handle Mechanically
 APPLICABLE FREIGHT CLASSIFICATION
 Shippers Use ST, Grade 10000 CO.

