
Page 1/92



TM

33 Lbs Not Wt TREATS

10,000 Sq. £t

FOLLOW DIRECTIONS CAREFULLY.

rthroha

Plus


CONTROLS THESE INSECTS IN TUFF

EUROPEAN CHAFFE AND JAPANESE BEETLE GRUB



TREAT AT RATE OF 66 LBS PER 1000 SQ # PACKAGE TREATS 5000 SQLFT 50 k100





(PACKAGE THEATS 10 000 SQ FT 100 V100 USE WHEN INSECTS ARE ACTIVE

Contains 11 essential elements to provide the best results regardless of grass type and soil quality.

APPLICATION DIRECTIONS

Treat when insects are active or as recommended by State Agricultural Experiment Station or State Extension Service Specialists.

Apply Arthroban at the rate of 6.6 (bs. per 1000 sq. ft. for European Chafer Grubs, Japanese Beetle Grubs and Hyperodes Weevil.

Apply Arthroban at the rate of 3.3 lbs. per 1000 sq. ft for Sod Webworms and Chinch Bugs. Immediately after spreading water lightly using 30 gals. per 1000 sq. ft.

After 2 to 3 days water again; apply about an inch of water. Rainfall during this period may be counted, but should be measured.

SPREADER SETTINGS

Due to variations in individual machines, these settings are intended to be a guide, it is important to make certain that your machine is streading the correct amount of material at the setting used.

Coverage .ft. 10,000 sq. ft. 5000 sq. ft. MAKE OF SPREADER Turf Line Rotary Cyclone Rotary Scott Conventiona' 3.75* 3.75* 5.0 2.75* 2.75* 3.5 "Based on a spreading width of 6 feet

CAUTION! KEEP OUT OF REACH OF CHILDREN AND ANIMALS MAY BE HARMFUL IF SWALLOWED

Avoid eye contact. Flush contaminated eyes with plenty of water and get medical attention ± imitation persists. Mash thoroughly after handling. Wash contaminated clothing before re-use. Keep away from food feedstuffs and comestic water supplies.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an amidote

Keep children and pets off treated areas until grass is dry after initial watering with 30 gals, per 1000 sq. ft

This product is fosic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of leves streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of exponent, or dispose of wastes. Apply this product only as specified on this label. Use only on home tawns, golf courses and other turn areas not used for pasture or hay. Do not reuse container. Bury with waste in a safe place. Do not burn.

Do not apply when wind conditions would cause drift from areas treated.

45



TM

TREATS

FOLLOW DIRECTIONS CAREFULLY.

Plus

CONTROLS THESE INSECTS IN TURF

EUROPEAN CHAFER AND JAPANESE BEETLE GRUB

HYPERODES WEEVIL



TREAT AT RATE OF 66 LBS. PER 1000 SQ. FT (PACKAGE TREATS 5000 SQ. FT. 50×100)

SOD WEBWORM





(PACKAGE TREATS 10,000 SQ FT. 100'x100') USE WHEN INSECTS ARE ACTIVE

Contains 11 essential elements to provide the best results regardless of grass Type and soil quality.

NET WEIGHT 33 LBS.

ACTIVE INGREDIENT

INERT INGREDIENTS 99.30%

*U. S. Patent No. 3,244,586

GUARANTEED ANALYSIS

GUARANTEED ANALYSIS
Total Nitrogen (N)
5.5 solvible nitrogen derived from unter and me
4.5x insolvible nitrogen derived from administer
2.0x solvible nitrogen derived from administer
2.0x solvible nitrogen derived from administer
5.0xuble Potash (KGO)
Suffer (S)
Magmesium (Mg)
from (Fe)
Manganese (Mn)
Cooper (Cu)
Zinc (Zn)
Boron (B)
Molybdenum (Mo)

KEEP OUT OF REACH OF CHILDREN

CAUTION SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

MANUFACTURED IN NEW YORK STATE FOR USE IN NEW YORK STATE

Manufactured By: HIGH POINT MILLS, INC. 1225 LEHIGH STATION ROAD HENRIETTA, NEW YORK 14467

EPA Establishment No. 8780-NY-1 EPA Registration No. 8780-42

Copyright 1976