

TREATS
10,000 SQ. FT.
(100 ft. X 100 ft.)

WEED KILLER

PLUS

TURF FERTILIZER

20-6-4


TREATS
10,000 SQ. FT.



KILLS
DANDELION
PLANTAIN
CHICKWEED

&

SIMILAR
BROADLEAF
AND
VINEY WEEDS



**MULTIPLES GRASS
AND MAKES IT GREEN
LONGER**



NET WEIGHT 50 LBS.

ACTIVE INGREDIENTS

2 (2-methyl-4-chlorophenoxy) propionic acid
(dichlorophenoxy herb)

1.80%

2,4-Dichlorophenoxyacetic acid sodium salt
(methyl ester)

0.87%

INERT INGREDIENTS

97.33%

*Figure is 0.70% 2 (2-methyl-4-chlorophenoxy) propionic acid
**Figure is 0.17% 2,4-dichlorophenoxyacetic acid

Manufactured By

HIGH POINT MILLS, INC.

1225 LEHIGH STATION ROAD

PERKINNETTA, NEW YORK 14527

EPA REGISTRATION NO. 147-115-1

1/24/74, 1/24/74, 1/24/74, 1/24/74

CAUTION



CONTAINS ELEVEN ESSENTIAL ELEMENTS INCLUDING
EXTENDED TIME RELEASE NITROGEN TO PRODUCE
EXCELLENT RESULTS REGARDLESS OF GRASS TYPE
OR SOIL QUALITY.

GUARANTEED ANALYSIS

Total Nitrogen	2.00%	Total Phosphorus	0.60%	Total Potassium	4.40%
Water Soluble Nitrogen	0.50%	Water Soluble Phosphorus	0.15%	Water Soluble Potassium	1.10%
Ammoniacal Nitrogen	0.10%	Ammoniacal Phosphorus	0.05%	Ammoniacal Potassium	0.20%
Plant Available Nitrogen	0.40%	Plant Available Phosphorus	0.10%	Plant Available Potassium	0.90%
Water Soluble Nitrogen	0.50%	Water Soluble Phosphorus	0.15%	Water Soluble Potassium	1.10%
Ammoniacal Nitrogen	0.10%	Ammoniacal Phosphorus	0.05%	Ammoniacal Potassium	0.20%
Plant Available Nitrogen	0.40%	Plant Available Phosphorus	0.10%	Plant Available Potassium	0.90%



TREATS
10,000 SQ. FT.
(25' x 40')

WEED KILLER

PLUS


TURF FERTILIZER

20-6-4 TREATS
10,000 SQ. FT.



KILLS

&



MULTIPLES GRASS
AND MAKES IT GREEN
LONGER



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

STATEMENT OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS

DIRECTIONS FOR USE

CAUTION

WHEN TO USE

SPRING TREATMENT

SUMMER TREATMENT

FALL TREATMENT

HOW TO APPLY


44-38861-1000

**PLUS
TURF FERTILIZER**

20-6-4 TREATS
10,000 SQ. FT.



SIMILAR
BROADLEAF
AND
VINEY WEEDS



MULTIPLES GRASS
AND MAKES IT GREEN
LONGER 



CONTAINS ELEVEN ESSENTIAL ELEMENTS, INCLUDING

20 0 10,000 SQ. FT.



KILLS
DANDELION
PLANTAIN
CHICKWEED

&

SIMILAR
BROADLEAF
AND
VINEY WEEDS

RED SOREL
WILD ONION
STITCHWORT
WILD GARLIC

ENGLISH DAISY
YARROW
RED CLOVER
WHITE CLOVER
MOUSE-EARED CHICKWEED

BUCKHORN
CURLED DOCK
GROUND IVY
KNOTWEED



**MULTIPLES GRASS
AND MAKES IT GREEN
LONGER**



NET WEIGHT 50 LBS.

ACTIVE INGREDIENTS:

2-(4-chlorophenyl)-4-chlorophenyl phosphonic acid, 0.50%

2, 4-dichlorophenyl phosphonic acid, sodium salt, 0.50%

0.50%

*Up to 0.50% 2-(4-chlorophenyl)-4-chlorophenyl phosphonic acid.

*Up to 0.50% 2, 4-dichlorophenyl phosphonic acid.

Manufactured By

HIGH POINT MILLS, INC.
1235 LEBSON STATION ROAD
MINNETTA, NEW YORK 14467

EPA Establishment No. 8780-NY-1
EPA Registration No. 8780-44

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



**CONTAINS ELEVEN ESSENTIAL ELEMENTS, INCLUDING
EXTENDED TIME-RELEASE NITROGEN, TO PRODUCE
EXCELLENT RESULTS REGARDLESS OF GRASS TYPE
OR SOIL QUALITY.**

GUARANTEED ANALYSIS

Total Nitrogen (N)	20.0%	Boron (B)	0.02%
15.1% Urea nitrogen		Molybdenum (Mo)	0.0000%
0.9% Ammonia-nitrogen		Copper (Cu)	0.05%
4.0% Water insoluble nitrogen		Iron (Fe)	1.00%
Available Phosphate Acid (P ₂ O ₅)	0.0%	Manganese (Mn)	0.05%
Soluble Potash (K ₂ O)	4.0%	Zinc (Zn)	0.05%
Magnesium (Mg)	0.0%		
Sulfur (S)	1.0%		

* Derived from ammoniated phosphate
** Derived from activated sludge and
Isobutylic acid

Copyright 1977



TREATS
10,000 SQ.
 (100 ft X 100 ft)

WEED KILLER

PLUS

TURF FERTILIZER


20-6-4 TREATS
 10,000 SQ. FT.



KILLS
 DANDELION
 PLANTAIN
 CHICK WEED

&

SIMILAR
 BROADLEAF
 AND
 VINEY WEEDS



**MULTIPLES GRASS
 AND MAKES IT GREEN
 LONGER**



PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS (4 DOMESTIC ANIMALS)

CAUTION

STATEMENT OF PRACTICAL TREATMENT

ENVIRONMENTAL HAZARDS

WHEN TO USE

SPRING TREATMENT

SUMMER TREATMENT

FALL TREATMENT



KILLS
DANDELION
PLANTAIN
CHICKWEED


&

SIMILAR
BROADLEAF
AND
VINEY WEEDS

REEDS, RIBS,
WEE, LAMB,
STICKLEWORT,
AND D. LAMB.

REEDS, RIBS,
WEE, LAMB,
STICKLEWORT,
AND D. LAMB.

REEDS, RIBS,
WEE, LAMB,
STICKLEWORT,
AND D. LAMB.



**MULTIPLES GRASS
AND MAKES IT GREEN
LONGER**



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)**

CAUTION

Avoid contact with skin, eyes, or clothing.
Avoid breathing dust.
Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water. For eyes,
flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, or ponds. Do not apply when weather
conditions favor drift from target area.

DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent
with its labeling.

STORAGE AND DISPOSAL

Store this product in a safe place beyond the reach of children and
pets. Do not contaminate water, food, or feed by storage, disposal,
or cleaning of equipment. Do not store near other pesticides or
seeds. Any excess material should be saved for later use in confor-
mation with the directions on this package. Do not reuse empty
container. Wrap container and put in trash collection.

WHERE TO USE

Use this product on home lawns or other turf areas not intended for pas-
ture or hay. Do not use on centipede grass, St. Augustine, or dichondra,
nor on lawns where desirable bluegrass is present. Do not apply to newly
seeded lawns until after two mowings.

WHEN TO USE

Use when weeds are present in lawns, golf courses, or other areas that
aren't growing normally.
Or, as applicable, on all areas where you wish to keep weeds from
growing.

Applications made when weeds are in the seed stage are most effective.

SPRING TREATMENT

Apply as soon as weeds appear.

SUMMER TREATMENT

Use if weeds are a problem in late summer. Summer treatments are ade-
quate to promote normal growth in the autumn. If it rains, it may be 3-5 days
before you can apply the next application at 4-6 degrees.

FALL TREATMENT

Use if weeds are a problem in late summer. Summer treatments are ade-
quate to promote normal growth in the autumn. If it rains, it may be 3-5 days
before you can apply the next application at 4-6 degrees.

HOW TO APPLY

Do not mow the lawn or garden before or after treatment. The grass
should be in the seed stage. Do not apply to the lawn or garden if it is raining or
if it is windy.

Do not apply or allow to drift on lawns, shrubs, ornamentals, trees, fruits,
or vegetables.

Use a garden type spreader or a backpack type spreader. The backpack type
spreader should be used on lawns. Use a backpack type spreader setting
of 10-15 lbs. per 1000 sq. ft. of area. Do not apply to the lawn or garden if it is
raining or if it is windy. Do not apply to the lawn or garden if it is raining or
if it is windy. Do not apply to the lawn or garden if it is raining or if it is windy.

Do not apply to the lawn or garden if it is raining or if it is windy. Do not
apply to the lawn or garden if it is raining or if it is windy. Do not apply to the
lawn or garden if it is raining or if it is windy.

SPREADER SETTINGS

TURF LITE	3
CYCLONE	3
SCOTT	4

*Based on 6 ft. spreading width.