

8780-51

07/06/1979

1/2

TURF LINE DIAZINON® 5G. LAWN INSECT CONTROL

Treats up to 12,000 square feet for general lawn insect control, higher rates are for white grubs and for use on heavy lawns. (See directions for use)

25 LBS. NET WEIGHT

ACTIVE INGREDIENT:

0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 5.0%

INERT INGREDIENTS: 95.0%


**KEEP OUT OF REACH OF CHILDREN
WARNING**

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS


Manufactured by: HIGH POINT MILLS, INC.
P.O. Box 36
1225 Lehigh Station Rd.
Henrietta, New York 14467

EPA Establishment No. 8780-NY-1
EPA Registration No. 8780-51
DIAZINON® is a trademark of
CIBA-GEIGY Corporation

JAPANESE BEETLE




GRUB




ADULT

CHINCH BUG




CHINCH BUG

BLUEGRASS BILLBUG



LARVA




ADULT

SOD WEBWORM




LARVA




ADULT

EUROPEAN CHAFER



GRUB



ADULT

TURF LINE DIAZINON 5G LAWN INSECT CONTROL KILLS:

ANTS
ARMYWORMS
BERMUDA MITES
BILLBUGS
BROWN DOG TICKS
CHIGGERS
CHINCH BUGS
CLOVER MITES

CRICKETS
CUTWORMS
DIGGER WASPS
EARWIGS
FLEAS
LAWN MOTHS
LEAFHOPPERS
MILLIPEDES

SPRINGTAILS
SOD WEBWORMS
SOWBUGS
WHITE GRUBS OF:
*EUROPEAN CHAFER
*JAPANESE BEETLE
*SOUTHERN CHAFER
*See back panel—

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS
AND DOMESTIC ANIMALS
WARNING**

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust and contact with eyes, skin, and clothing. Keep children and pets off treated areas until this product is washed into the soil and the grass is dry.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink one or two glasses of water and induce vomiting by touching the back of the throat with a finger. Do not induce vomiting by giving anything by mouth to an unconscious person. Get medical attention immediately.

If Inhaled: Move victim to fresh air. Apply respiration if necessary.

If on Skin: Wash thoroughly with soap and water.

If in Eyes: Flush eyes with water for at least 15 minutes. If irritation persists get medical attention.

NOTE TO PHYSICIAN: If symptoms of cholinergic poisoning occur, administer atropine therapeutically.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply where runoff is likely to occur. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

OVERALL LAWN APPLICATION

To control the following lawn insects on both grass and dichondra lawns: Chinch bugs, Earwigs, Ants, Lawn moths (Sod webworms), Bermuda mites, Crickets, Armyworms, Brown dog ticks, Chiggers, Clover mites, Cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails (Collembola).

Apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50'). For heavy lawns such as St. Augustine grass, application should be increased to 3 lbs. per 1000 sq. ft. Water lawn lightly after application.

To control the following insects in lawns:

Billbugs, White grubs of European chafer, Japanese beetle, and Southern chafer.

Apply at the rate of 2½ lbs. per 1000 sq. ft. Water lawn thoroughly after application. For grub control make application any time between late July and early October. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

TO CONTROL:

HYPERODES WEEVIL (NEW YORK ONLY)

Apply 2 lbs. per 1000 sq. ft. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

**IT IS RECOMMENDED THAT ALL
APPLICATIONS BE MADE WITH A
FERTILIZER SPREADER**

**SPREADER SETTINGS FOR OVERALL
LAWN APPLICATIONS:**

NOTE: The spreader settings listed are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting if necessary in order to assure the correct rate of application.

Spreader make	Spreader setting number		
	2 lbs. per 1000 sq. ft.	2½ lbs. per 1000 sq. ft.	3 lbs. per 1000 sq. ft.
Scott Model 35	4	4½	5
Cyclone Model B Rotary	4½	4¾	5

SPOT TREATMENT

Ants: Apply 1½ teaspoons over and around each individual ant hill. Digger Wasps: Apply 1½ teaspoons in and around each ground nest opening. Make application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants, Brown dog ticks, Clover mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails (Collembola).

For control of these insects, apply a five-foot band of Turf Line Diazinon 5G Lawn Insect Control next to the foundation of the house. Distribute the granules uniformly using a rate of 3.2 oz. for each 5' by 20' area to be treated.

STORAGE AND DISPOSAL

Store this product in a safe place beyond the reach of children and pets. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not store near other pesticides or seeds. Any excess material should be saved for later use in confirmation with the directions on this package. Do not reuse empty container. Wrap container and put in trash collection.

