


PM 15
8780-51



TM



THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOO FARMS

TURF LINE DIAZINON 5G. LAWN INSECT CONTROL

Treats up to 12,000 square feet for general lawn insect control. Higher rates are for white grubs and for use on heavy lawns. (See directions for use)

25 LBS. NET WEIGHT

ACTIVE INGREDIENT
G. 0-dimethyl 0-(2-diisopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 5.0%
INERT INGREDIENTS 95.0%

**KEEP OUT OF REACH OF CHILDREN
WARNING**
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Manufactured By: HIGH POINT MILLS
a division of High Point Mills Corporation Inc.
1225 Lathrop Station Road
Hawthorne, New York 14457

EPA Establishment No. 6780-NY-1
EPA Registration No. 6780-51

JAPANESE BEETLE		CHINCH BUG	BLUEGRASS BILLBUG	
				
GRUB	ADULT	CHINCH BUG	LARVA	ADULT
SOD WEBWORM		EUROPEAN CHAFER		
				
				
ADULT		GRUB ADULT		

TURF LINE

TM

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOG FARMS

TURF LINE DIAZINON 5G. LAWN INSECT CONTROL

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS WARNING

Avoid breathing dust and contact with eyes, skin and clothing. Keep children and pets off treated areas until the product is washed into the soil and the grass is dry.

STATEMENT OF PRACTICAL TREATMENT
If SWALLOWED: Drink one or two glasses of water and induce vomiting by touching the back of the throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

If INHALED: Move victim to fresh air. Apply respiration if necessary.

If on SKIN: Wash thoroughly with soap and water.

If in EYES: Flush eyes with water for at least 15 minutes. If children persons get medical attention.

NOTE TO PHYSICIANS: If symptoms of cholinergic poisoning occur, administer atropine intravenously.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply where runoff is likely to occur. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

OVERALL LAWN APPLICATION

To control the following lawn insects on both grass and sclerophyllous leaves: Chinch Bugs, Earwigs, Ants, Lawn moths (Red & Green), Brownish Meas, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Carpet Weas, Caterpillars, Fleas, Leafhoppers, Sawflies, etc., and Springtails (Collembola).

Apply at the rate of 2 lbs. per 1000 Sq. Ft. (or area 20 x 50). For heavy infestations such as St. Augustine grass, application should be increased to 3 lbs. per 1000 Sq. Ft. Water lawn lightly after application.

To control the following insects in the house:

Blattels, White grubs of European, Chorio, Japanese Beetles & Southern Chorio.

Apply at the rate of 2 x 1 lbs. per 1000 Sq. Ft. Water lawn thoroughly after application. For grub control, hand application only from late July and early October. For Bling control, apply when activity is first observed or when chorio or other grubs indicate damage from these insects. Repeat as necessary.

TO CONTROL:

HYPERODES WISEVI (NEW YORK ONLY)

Apply 2 lbs. per 1000 Sq. Ft. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

IT IS RECOMMENDED THAT ALL APPLICATIONS BE MADE WITH A FERTILIZER SPREADER.

SPREADER SETTINGS FOR OVERALL LAWN APPLICATIONS:

NOTE: The spreader settings listed are approximate. Automatically weigh a few pounds into your spreader and apply to a measured area. Then adjust your spreader setting if necessary in order to ensure the correct rate of application.

Spreader make	Spreader setting number		
	2 lbs. per 1000 sq. ft.	2 1/2 lbs. per 1000 sq. ft.	3 lbs. per 1000 sq. ft.
Scott Model 25	4	4 1/2	5
Cyprus Model B Heavy	4 1/2	4 1/2	5

SPOT TREATMENT

Ants: Apply 1 1/2 teaspoons over and around each individual ant hill. Dig out the nest. Apply 1 1/2 teaspoons in and around each ground nest opening. Most applications should be made when the ants are not active.

BARB TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Control of these insects may enter the house.

Ants, Brown dog ticks, Chorio, Crickets, Earwigs, Fleas, Meas, Sawflies, Springtails (Collembola).

For control of these insects, apply a two-foot band of Turf Line Diazinon 5G Lawn Insect Control near the foundation of the house. Distribute the granules uniformly using a rate of 3.2 oz. for each 5 x 20 area to be treated.

STORAGE AND DISPOSAL

Store this product in a safe place beyond the reach of children and pets. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not store near other pesticides or poisons. Any device material should be used for later use in conjunction with directions on this package. Do not reuse empty container. Wrap container and put in trash collection.