

Reg # 8780-55

PM-13

Treats up to ^{1 1/2} 2
5,000 SQ. FT.
[50'x100']

BEST AVAILABLE COPY

15 LBS.
NET WT.



TM

ACCEPTED

FEB 16 1993

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticides
 registered under
 EPA Reg. No. 8780-55

**TURF LINE DIAZINON
 LAWN INSECT CONTROL PLUS FERTILIZER.**

Treats up to 5000 Square Feet

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS
 For outdoor use on the home lawn

DIAZINON LAWN INSECT CONTROL KILLS:

Ants	Chiggers	Earwigs	Millipedes
Armyworms	Chinch Bugs	European Chafar Grubs	Springtails
European Weevils	Clover Mites	Flies	Red Worms
Billbugs	Crickets	Japanese Beetle Grubs	Thrips
Brown Dog Ticks	Cutworms	Leafhoppers	Southern Chafar Grubs

Ready-to-use Granules. Contains 3.3% diazinon insecticide.
Net Weight 15 pounds

Active Ingredient:

Diazinon 6,6-Diethyl 6-(2-propyl 1,4-methyl-4-pyridinyl)
 phosphorothioate 3.3%

Inert Ingredients: 96.7%

Total Ingredients: 100.0%

GUARANTEED ANALYSIS

Total Nitrogen (N) 22.0%

Available Phosphoric Acid (P₂O₅) 4.0%

Soluble Potash (K₂O) 0.5%

Magnesium (Mg) 0.5%

Sulfur (S) 1.5%

Iron (Fe) 0.5%

KEEP OUT OF REACH OF CHILDREN.
CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Manufactured By: **HIGH POINT MILLS**
 A Division of Progressive Lawn Research, Inc.
 Henrietta, NY 14467

EPA Reg No 8780-55
 EPA Est No 8780-NY-1



TM

BEST AVAILABLE COPY

Treats up to ²⁹ 2
5,000 SQ. FT.
[50'x100']
15 LBS.
NET WT.

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of High Point Mills or the seller. All such risks shall be assumed by the Buyer.

High Point Mills warrants that this product conforms to the Chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. High Point Mills makes no other express or implied warranty of Fitness of Merchantability or any other express or implied warranty. In no case shall High Point Mills or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. High Point Mills and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of High Point Mills.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

OVERALL LAWN APPLICATION

To control the following lawn insects on both grass and dichondra lawns: Chinch Bugs, Earwigs, Ants, Lawn Moths (Sod Webworms), Bermudagrass Mites, Cutworms, Fleas, Leafhoppers, Sodbugs, Millipedes, and Springtails (Collembola).

Precautions: Birds, especially waterfowl feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping treated areas. Where thorough wetting is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected to enter the treated area after treatment, (except spot treatments) water lawn with at least 1/4 inch of water immediately after applying the product, however, stop watering before puddling occurs.

Apply at the rate of 3 lbs. per 1,000 sq. ft. (an area 20' X 50'). Apply when grass is dry. Water lawn lightly after application.

To control the following insects in the lawn:

Billbugs, White Grubs of European Chaffer, Japanese Beetles, & Southern Chaffer

Apply at the rate of 3 lbs. per 1,000 sq. ft. (an area 20' X 50'). Water lawn thoroughly after application. For grub control make application any time between late July and early October or between late March and early May. For Billbug control, apply when activity is first observed or when chawed or brown grass indicates damage from these insects. Repeat as necessary.

To Control HYPERODES WEEVIL:

Apply 3 lbs. per 1,000 sq. ft. Water grass thoroughly after application. Treat problem areas in Mid-April and again Mid-May.

IT IS RECOMMENDED THAT ALL APPLICATIONS BE MADE WITH A FERTILIZER SPREADER

SPREADER SETTINGS FOR OVERALL LAWN APPLICATIONS:

NOTE: The spreader settings listed are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust your spreader settings if necessary so that the desired rate is obtained.

SPREADER NAME SPREADER SETTING NUMBER To deliver 3 lbs. per 1,000 sq. ft.)

Scott Precision Flow Spreader Model PF-4	3.25
Cyclone Model Bt Rotary	2.25
Scott Rotary	A
Central Imperial	3.25
Central Rotary	1.80

STORAGE AND DISPOSAL

Storage: Store in original container in a cool, dry, well-ventilated area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. Do not reuse bag. Discard bag in trash.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash.

PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin, or clothing. Wash thoroughly after handling and before eating or using. May cause contact sensitization following repeated contact with skin of susceptible individuals. If sensitization reactions result, contact a physician. Do not store near food or feed products. Do not contaminate ornamental fish ponds. Do not allow children or pets onto the treated area until granules have been watered in and the grass has dried.

Statement of Practical Treatment

If swallowed, call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or comatose.

If inhaled, remove victim to fresh air and apply artificial respiration if indicated.

If on skin, wash promptly with soap and water. Rinse thoroughly.

If in eyes, rinse eyes with plenty of water and call a physician immediately.

Note to Physicians:

This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidote. 2-PAM is also antidote and may be administered, but only in conjunction with atropine.

Environmental Hazards

This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of granules. For lawns where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected to enter the treated area after treatment, (except spot treatments) water lawn with at least 1/4 inch of water immediately after applying the product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to untreated areas below the mean high water mark. Punctil may be hazardous to a public organism in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.