

GENERAL INFORMATION

HABCO BROMEX is a granular semi-soil sterilant non-selective weed killer containing 4% Diuron and 2% Bromacil. This product is designed for use on railroads and individual sites and other similar areas to kill weeds and grasses. The product has residual weed control properties when applied according to directions.

DIRECTIONS FOR USE

HABCO BROMEX is effective in controlling Paragrass, Maidencane grass, Crabgrass, Water grass, Bermuda grass, and certain broad-leafed weeds such as Primrose Willow, Sea Myrtle, Pigweed, Purslane, Spanish Needle, Ragweed, Chickweed, Wild Mustard, Annual Morning Glory, and Lambs' quarters.

GENERAL RECOMMENDATIONS: May be used on non cropland areas such as railroad rights-of-way, along pipe lines, around telephone and power poles and other such areas where total plant kill is desired.

RECOMMENDATIONS: For general weed control, spread evenly by mechanical spreader as follows: For Initial Application in Southern areas with long

CAUTION

MAY CALLE IRREPARENT OF EVENING OF THE WAT AND KIN AVE EDB BORD BUT AF SECOND which is the work, remains an equ

IMPORTANT

DO NOT APPLY on or near desirable trees or other plants, or on other areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do NOT use on Lawns, Walks, Driveways, Tennis Courts, and other similar areas. DO NOT apply near desirable plants. DO NOT contaminate irrigation or domestic waters. Keep from contact with fertilizers, insecticides, fungicides, and seeds. Thoroughly clean application equipment after use. DO NOT RE-USE CONTAINER. DESTROY BY BURYING IN A SAFE PLACE. DO NOT BURN Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ACTIVE INGREDIENTS

Diuron (3-(3, 4-Dichloropheny 1)-1, 1-dimethylurea) Bromaci! (5-bromo-3-sec-butyl-6--methyluracil) INERT INGREDIENTS

CAUTION: Keep out of the reach of Children. See other cautions on this label.

DPART B

E.F.A. EST 2342 FL 1

EPA PED NO SHEEK

Products for Kailroads and Industry

BROMEX A Semi-Soil Sterilant Type Weed Killer

Product No. 20-738981

2.0% 94.0% Total --- 100.0 %

4.0%

growing season, apply 400 to 800 lbs. per acre (1 to 2 lbs. per 100 sq. ft.).)

In Northern areas with shorter growing season, apply 200 to 800 lbs. per acre ($\frac{1}{2}$ to 2 lbs. per 100 sq. ft.).

For Follow Up Treatments: Northern and Southern areas — apply 200 to 600 lbs. per acre ($\frac{1}{2}$ to $1\frac{1}{2}$ lbs. per 100 sq. ft.).


The amount to be used will depend on the type of weeds present, the degree of control and duration of effect required. These depend on soil type, rainfall and other conditions.

WHEN TO APPLY: May be applied at any time of the year, except when ground is frozen, as long as adequate moisture is supplied, either by rainfall or irrigation. Best results are obtained on annual and perennial weeds in high rainfall areas when applied just before weed growth begins.

To avoid fire hazard from dead and dry vegetation, apply when weeds and grass are small.

> 1997 - 1997 - 🛃 🕺 CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufac turer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.



GENERAL INFORMATION

HABCO BROMEX is a granular semi-soil sterilant non-selective weed killer containing 4% Diuron and 2% Bromacil. This product is designed for use on railroads and individual sites and other similar areas to kill weeds and grasses. The product has residual weed control properties when applied according to directions.

DIRECTIONS FOR USE

HABCO BROMEX is effective in controlling Paragrass, Maidencane grass, Crabgrass, Water grass, Bermuda grass, and certain broad-leafed weeds such as Primrose Willow, Sea Myrtle, Pigweed, Purstane, Spanish Needle, Ragweed, Chickweed, Wild Mustard, Annual Morning Glory, and Lambs' quarters.

GENERAL RECOMMENDATIONS: May be used on non-cropland areas such as railroad rights-of-way, along pipe lines, around telephone and power poles and other such areas where total plant kill is desired.

RECOMMENDATIONS: For general weed control, spread evenly by mechanical spreader as follows: For Initial Application in Southern areas with long

CAUTION

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT, AND SKIN. Avoid breathing dust. Avoid contect with skin, eyes, and clothing.

IMPORTANT

DO NOT APPLY on or near desirable trees or other plants, or on other areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do NOT use on Lawns, Walks, Driveways, Tennis Courts, and other similar areas. DO NOT apply near desirable plants. DO NOT contaminate irrigation or domestic waters. Keep from contact with fertilizers, insecticides, fungicides, and seeds. Thoroughly clean application equipment ofter use. CO NOT RE-USE CONTAINER. DESTROY BY BURYING IN A SAFE PLACE, DO NOT BURN.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ACTIVE INGREDIENTS

Diuron (3-(3, 4-Dichloropheny 1)-1,	
1-dimethylurea)	4.0%
Bramacil (5-bromo-3-sec-butyl-	
6methyluracil)	2.0%
INERT INGREDIENTS	94.0%
Total -	- 100.4%

CAUTION: Keep out of the reach of Children.

See other cautions on this label.

E P.A. REG. NO 5823 38

growing season, apply 400 to 800 lbs. per acre (1 to 2 lbs. per 100 sq. ft.).)

In Northern areas with shorter growing season, apply 200 to 800 lbs, per acre (1/2 to 2 lbs, per 100 sg. ft.).

For Follow Up Treatments: Northern and Southern areas - apply 200 to 600 lbs. per acre (1/2 to 1 1/2 lbs. per 100 sq. ft.).

The amount to be used will depend on the type of weeds present, the degree at control and duration of effect required. These depend on soil type, rainfall and other conditions.

WHEN TO APPLY: May be opplied at any time of the year, except when ground is frozen, as long as adequate moisture is supplied, either by rainfall or irrigation. Best results are obtained on annual and perennial weeds in high rainfall aceas when applied just before weed growth begins.

To avoid fire hazard from dead and dry vegetation, apply when weeds and grass are small.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no worranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read oll directions carefully.

