

435763

Ms. Kathie Tryson
Spectrum Group
Division of United Industries Corp.
P.O. Box 15842
St. Louis, MO 63114

JAN 25 1993

Dear Ms. Tryson:

Subject: Amendment - Change to "Caution" Formula
Spectracide Lawn and Garden Insect Control
EPA Reg. No. 8845-92
Your submission dated July 24, 1992

The amendment referred to above, submitted in connection with a registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable provided the following revisions are made prior to release for shipment under the amended labeling:

- 1) The prohibition "This product must not be used on golf courses and sod farms" must appear at the top of the front panel. At its broadest interpretation, this would mean at least in the top half of the label. For your label that would be above "WIDE RANGE OF CONTROL". You may choose to relocate the prohibition to beneath the product name, or else above or directly below "Protects lawns from grubs and other insects" (provided it is still in the top half of the label). You may wish to relocate the front panel limitation to home lawn and home garden use only to appear with the golf course/sod farm prohibition, wherever you relocate it, although such placement is not required.
- 2) On both the container label and the booklet label, revise the footnote under the ingredients statement to read as follows:

* Contains aromatic petroleum distillates
- 3) For consistency with the capitalization of the other multi-part pest names on the label, for the crop "BEANS", the pest name should appear as "Cucumber Beetles", rather than "Cucumber beetles".

A stamped copy of the label is enclosed for your records. The replacement Confidential Formula (CF) dated July 24, 1992 is acceptable and has been added to this product's registration file,

NOTE - The label and associated Confidential Statement of Formula accepted at this time supersede all previous labels and formula statements on file for this product.

Sincerely yours,

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (H7505C)

Spectracide® Lawn & Garden Insect Control CONCENTRATE

PROTECTS MANY AREAS
**PROTECTS: Lawns, Vegetables, Fruits & Nuts, Roses,
Flowers, Trees, Shrubs**

Protects lawns from grubs and other insects

WIDE RANGE OF CONTROL
**CONTROLS: Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants¹,
Flea Beetles, Grubs, Mites, Sod Webworms and many others**
¹Aids in control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate • Contains diazinon insecticide
Diazinon Concentrate

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-
4-pyrimidinyl) phosphorothioate 25.0%

Inert Ingredients * : 75.0%

* Contains xylene range aromatic solvent

KEEP OUT OF REACH OF CHILDREN!

CAUTION: See side panel for additional precautionary statements

See full label in booklet attached to bottle

Net Contents: 1 Pint

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 25 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
8845-92

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information: Spectracide Lawn & Garden Insect Control is an emulsifiable concentrate product which must be diluted with water before application. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits and insects which

annoy pets and
LET FOR COM
AND PESTS O

Notes: (1) USE
AFTER OPENING
POSITIVE DIREC
home. (3) Do not
cus, papaya, p
plant injury ma
succulent peas
dried pea or be
food crops grow

After spraying, wait the following number of days before harvesting

1 day	Tomatoes
3 days	Melons, Winter Squash
5 days	Peppers, Strawberries
7 days	Beans, Broccoli, Blackberries, Boysenberries, Dewberries, Loganberries, Peas, Raspberries
10 days	Chinese Broccoli, Chinese Cabbage, Collards
14 days	Carrots, Endive (Escarole), Lettuce, Onions, Radishes, Spinach, Turnips
21 days	Apricots, Cabbage, Cherries, Nectarines, Peaches
28 days	Grapes
35 days	Potatoes (Irish)
45 days	Walnuts
—	Almonds (one dormant spray only)

Precautions: (1) Wear a long-sleeved shirt and long-legged pants with the wind to your back. (2) Do not spray on windy days. (3) If clothing becomes soiled, use, remove clothing after spraying, wash affected body areas thoroughly, and launder clothing before wearing again.

BEST AVAILABLE COPY

Problem Area	Pests*
Nuisance Pests in Outside Areas	Ants, Brown Dog Ticks, Chiggers, Cockroaches, Crickets, Fleas, Fire Ants, Spiders and others*
Lawns	Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs and others*
Vegetables: Foliar Application	
Beans, Peas (except dried varieties)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites
Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale	Aphids, Flea Beetles, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms
Endive (Escarole), Lettuce, Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites
Peppers, Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms (Tomatoes), Flea Beetles, Leafhoppers, Southern Armyworms (Tomatoes), Vinegar Flies - <i>Drosophila</i> spp. (Tomatoes)
Onions (Bulb and Green)	Onion Thrips
Radishes, Turnips, Carrots, Parsnips, Potatoes	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Colorado Potato Beetles (Potatoes), Southern Armyworms (Potatoes)
Sweet Corn	Aphids, Flea Beetles, Grasshoppers, Spider Mites
Melons, Squash (Summer, Winter)	Aphids, Thrips, Spider Mites
Vegetables: Preplant Soil Application	
Beans, Peas (except dried varieties)	Cutworms, Mole Crickets, Wireworms (Snap Beans)
Beets, Red (Table)	Cutworms, Mole Crickets
Cabbage	Cutworms, Root Maggots, Wireworms
Carrots, Sweet Corn	Cutworms, Mole Crickets, Wireworms

Lettuce, Tomatoes	Cutworms, Wireworms
Radishes, Turnips	Cutworms
Fruits & Nuts	
Almonds (California only)	Scale Crawlers
Walnuts (California only)	Aphids, Scale Crawlers, Spider Mites and others*
Apricots, Peaches, Nectarines	Aphids, Apricot Mealybugs (Apricots), Spider Mites, Scale Crawlers
Cherries	Black Cherry Aphids, Cherry Fruitflies, Cherry Rust Mites, Scale Crawlers and others*
Plums, Prunes	Aphids, Spider Mites
Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Aphids, Leafhoppers, Two-spotted Spider Mites, Thrips, Raspberry Fruitworms
Strawberries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers
Grapes	Aphids, Grape Berry Moths, Grape Leafrollers, Leafhoppers
Roses & Flowers	
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Rhododendron, Roses	Aphids, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Two-spotted Spider Mites, Thrips, Whiteflies and others*
Ornamental Trees & Shrubs	
Arbovitae, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Spruce, Willow	Aphids, Mites, Scale Crawlers, Whiteflies, Tent Caterpillars and others*

*This table is only a partial listing of the pests controlled by this product. See attached booklet for complete directions for use and pests controlled.

BEST AVAILABLE COPY

4
2
17

STORAGE AND DISPOSAL

Storage: Do not store near heat or open flame. Store in original container in an area inaccessible to children or pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.** **Disposal:** Do not reuse empty container. Thoroughly rinse container, securely wrap in several layers of newspaper, and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate or contaminate foodstuffs. Food utensils such as spoons and measuring cups must be cleaned for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. **If inhaled:** Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention. **If on skin:** Wash promptly with soap and water. Rinse thoroughly. Get medical attention. **If in eyes:** Rinse eyes with plenty of water and call a physician immediately. **Note to Physician:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is anticholinergic. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast

waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if water flow, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Do not apply to lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call toll free 800-332-5553 weekdays 9-5 Central Time

Manufactured in the USA for Spectrum Group, Division of United Industries Corporation, St. Louis, MO 63114-5842

EPA Reg. NO. 8845-92
EPA Est. No. 9688-MO-1

Form: A1/Reg ©1992

BEST AVAILABLE COPY

5
9
17

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Spectracide Lawn and Garden Insect Control is an emulsifiable concentrate product which must be diluted with water before application. Refer to the section How to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Spectracide works fast and applications should be repeated only as directed to maintain control.

- Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do not apply to food crops grown in greenhouses.

- Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

ACCEPTED with COMMENTS in EPA Letter Dated

JAN 25 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 8845-92

Spectracide® Lawn & Garden Insect Control CONCENTRATE

PROTECTS MANY AREAS PROTECTS: Lawns, Vegetables, Fruits & Nuts, Roses, Flowers, Trees, Shrubs

Protects lawns from grubs and other insects

WIDE RANGE OF CONTROL CONTROLS: Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants, Flea Beetles, Grubs, Mites, Sod Webworms and many others Aids in control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate • Contains diazinon insecticide Diazinon Concentrate

Active Ingredient: Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 25.0% Inert Ingredients*: 75.0% * Contains xylene range aromatic solvent

KEEP OUT OF REACH OF CHILDREN CAUTION:

See end of booklet for additional precautionary statements

Net Contents: 1 Pint

How to Mix and Apply

Step #1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

Step #2: Use clean sprayer. Carefully measure and mix the amount of Spectracide Lawn & Garden Insect Control and water as suggested in the rate chart.

Step #3: Spray as directed in rate chart. Thorough coverage is important. Flush sprayer with clean water after each use. See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Measurement Conversions

- 1 tablespoon = 3 teaspoons 1 fl. oz. = 6 teaspoons or 2 tablespoons 6 fl. oz. = 3/4 cup 1 1/2 tablespoons in 3 gals. of water = 1 1/2 teaspoons in 1 gal. of water 2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

Lawn Pest Control

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e. ducks or geese can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustinegrass.

To control pests listed below, apply recommended rates.

BEST AVAILABLE COPY

6 9 17

Nuisance Pests Outside the Home	
Pests	Remarks
Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola)	<p>Mix 6 fl. oz. in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.</p> <p>To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary.</p> <p>Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.</p>
Fire Ants	<p>To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>NOTE: For best results, apply in cool weather 65-80°, or in early morning or late evening hours. Treat new mounds as they appear.</p>

Insect Control on Ornamental Trees, Shrubs and Flowers			
Trees and Shrubs	Insects	Rate per 3 Gals. of Water	Timing/Special Directions
Arborvitae, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Spruce, Willow	Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Obscure Root Weevils, Omnivorous Leafiers, Scale Crawlers (Cottonycushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft)	2 tablespoons	Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray.
	Webworms (Juniper, Mimosa), Whiteflies	2 tablespoons	Apply as a thorough foliar spray. Mimosa Webworms: apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.
	Apple-and-Thorn Skeletonizers, Hemlock Chermes, Oak Loopers, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars, Webworms (Cottoncater, Fall)	2 tablespoons	Apply as a thorough foliar spray.

BEST AVAILABLE COPY

8
2
17

Roses and Flowers	Insects	Rate Per 3 Gals. of Water	Timing/Special Directions
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Rhododendron, Roses	Aphids, Euonymus Scale Crawlers, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies	2 tablespoons	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.

Fruits and Nuts	
Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted. Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.	
Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums	21
Prunes	21
Raspberries	7
Strawberries	5
Walnuts	15

* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.
 ** Dormant spray only.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

Crop	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose scale, Parlatoria scale	2 tablespoons in 3 gals. of water	Apply only as a dormant spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard.
NOTE: This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.			
Apricots	Aphids, Apricot Mealybugs, Mites (Brown Two-spotted Spider), Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 5 gallons per 500 sq. ft. of field. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest.
NOTE: This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states.			
Cherries	Black Cherry Aphids, Cherry Fruitflies, Cherry Fruitworms (Maggots), Fruittree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.

BEST AVAILABLE COPY

9
8
17

Crop	Pest	Rate	Timing/Special Directions
Grapes	Aphids, Leafhoppers, Grape Berry Moths, Grape Leafhoppers	4 1/2 teaspoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of vineyard. Do not make more than 5 applications per season a minimum of 7 days apart. Do not apply within 28 days of harvest.
Nectarines	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Plums and Prunes	Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 500 sq. ft. of orchard. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 21 days of harvest.

Crop	Pest	Rate	Timing/Special Directions
Strawberries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 500 sq. ft. of field. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 6 gallons per 500 sq. ft. of orchard. Do not make more than 3 applications per season 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest.
NOTE: This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.			

BEST AVAILABLE COPY

10 2 17

Vegetables: For foliar or soil applications, Spectracide Lawn and Garden Insect Control should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil. Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Interval (PHI*) for Foliar Applications	
Crop	PHI* (Days)
Beans ¹ (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cariboupeas	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Cranshaws	3
Cucumbers	7
Endive (Escarole)	14
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Peas ¹	7
Peppers	5
Potatoes (Irish)	35
Radishes, Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	1
Turnips (Roots and Tops)	14
Watermelons	3

*PHI-Pre-Harvest Interval, or the number of days between the last application and harvest. ¹Do not use this product on dried peas or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas.

Vegetables
For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing/Special Directions
Beans: Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.
	NOTES: (1) Do not apply this product to dried bean varieties such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.		
Beets, Red (Table)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.

BEST AVAILABLE COPY

11 2 17

Crop	Pest	Rate	Timing/Special Directions
Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.

Crop	Pest	Rate	Timing/Special Directions
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Cauliflower	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			

BEST AVAILABLE COPY

12 of 17

Crop	Pest	Rate	Timing/Special Directions
Chinese Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
Collards	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			

Crop	Pest	Rates	Timing/Special Directions
Corn (see sweet corn)			
Cucumbers	Aphids, Thrips, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar insecticides.			

BEST AVAILABLE COPY

13
of
17

Crop	Pest	Rate	Timing/Special Directions
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons	Aphids, Thrips, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Onions (Bulb and Green)	Onion Thrips	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.

Crop	Pest	Rate	Timing/Special Directions
Parsnips	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat applications every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
Peas (succulent only)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
NOTES: (1) Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.			

BEST AVAILABLE COPY

14
8
17

Crop	Pest	Rate	Timing/Special Directions
Peppers	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
Radishes	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

Crop	Pest	Rate	Timing/Special Directions
Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.			
Squash, Summer	Aphids, Thrips, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Squash, Winter	Aphids, Thrips, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.

BEST AVAILABLE COPY

15 of 17

Crop	Pest	Rate	Timing/Special Directions
Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
	NOTE: To protect bees, do not apply to corn foliage during the pollen shed period.		

Crop	Pest	Rate	Timing/Special Directions
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (<i>Drosophila</i> spp.)	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-8 inches.
Turnips and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
	NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.		
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

BEST AVAILABLE COPY

16
2
17

STORAGE AND DISPOSAL**Storage:**

Do not store near heat or open flame. Store in original container in an area inaccessible to children and pets. USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.

Disposal:

Do not reuse empty container. Thoroughly rinse container, securely wrap it in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.


Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions or if waterfowl, i.e., duck or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal area below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters. This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.



NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions. Questions or comments? Call toll free 800-332-6663 weekdays 9-5 Central Time. Manufactured in the USA for Spectrum Group, Division of United Industries Corp., St. Louis, MO 63114-5942. EPA Reg. NO. 8845-92 Item: EPA Est. No. 9808-MO-1 Form: A1/REG 01982

BEST AVAILABLE COPY

17 of 17