

8845-92

2/19/2001

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 11-30-93

4/11



United States Environmental Protection Agency
Office of Pesticide Programs (H7505C)
Washington, DC 20460

Application for Pesticide:

Registration
Amendment
Other

OPP Identifier Number

Section I

1. Company/Product Number 8845-92	2. Product Manager G. LaRocca	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Spectracide Lawn & Garden Insect Control Concentrate	PM# 13	
5. Name and Address of Applicant (include ZIP Code) Spectrum Group Division of United Industries Corp. PO Box 15842 St. Louis, Missouri 63114-0842 <input type="checkbox"/> Check if this is a new address	6 Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labelling to: EPA Reg. No. _____ Product Name _____	

Section II

Amendment - Explain below
 Resubmission in response to Agency letter dated _____
 Notification - Explain below.

Final printed label in response to Agency letter dated _____
 "ME TOO" Application.
 Other - explain below

NOTIFICATION

FEB 19 2001

Explanation: Use additional page(s) if necessary. (For section I and Section II)

Notification under PR Notice 98-10 of optional marketing claim "GreenThumb Satisfaction Guarantee" with explanatory statement.

Section III

1. Material in Which This Product Will Be Packaged:				2. Type of Container	
<input type="checkbox"/> Yes * <input type="checkbox"/> No * Certification must be submitted.	<input type="checkbox"/> Yes <input type="checkbox"/> No If "YES," Unit package wgt _____ No. per container _____	<input type="checkbox"/> Yes <input type="checkbox"/> No If "YES," Package wgt _____ No. per container _____	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label Is Affixed To Product <input type="checkbox"/> Lithographed <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section IV

1. Contact Point (Complete items directly below for identification of the individual to be contacted, if necessary, to process this application.)		
Name Glenda L. Mah	Title Regulatory Affairs Specialist	Telephone No. (include Area Code) 800-242-1166 Ext. 3415
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate, and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6 Date Application Received (Stamped)
2. Signature <i>Glenda L. Mah</i>	3. Title Regulatory Affairs Specialist	
4. Typed Name Glenda L. Mah	5. Date 2-14-01	



Lawn & Garden Insect Control CONCENTRATE

This product must not be used on golf courses and sod farms.
For outdoor use on the home lawn and the home garden only.

[PROTECTS MANY AREAS: [PROTECTS] Lawns, Vegetables, Fruits & Nuts, Roses, Flowers, Trees, Shrubs • Protects lawns from grubs and other insects • Controls/kills lawn & garden insects and soil pests

WIDE RANGE OF CONTROL: [CONTROLS] Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants¹, Flea Beetles, Grubs, Mites, Sod Webworms & many others

Controls tough outdoor insects • Kills lawn and garden insects ... including grubs, chinch bugs, fleas, sod webworms, aphids, beetles, whiteflies, ants, mites, brown dog ticks and other unwanted insects • Kills grubs • Connect and Spray (Optional for hose-end sprayer) • Spray concentrate • Contains diazinon insecticide • Diazinon Concentrate • Diazinon Insect Spray • Concentrated Insect Spray • Kills 90+ Pests • *Guaranteed Results or Your Money Back • GreenThumb® Satisfaction Guarantee ¹Aids in control]

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 25.0%

Inert Ingredients 75.0%

* Contains aromatic petroleum distillates

KEEP OUT OF REACH OF CHILDREN

CAUTION See end of booklet for additional precautionary statements

(Optional for Roblet Label: See back booklet for additional precautionary statements)

NET CONTENTS 1 PINT

[OPTIONAL PRODUCT FACT BOX]

PRODUCT FACTS

What It Does:

Kills Grubs, Ants, Fleas, Chinch Bugs, Sod Webworms, Aphids, Beetles, Whiteflies, Mites, Brown Dog Ticks and Other Listed Insects

Where to Use: (icon outdoors)

Questions & Comments: (phone icon) (computer icon)

Call 1-800-332-5553 or Visit our Website at www.spectracide.com

NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

STOP. (optional) READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, PLANT INJURY, OR ILLEGAL RESIDUES.

GENERAL INFORMATION

Spectracide Lawn and Garden Insect Control is an emulsifiable concentrate product which must be diluted with water before application. Refer to the section **How to Mix and Apply**. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy people and pets. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Spectracide works fast and applications should be repeated only as directed to maintain control.

Notes:

- (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS.
- (2) Do not use inside the home.
- (3) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since injury may occur.
- (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties.
- (5) Do not apply to food crops grown in greenhouses.

Precautions:

- (1) *Wear a long-sleeved shirt and long-legged pants when spraying.*
- (2) *Spray with the wind to your back.*
- (3) *Do not spray on windy days.*
- (4) *If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.*

FEB 19 2001

NOTIFICATION

2/11

HOW TO MIX AND APPLY

Step #1: Determine size of spraying job. For lawns, fruits, nuts and vegetables, measure the area to be sprayed. For trees, shrubs and flowers, apply as thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.
Step #2: Use clean sprayer. Carefully measure and mix the amount of Spectracide Lawn & Garden Insect Control and water as indicated in the rate chart.
Step #3: Spray as directed in rate chart. Thorough coverage is important.
Step #4: Flush sprayer with clean water after each use.

See chart for complete list of use sites, plants and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or plant you are spraying.

MEASUREMENT CONVERSIONS

1 tablespoon = 3 teaspoons 1 fl. oz. = 6 teaspoons or 2 tablespoons 6 fl. oz. = 3/4 cup
 1 1/2 tablespoons in 3 gals. of water = 1 1/2 teaspoons in 1 gal. of water
 2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

LAWNS

PRECAUTIONS: *Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.*

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustine grass.

FOR INSECT CONTROL: Apply as indicated below. Use 1-1/2 tablespoons in 3 gallons of water to cover 125 sq. ft. or 6 fl. oz. in 24 gallons of water to cover 1,000 sq. ft.

SURFACE INSECTS (icon)

Including: Ants, armyworms, bermudagrass mites, brown dog ticks, chiggers, chinch bugs, clover mites, crickets, cutworms, earwigs, fleas, fruit flies, lawn moths (sod webworms), leafhoppers, millipedes, rhodesgrass scales, sowbugs and springtails (collembola) Thoroughly wet down grass a few hours before applying. If chinch bugs in the North are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gallons of water to cover 125 sq. ft. or 4 fl. oz. in 24 gallons of water to cover 1,000 sq. ft. For fruit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.

UNDERGROUND INSECTS (icon)

Including: Billbugs, white grubs (Japanese beetle, European chafer, Southern chafer), dung beetle and hyperodes weevils. Water grass thoroughly after application.

WHEN TO TREAT: Treat the following pests during the months indicated or when first noticed.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Billbugs

Grubs

Dung beetle grubs

Hyperodes weevils

Treat Ants and other insects when they first appear or you notice lawn damage.

NUISANCE PESTS OUTSIDE THE HOME

Pests	Directions
Ants, Boxelder bugs, Brown dog ticks, Chiggers, Clover mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola)	Mix 2 fl. oz. in 1 gallon of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide. To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2 - 3 feet. Repeat application as necessary. Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.
Fire ants	To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gallon of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. NOTE: For best results, apply in cool weather 65 - 80 °, or in early morning or late evening hours. Treat new mounds as they appear.

W
11

ORNAMENTAL TREES, SHRUBS AND FLOWERS Spray when listed pests first appear. Mix 2 teaspoons in 1 gallon of water. Apply as a thorough foliar spray. Wet leaves and branches to the point of dripping. Try to spray underside of leaves and penetrate dense foliage. Repeat application as necessary.

TREES AND SHRUBS	
Plants	Pests
Arborvitae, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Spruce, Willow	Aphids, Apple-and-Thorn skeletonizers, Bagworms, Dipterous leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Hemlock chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European red, Privet, Two-spotted spider), Moths (European pine shoot, Holly bud), Oak looppers, Obliquebanded leafrollers, Obscure root weevils, Omnivorous leafhoppers, Scale crawlers (Cottonycushion, Euonymus, Lecanium, Pine needle, Pit, San Jose, Soft), Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies NOTE: For Mimosa Webworms, apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.
ROSES AND FLOWERS	
Plants	Pests
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Rhododendron, Roses	Aphids, Euonymus scale crawlers, Flea beetles, Dipterous leafminers, Leafhoppers, Mealybugs, Mites (Carnation bud, Carnation shoot, cyclamen, European red, Two-spotted spider), Thrips, Whiteflies

FRUITS AND NUTS Spectracide Lawn and Garden Insect Control should be applied in the amount of water recommended in the table below. Start spraying after blooming. Treat when aphids, mites, and crawler stages of scale insects or other insects are first noticed. Apply as a full cover foliar treatment, thoroughly spraying all parts of the plant. Spray until plants just begin to drip. Observe stated time intervals between last application and harvest, as well as intervals between applications and the total number of applications that can be made per season. NOTE: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST					
Fruits & Nuts	PHI*	Fruits & Nuts	PHI*	Fruits & Nuts	PHI*
Almonds	**	Cherries	21	Peaches	21
Apricots	21	Dewberries	7	Plums	21
Blackberries	7	Grapes	28	Prunes	21
Boysenberries	7	Honeydew Melons	3	Raspberries	7
Cantaloupes	3	Loganberries	7	Strawberries	5
Casaba Melons	3	Muskmelons	3	Walnuts	45
Crenshaw Melons	3	Nectarines	21	Watermelons	3

* PHI = Pre-Harvest Interval. ** Dormant spray only.

Fruits & Nuts	Pests	Rate	Directions
Almonds (CA only)	Scales (Parlatoria, San Jose)	2 teaspoons in 1 gal. of water	Apply only as a dormant spray (before the leaves come out). Do not apply more than 6 gallons per 500 sq. ft. of home orchard.
NOTE: This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.			

Fruits & Nuts	Pests	Rate	Directions
Apricots, Nectarines, Peaches	Aphids, Apricot mealybugs, Mites (Brown Two-spotted spider), Scale crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Do not apply more than 4 gallons per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart.
Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry fruitworms, Thrips, Two-spotted spider mites	2 teaspoons in 1 gal. of water	Do not apply more than 5 gallons per 500 sq. ft. of caneberry patch. Do not make more than 5 applications per season a minimum of 14 days apart.
NOTE: This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states.			
Cherries, Plums, Prunes	Black cherry aphids, Cherry fruitflies, Cherry fruitworms (maggots), Fruit tree leafrollers, Mites (Cherry rust), Scale crawlers (San Jose)	2 teaspoons in 1 gal. of water	Do not apply more than 4 gallons per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 7 days apart.
Grapes	Aphids, Grape berry moths, Grape leaffolders, Leafhoppers	1-1/2 teaspoons in 1 gal. of water	Do not apply more than 3 gallons per 500 sq. ft. of home vineyard. Do not make more than 5 applications per season a minimum of 7 days apart.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons	Aphids, Spider mites, Thrips	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	Repeat application every 7 days up to 5 applications per season.
Strawberries	Aphids, Strawberry leafrollers, Two-spotted spider mites	2 teaspoons in 1 gal. of water	Do not apply more than 3 gallons per 500 sq. ft. of strawberry patch. Do not make more than 3 applications per season a minimum of 7 days apart.

Fruits & Nuts	Pests	Rate	Directions
Walnuts (CA only)	Aphids, Scale crawlers (Walnut, Italian pear, Frosted), Spider mites, Twig borers	2 teaspoons in 1 gal. of water	Do not apply more than 6 gallons per 500 sq. ft. of home orchard. Do not make more than 3 applications per season 14 days apart. Do not apply after husks open or split.
NOTE: This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.			

VEGETABLES For foliar or soil applications, Spectracide Lawn and Garden Insect Control should be applied in the amount of water recommended in the table below. For application to the foliage, thoroughly spray all plant parts. Wet the plant just to the point of dripping. Application to the soil should be mixed into the soil immediately after application. Start treating when insects are first noticed. Observe stated time intervals between last application and harvest as well as intervals between applications and the maximum number of applications per season. Do not apply to vegetables grown in greenhouses. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST					
Vegetables	PHI*	Vegetables	PHI*	Vegetables	PHI*
Beans ¹ (succulent varieties only: Lima, Pole, Snap)	7	Cucumbers	7	Potatoes (Irish)	35
Broccoli	7	Endive (Escarole)	14	Radishes,	14
Cabbage	21	Kale	10	Spinach	14
Carrots	14	Lettuce	14	Squash, Summer	7
Cauliflower	7	Onions (Bulb, Green)	14	Squash, Winter	3
Chinese Broccoli	10	Parsnips	14	Sweet Corn	7
Chinese Cabbage	10	Peas ¹	7	Tomatoes	1
Collards	10	Peppers	5	Turnips	14

*PHI=Pre-Harvest Interval. ¹Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas.

Underground insects (cutworms, mole crickets, root maggots and wireworms) are treated with a soil application which is immediately mixed into the soil at specific depth for each insect. Refer to the following table for the required depth for each insect listed

SOIL INCORPORATION TABLE

Insect	Depth of Mixing Into Soil
Surface cutworms, Root maggots	2-3 inches
Subterranean cutworms	3-6 inches
Mole crickets	1-2 inches
Wireworms	4-8 inches

Vegetables	Pests	Rate	Directions
Beans: Lima, Pole, Snap (succulent varieties only); Peas (succulent only)	Aphids, Beetles (Cucumber, Mexican bean), Grasshoppers, Leafhoppers, Spider mites	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 5 days, up to 3 applications per season.
	Cutworms (surface, subterranean), Wireworms (For peas, snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly just before planting. Use for treating wireworms on peas and snap beans only.
	Mole crickets	2 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly 1-2 days before planting.
NOTES: (1) Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if the beans, peas or weeds in the treatment area are in bloom.			
Beets, Red (Table)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly just before planting.
	Mole crickets	2 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly 1-2 days before planting.

Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage (Bok Choy, Napa), Collards, Cucumbers, Endive (Escarole), Kale, Parsnips, Peppers, Potatoes, Spinach, Squash (Summer, Winter)	Aphids, Armyworms (Southern), Beetles (Colorado potato, Flea) Diamondback moths, Dipterous leafminers, Imported cabbageworms, Leafhoppers, Spider mites, Thrips	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 7 days, up to 5 applications per season.
Cabbage	Aphids, Diamondback moths, Dipterous leafminers, Flea beetles, Imported cabbageworms	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 7 days, up to 5 applications per season.
	Cutworms (surface, subterranean), Root maggots, Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly just before planting.
Carrots, Corn (Sweet corn)	Aphids, Dipterous leafminers, Flea beetles, Grasshoppers, Leafhoppers, Spider mites, Thrips	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 7 days, up to 5 applications per season.
	Cutworms (surface, subterranean), Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly just before planting.
	Mole crickets	2 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly 1-2 days before planting.
NOTE: To protect bees, do not apply to corn foliage during the pollen shed period.			
Lettuce, Tomatoes	Aphids, Armyworms (Fall, Southern), Beetles (Colorado potato, Flea) Dipterous leafminers, Leafhoppers, Spider mites, Vinegar flies (<i>Drosophila spp.</i>)	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 7 days, up to 5 applications per season.
	Cutworms (surface, subterranean), Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly just before planting.
Onions (Bulb, Green)	Onion thrips	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 7 days, up to 3 applications per season.

Radishes	Aphids, Dipterous leafminers, Flea beetles, Imported cabbageworms, Leafhoppers	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 7 days, up to 3 applications per season.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly just before planting.
Turnips	Aphids, Dipterous leafminers, Flea Beetles, Imported cabbageworms, Leafhoppers	2 teaspoons in 1 gal. of water to cover 350 sq. ft. of garden	If necessary, treat every 7 days, up to 5 applications per season.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1-1/2 gals. of water	Spray the soil evenly just before planting.

STORAGE AND DISPOSAL

Storage: Do not store near heat or open flame. Store in original container in an area inaccessible to children and pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.** **Disposal:** Do not pour the concentrate, the spray mixture or the rinse water down the drain or sewer. Do not reuse empty container. Thoroughly rinse container, securely wrap it in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention. If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention. If in eyes: Rinse eyes with plenty of water, and call a physician immediately. **Note to Physician:** This product is an organophosphorus insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction

with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters. This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call 800-332-5553

[*If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]

[Satisfaction Guaranteed: If for any reason you are not completely satisfied with the performance or results of this GreenThumb® product, it will be repaired or replaced free of charge. Simply provide proof of purchase and return the product and/or unused portion to the place of purchase.]

OPTIONAL READY SPRAY NOZZLE DIRECTIONS

Made in the USA for
Spectrum Group, Division of United Industries Corporation,
P. O. Box 15842, St. Louis, MO 63114-0842
EPA Reg. NO. 8845-92 EPA Est. No. 9688-MO-1 8 2000

NOTE: this product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, PLANT INJURY, OR ILLEGAL RESIDUES.

GENERAL INFORMATION

This product is an emulsifiable concentrate product which must be diluted with water before application. Refer to the section *How to Apply*. It kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy people and pets. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Spectracide works fast and applications should be repeated only as directed to maintain control.

- Notes:**
- (1) USE CONTENTS WITHIN 6 MONTHS AFTER FIRST USE. SEE STORAGE AND DISPOSAL DIRECTIONS.
 - (2) Do not use inside the home.
 - (3) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia or orchids since injury may occur.
 - (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties.
 - (5) Do not apply to food crops grown in greenhouses.

- Precautions:**
- (1) Wear a long-sleeved shirt and long-legged pants when spraying.
 - (2) Spray with the wind to your back.
 - (3) Do not spray on windy days.
 - (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

HOW TO APPLY

Determine the size of the spraying job. For lawns, fruits, nuts and vegetables, measure the area to be sprayed. For trees, shrubs and flowers, apply as a thorough cover spray. Begin treating when insects first appear. Applications should be repeated only as directed to maintain control. Apply as directed in the rate chart. Remember, thorough coverage is important. Be sure to follow the important use limitations for the specific fruit or vegetable you are treating, including the amount of spray that can be applied, the interval between treatments and the days to wait before harvest.

USING THE READY SPRAY NOZZLE (Optional Subhead)

1. Shake bottle well before using.
2. Connect garden hose to the ready spray nozzle. Make sure the black water control dial on the nozzle is in the "OFF" position (tab is vertical).
3. Turn on water faucet. Extend hose to the farthest area to be treated and work back towards faucet so that you do not walk over treated areas. Twist the small, star shaped black product control button to align the flat portion with the lock tab. Push the button in to open.
4. To BEGIN spraying, point nozzle towards the area to be treated and turn the black water control dial to the "ON" position (tab is horizontal). Water will automatically mix with the product. Walk at a steady pace while spraying using an even sweeping motion. Refer to site gauge on bottle to determine amount of product being applied.

5. To STOP spraying turn the black water control dial to the "OFF" position. Push the small, black product control button back in to close. Twist the small, star shaped black product control button so that it is no longer aligned with the flat portion of the lock tab. Turn off the water at the faucet. To relieve pressure, turn the black water control dial to the "ON" position until water stops spraying before removing the nozzle from the hose.
6. To STORE unused product, place in a cool area away from heat, direct sunlight or open flame and out of reach of children or pets.

LAWNS

PRECAUTIONS: *Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.*

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion and St. Augustinegrass.

FOR INSECT CONTROL: Apply as indicated below. Use 1 quart to cover 5,332 sq. ft. of lawn (8 fl. oz. covers 1,333 sq. ft.; a pint covers 2,666 sq. ft.)

SURFACE INSECTS (icon)

Including: Ants, armyworms, bermudagrass mites, brown dog ticks, chiggers, chinch bugs, clover mites, crickets, cutworms, earwigs, fleas, fruit flies, lawn moths (sod webworms), leathoppers, millipedes, rhodesgrass scales, sowbugs and springtails (collembola). Thoroughly wet down grass a few hours before applying. If chinch bugs in the North are the only problem, a lower rate of application can be used: 1 quart covers 8,000 sq. ft. (1 pint covers 4,000 sq. ft.). For fruit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.

UNDERGROUND INSECTS (icon)

Including: Billbugs, white grubs (Japanese beetle, European chafer, Southern chafer), dung beetle and hyperodes weevils: Water grass thoroughly after application.

WHEN TO TREAT: Treat the following pests during the months indicated or when first noticed.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Billbugs

Grubs

Dung beetle grubs

Hyperodes weevils

Treat Ants and other insects when they first appear or you notice lawn damage.

ORNAMENTAL TREES, SHRUBS AND FLOWERS

TREES AND SHRUBS	
Arborvitae, Azaleas, Birch, Boxwood, Camellias, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Spruce, Willow	
Insects	Directions
Aphids, Apple-and-Thorn skeletonizers, Bagworms, Hemlock chermes, Dipterous leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European red, Privet, Two-spotted spider), Moths (European pine shoot, Holly bud), Oak loppers, Obliquebanded leafrollers, Obscure root weevils, Omnivorous leafhoppers, Pear slugs, Scale crawlers (Cottoncushion, Euonymus, Lecanium, Pine needle, Pit, San Jose, Soft), Tent caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies	Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray. Mimosa webworms: Apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.
ROSES AND FLOWERS	
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Rhododendron, Roses	
Aphids, Euonymus scale crawlers, Flea beetles, Dipterous leafminers, Leafhoppers, Mealybugs, Mites (Carnation bud, Carnation shoot, Cyclamen, European red, Two-spotted spider), Thrips, Whiteflies	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.

FRUITS AND NUTS Start spraying after blooming. Treat when aphids, mites, and crawler stages of scale insects or other insects are first noticed. Spray until plants just begin to drip. Observe stated time intervals between last application and harvest, as well as intervals between applications and the total number of applications that can be made per season. NOTE: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST					
Fruits & Nuts	PHI*	Fruits & Nuts	PHI*	Fruits & Nuts	PHI*
Almonds	**	Cherries	21	Peaches	21
Apricots	21	Dewberries	7	Plums	21
Blackberries	7	Grapes	28	Prunes	21
Boysenberries	7	Honeydew Melons	3	Raspberries	7
Cantaloupes	3	Loganberries	7	Strawberries	5
Casaba Melons	3	Muskmelons	3	Walnuts	45
Crenshaw Melon	3	Nectarines	21	Watermelons	3

* PHI = Pre-Harvest Interval. ** Dormant spray only.

Fruits & Nuts	Pests	Rate	Directions
Almonds (CA only)	Scales (Parlatoria, San Jose)	1 fl. oz. covers 250 sq. ft (8 fl. oz. covers 2,000 sq. ft.; 1 quart covers 8,000 sq. ft.)	Apply only as a dormant spray (before the leaves come out).
NOTE: This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.			
Apricots, Nectarine, Peaches	Aphids, Apricot mealybugs, Mites (Brown, Two-spotted spider), Scale crawlers (Olive, San Jose)	1 fl. oz. covers 375 sq. ft (8 fl. oz. covers 3,000 sq. ft.; 1 quart covers 8,000 sq. ft.)	Do not make more than 3 applications per season a minimum of 14 days apart.
Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry fruitworms, Thrips, Two-spotted spider mites	1 fl. oz. covers 300 sq. ft. (8 fl. oz. covers 2,400 sq. ft.; 1 quart covers 9,600 sq. ft.)	Do not make more than 5 applications per season a minimum of 14 days apart.
NOTE: This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states.			
Cherries, Plums, Prunes	Aphids (Black cherry, Leafcurl plum, Mealy plum, Thistle), Cherry fruitflies, Cherry fruitworms (maggots), Fruittree leafrollers, Mites (Brown, Cherry rust, European red), Scale crawlers (San Jose)	1 fl. oz. covers 375 sq. ft (8 fl. oz. covers 3,000 sq. ft.; 1 quart covers 12,000 sq. ft.)	Do not make more than 3 applications per season a minimum of 7 days apart.
Grapes	Aphids, Grape berry moths, Grape leafhoppers, Leafhoppers	1 fl. oz. covers 667 sq. ft. (8 fl. oz. covers 5,332 sq. ft.; 1 quart covers 21,332 sq. ft.)	Do not make more than 5 applications per season a minimum of 7 days apart.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons	Aphids, Spider mites, Thrips	1 fl. oz. covers 1,000 sq. ft. (8 fl. oz. covers 8,000 sq. ft.; 1 quart covers 32,000 sq. ft.)	Repeat application every 7 days up to 5 applications per season.

Fruits & Nuts	Pests	Rate	Directions
Strawberries	Aphids, Strawberry leafrollers, Two-spotted spider mites	1 fl. oz. covers 500 sq. ft. (8 fl. oz. covers 4,000 sq. ft.; 1 quart covers 16,000 sq. ft.)	Do not make more than 3 applications per season a minimum of 7 days apart.
Walnuts (CA only)	Aphids, Scale crawlers (Walnut, Italian pear, Frosted), Spider mites, Twig borers	1 fl. oz. covers 250 sq. ft.; (8 fl. oz. covers 2,000 sq. ft.; 1 quart covers 8,000 sq. ft.)	Do not make more than 3 applications per season 14 days apart. Do not apply after husks open or split.
NOTE: This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.			

VEGETABLES Start treating when insects are first noticed. Spray to assure thorough coverage of all parts of the plant. Wet the plant just to the point of dripping. Observe stated time intervals between last application and harvest as well as intervals between applications and the maximum number of applications per season. Do not apply to vegetables grown in greenhouses. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST					
Vegetables	PHI*	Vegetables	PHI*	Vegetables	PHI*
Beans ¹ (succulent varieties only: Lima, Pole, Snap)	7	Cucumbers	7	Potatoes (Irish)	35
Broccoli	7	Endive (Escarole)	14	Radishes,	14
Cabbage	21	Kale	10	Spinach	14
Carrots	14	Lettuce	14	Squash, Summer	7
Cauliflower	7	Onions (Bulb and Green)	14	Squash, Winter	3
Chinese Broccoli	10	Parsnips	14	Sweet Corn	7
Chinese Cabbage	10	Peas ¹	7	Tomatoes	1
Collards	10	Peppers	5	Turnips	14

*PHI=Pre-Harvest Interval. ¹Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas.

Vegetables	Pests	Rate	Directions
Beans: Lima, Pole, Snap (succulent varieties only), Peas (succulent only)	Aphids, Beetles (Cucumber, Mexican bean), Grasshoppers, Leafhoppers, Spider mites	1 fl. oz. covers 1,000 sq. ft. (8 fl. oz. covers 8,000 sq. ft.; 1 quart covers 32,000 sq. ft.)	If necessary, treat every 5 days. Make no more than 3 applications per season
NOTES: (1) Do not use this product on dried pea or bean varieties such as pinto beans, dried limas, split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if the beans, peas or weeds in the treatment area are in bloom.			
Broccoli, Cabbage, Cauliflower, Carrots, Chinese Broccoli, Chinese Cabbage (Bok Choy, Napa), Collards, Corn (Sweet corn), Cucumbers, Endive (Escarole), Kale, Lettuce, Parsnips, Peppers, Potatoes, Spinach, Squash (Summer, Winter), Tomatoes, Turnips	Aphids, Armyworms (Fall, Southern), Beetles (Colorado potato, Flea), Diamondback moths, Dipterous leafminers, Grasshoppers, Imported cabbageworms, Leafhoppers, Spider mites, Thrips, Vinegar flies (<i>Drosophila spp.</i>)	1 fl. oz. covers 1,000 sq. ft. (8 fl. oz. covers 8,000 sq. ft.; 1 quart covers 32,000 sq. ft.)	If necessary, treat every 7 days. Make no more than 5 applications per season.
NOTE: To protect bees, do not apply to corn foliage during the pollen shed period.			
Onions (Bulb, Green)	Onion thrips	1 fl. oz. covers 1,000 sq. ft. (8 fl. oz. covers 8,000 sq. ft.; 1 quart covers 32,000 sq. ft.)	If necessary, treat every 7 days. Make no more than 3 applications per season
Radishes	Aphids, Dipterous leafminers, Flea beetles, Imported cabbageworms, Leafhoppers	1 fl. oz. covers 1,000 sq. ft. (8 fl. oz. covers 8,000 sq. ft.; 1 quart covers 32,000 sq. ft.)	If necessary, treat every 7 days. Make no more than 3 applications per season.

STORAGE AND DISPOSAL

Storage: Be sure the small, star shaped black product control button is in the closed position. Do not store near heat or open flame. Store in original container in an area inaccessible to children and pets. **USE CONTENTS WITHIN 6 MONTHS AFTER FIRST USE.** **Disposal:** Do not pour the concentrate, the spray mixture or the rinse water down the drain or sewer. Be sure the small, star shaped black product control button is in the closed position. Do not reuse empty container. Securely wrap it in several layers of newspaper and discard in trash.

11/11

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds.

Statement of Practical Treatment

If swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. **If inhaled:** Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention. **If on skin:** Wash promptly with soap and water. Rinse thoroughly. Get medical attention. **If in eyes:** Rinse eyes with plenty of water and call a physician immediately. **Note to Physician:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters. This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

11/11