



· · T 1:3

DIRECTIONS FOR RATS Place 4 ounces to 1 pound of Safeguard Brand Mice and Rat Killer in locations frequented by rats. Replace bait as consumed so that feeding is not interrupted for more than 1 day or two at the most. Use no decomposed, mouldy or sour bait. Where a continuous source of infestation prevails from nearby dumps or fields, permanent bait stations should be used and Safequard Brand Mice and Rat Killer replenished as needed. Baiting should continue until complete lack of feeding is noted. This should be from Ten to Fourteen Days. Be sure that sufficient Safeguard Brand Mice and Rat Killer is at hand to complete a continuous 14 day feeding program. If bait is exhausted before rats have been completely eliminated obtain additional Safeguard Brand Mice and Rat Killer promptly. Continuous uninterrupted feeding is what gives you control.

DIRECTIONS FOR MICE. For controlling mice, follow the same general directions as for rats except bait placements of tablespoon amounts (1/4 to 1/2 oz.) of bait should be placed at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 5 days.

CAUTION: Safeguard Brand Mills and Rat should not be ploted will are chuldren or eat it. If this cossible to exists where it connot the reduned in eating the three point of such it with insufer suid repeat left in introphysican m vort t 👎 . . .

USDA Reg. No. 8848-2

MANUF ACTURED BY SAFEGUARD CHEMICAL CORPORATION NEW YORK, N. Y. 1940

NOTE FOR PHYSICIANS: when a human has been known to have ingested Safeguard Brand Mice and Rat Killer, blood

transfusions combined with intravenous injections and oral doses of Vitamin K are indicated us in the case of hemorrhage caused by everdoses of Dicumarol.

ACTIVE INGREDIENTS:
WARFARIN-3-(alpha-
acetonylbenzyl+-4-
hydroxycoumarin 0.025%
INERT INGREDIENTS
100.000%



