

# BLACK JACK LIQUID MULTI-PURPOSE INSECT KILLER

- STOPS BUGS FROM COMING THROUGH SCREENS
- CONTROLS: HOUSE FLIES, MOSQUITOES, GNATS, SPIDERS, ANTS, FLEAS, TICKS, ROACHES, PALMETTO BUGS, SILVERFISH, CRICKETS
- CONTROLS SPIDERS IN THE HOME
- WATER BASED FORMULATION
- FOR DOMESTIC USE BOTH INDOORS AND OUTDOORS
- A READY-TO-USE PERMETHRIN STABLE EMULSION INSECTICIDE
- FOR CONTROL OF FLEAS, BROWN DOG TICKS AND LICE ON PREMISES
- FOR CONTROL OF PESTS ON LIVESTOCK
- WITH COUNTRY FRESH FRAGRANCE
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF FLEAS
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF STORED PESTS

### ACTIVE INGREDIENT

\*Permethrin..... 0.25%

**INERT INGREDIENTS**..... 99.75%  
100.00%

\*(3-phenoxyphenyl) methyl (+-) 3-(2,2-dichloroethenyl) 2,2-dimethyl cyclopropane carboxylate. Cis/trans ratio: min. 35% (+-) cis and max. 65% (+-) trans.

EPA REG. NO. 8848-67 EPA EST. NO. 8848-NY-1

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

See Side Panel For Additional Precautions

**NET CONTENTS: 16 FL. OZ.**

**A C C E P T E D**  
  
 JUN - 1 2004  
  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 8848-67

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry. Do not use on pets.

FIRST AID	
<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give any liquid by mouth to an unconscious person.</li> </ul>
<b>If Inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If On Skin Or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor or doctor for treatment advice.</li> </ul>
<b>If In Eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center for treatment advice.</li> </ul>
<p>Contains petroleum distillate—vomiting may cause aspiration pneumonia</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may contact 1-800-858-7378 for emergency medical treatment information.</p>	

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment wash waters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **If Empty:** Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

Remove cap, affix trigger, pump or other type sprayer to container for application. Do not transfer to another container. SHAKE WELL BEFORE USING

READ ENTIRE LABEL BEFORE EACH USE

### INDOOR APPLICATION

For spot application only. Do not use as a space indoors.  
SURFACE SPRAYING: FOR 60-DAY (TWO MONTHS) CONTROL OF: COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Direct the spray into hiding places, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

**ADDITIONAL ALTERNATE TEXT**  
**Screen Treatment:** Remove, clean and dry screens. Applying outdoors, hold bottle upright and spray 8-10 inches from surface with sweeping movement to apply evenly. Spray both sides of screen to assure complete coverage. Two may be treated together. Wipe any excess liquid with lint-free cloth. Use on any type of screening. Repeat as necessary.

---

**ADDITIONAL ALTERNATE TEXT**  
**SURFACE SPRAYING**  
**To Control Spiders:** Apply around infested areas, doorframes, pipes under sinks, cracks, crevices, carpets and rugs. It can be used indoors and outdoors on paint, metal, fiberglass, concrete and masonry. Reapply as needed.

**TO CONTROL CARPET BEETLES:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat as necessary.

**TO CONTROL BEDBUGS:** Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

**TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE):** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed. Allow to dry thoroughly before allowing pets to contact treated areas.

**TO CONTROL CLOTHES MOTHS:** Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

**TO KILL LICE AND LOUSE EGGS:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry-cleaned. Allow all treated particles to dry thoroughly before use.

**ON LIVESTOCK:** To protect cattle (beef and dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

**TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES:** Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

**TO CONTROL BLOOD-SUCKING LICE:** Apply to the infested areas of the animal, using a stiff brush to get the

spray to the base of the hair. Repeat every 3 weeks if required.

**TO CONTROL POULTRY LICE:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

**TO CONTROL BEDBUGS AND MITES IN POULTRY HOUSES:** Spray crevices of roost poles, cracks in walls, and cracks in nests where the Bedbugs and Mites hide.

**OUTDOOR APPLICATION**  
 To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

**FOR CONTROL OF GARDEN INSECTS:** For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray as well as Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leaf miners. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 degrees F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

**Safeguard Chemical Corporation**  
**Bronx, NY 10454**  
**Made in USA**