

## SUPER SHOCK

**DIRECTIONS FOR USE**  
Swimming Pool Disinfectant: Begin with fresh water. Use 6½ oz. ARCO SUPER SHOCK for each 10,000 gallons of water to get a 0.6 ppm. This dosage should be repeated until a chlorine residual of 0.6 to 1.0 ppm is obtained as measured by a test kit.

Mix required amount of ARCO SUPER SHOCK with about one part of water and feed solution through a chlorinator, or dilute and scatter over surface of pool. If warm weather prevails or pool has unusually heavy use, maintenance dosage may have to be repeated more frequently.

Adjust PH in the measurement of the acid or alkaline content of the pool water.



# SUPER SHOCK LIQUID CHLORINE FOR SWIMMING POOL AND WATER TREATMENT

### ACTIVE INGREDIENTS

Sodium hypochlorite . . . . . 12.5%

INERT INGREDIENTS . . . . . 87.5%

TOTAL . . . . 100.0%

Weight of gallon 9.8 lb. Equivalent chlorine per gallon 1.3 lb.

**CAUTION:** IRRITANT. HARMFUL IF SWALLOWED.  
KEEP OUT OF REACH OF CHILDREN.  
IRRITATING TO EYES AND SKIN. SEE ANTIDOTE STATEMENT  
AND OTHER REQUIRED STATEMENTS ON SIDE PANEL.

NET CONTENTS: 1 U. S. GALLON (128 FL. OZ.)

EPA Reg. No. 8866-1

MANUFACTURED BY

**ARCO CHEMICAL CORP.**  
MILWAUKEE, WISCONSIN 53225

## SUPER SHOCK

### ANTIDOTE

**EXTERNAL:** Wash immediately with water. Flood eyes and skin with cold water.

**INTERNAL:** Feed gruel or cooked cereal followed by olive oil or cooking oil.

**CALL A PHYSICIAN.**  
RINSE PRODUCT FROM SKIN. KEEP AWAY FROM CHILDREN.

**RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT.**

Degrades with age. Use test kit and increase dosage as necessary to obtain required level of available chlorine.