



ACCEPTED
4-26-67
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 8934-45

Ring Around Brand

10% CHLORDANE DUST

INGREDIENTS

ACTIVE INGREDIENTS:

Technical Chlordane*	10%
INERT INGREDIENTS	90%
Total	100%

*Equivalent to 6% octachloro-4,7-methanoindane and 4% related compounds.

CAUTION → 18 point and in ^{Red} ~~Red~~

12 point and in ^{Red} ~~Red~~ ← KEEP OUT OF REACH OF CHILDREN
See additional CAUTIONS statements on front panel.

CAUTION → 18 point and in red -

Avoid breathing dust or vapor. Avoid contamination of feed and foodstuffs.

TOXIC to Fish and Wildlife. Do not contaminate any body of water.

Birds feeding on treated areas may be killed.

Apply this product only as specified on label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be...

foodstuffs.

TOXIC to Fish and Wildlife. Do not contaminate any body of water.

Birds feeding on treated areas may be killed.

Apply this product only as specified on label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning (keep out of smoke).

DIRECTIONS FOR USE

ON LAWNS

ANTS, CHIGGERS, SOD WEBWORMS, CRICKETS, CUTWORMS, AND EARWIGS—Apply 12½ to 15 lb. per acre (3 oz. per 500 sq. ft. of area) directly into the grass. Repeat as necessary.

For imported fire ant, break open hard surface of mound and pour in 1 to 2 cupfuls per mound; sprinkle surrounding soil as for other ants. If ants survive, repeat in 2 weeks.

WHITE GRUBS AND JAPANESE BEETLE GRUBS—Apply 75 to 100 lb. per acre (1 lb. per 500 sq. ft. of area) in early spring or late fall. Repeat annually if necessary.

Keep children and pets off treated areas until the insecticide has been washed into the soil, either by rainfall or thorough sprinkling, and the grass has completely dried. DO NOT feed grass from treated areas to poultry, dairy animals, or animals being finished for slaughter. Do not pasture such animals on treated areas.

PREPLANTING SOIL APPLICATIONS

CUTWORMS—Apply 12½ to 15 lb. per acre (3 oz. per 500 sq. ft.) uniformly over surface of prepared land prior to setting of tomato, cabbage, pepper, or tobacco transplants. If desired, place 1 to 1½ teaspoonfuls in a ring around each transplant but not in direct contact with plant.

WIREWORMS—Apply 35 to 50 lb. per acre (½ to 2/3 lb. per 500 sq. ft.) uniformly over surface of prepared land, work into top 3 or 4 inches of soil prior to planting of corn, sweet corn, tobacco, or strawberries.

CABBAGE ROOT MAGGOT—Apply 35 to 50 lb. per acre (½ to 2/3 lb. per 500 sq. ft.). Work into top 3 inches of soil just before setting of cabbage, broccoli, or cauliflower transplants.

SEED CORN MAGGOT—Apply 12½ to 15 lb. per acre (3 oz. per 500 sq. ft.). Work into top 3 inches of seed bed ahead of seeding of peas, beans, lima beans, sweet corn.

WHITE GRUBS—Uniformly apply 75 to 100 lb. per acre (1 lb. per 500 sq. ft.) of prepared soil and work into top 4 to 6 inches of soil just before setting of strawberry or ornamental transplants.

USE FOR FLY REDUCTION

MAGGOTS OF HOUSEFLIES—Apply powder to surface of manure piles, dropping pits, garbage, and dumps in which flies breed. A minimum of 1 pound should be used to cover 250 square feet of surface.

NOTE: 1 ounce is approximately equivalent to 2 level tablespoonfuls.

NOTICE

Seller makes no warranty of any kind, expressed or implied, concerning the use of this product by the buyer or others. Seller shall not be liable for any claims arising out of the use of such materials, whether used singly or in combination with other substances. Buyer assumes all risk from use or handling of this product. Use only for the purpose and in the manner stated in this label.

NET WEIGHT
50 POUNDS

STEGALL - SYLVEST SEED CO., INC.
Montgomery, Ala.
USDA Reg.No.

PRODUCT NO. CH 10