

DIRECTIONS—Continued

and water. Usually 1 quart of VET'S MANGE & LICE diluted with 10 gallons (2 table-spoons in 2½ pints) of water will cover 5,000 square feet of surface or 500 square feet per gallon of diluted material. Either remove or cover all milking utensils before spraying. Cover all feed troughs. Remove animals from barn while spraying.

CAUTION

MAY BE ABSORBED THROUGH SKIN. AVOID INHALATION AND SKIN CONTACT. IN CASE OF CONTACT, WASH IMMEDIATELY WITH SOAP AND WATER. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. DO NOT USE ON HOUSEHOLD PETS OR HUMANS. HARMFUL IF SWALLOWED. KEEP OUT OF REACH OF CHILDREN. DO NOT USE ON FEED OR FORAGE TO BE FED TO DAIRY ANIMALS, OR ANIMALS BEING FINISHED FOR SLAUGHTER. THIS PRODUCT WILL KILL FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. DO NOT USE IN POULTRY HOUSES. DO NOT DIP ANIMALS.

NOTICE: No warranty expressed or implied is made concerning this product or its use except that it conforms to the chemical description on this label and no person is authorized to make any other representation.

U.S.D.A. Reg. No. 8881-4

VETS

LIVESTOCK

Mange & Lice CONTROL

**CONTAINS
12.5% LINDANE**

**CAUTION
KEEP OUT OF
REACH OF
CHILDREN**

ACTIVE INGREDIENTS:

Lindane 12.5%
Inert Ingredients 87.5%

INERT INGREDIENTS:

NET CONTENTS: ONE QUART

SEE VET'S MANGE & LICE
FOR ADDITIONAL DIRECTIONS

DIRECTIONS Before Use Read Caution and Notice Statements Carefully

ANIMAL TREATMENT

Do not apply to beef cattle and hogs within 30 days of slaughter. Do not treat animals under 3 months of age. Do not use on dairy animals. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

To control psoroptic mites (responsible for sheep scabies), beef cattle & hog lice, ticks on beef cattle & sheep ticks, thoroughly cover animals with a spray consisting of 1 qt. VET'S MANGE & LICE mixed with 100 gallons (1 tablespoon in 1½ gallons) of water. Repeat applications as required.

Sarcoptic mange on hogs may be effectively reduced or controlled by the use of 1 quart of VET'S MANGE & LICE in 60 gallons (1 tablespoon in 1 gallon) of water. A second application should follow the first by one week. Two quarts of diluted spray per animal is required for adult horses and cattle and one quart for calves. Avoid spraying feed, forage or feed and drinking troughs.

Residual Spray

Flies, mosquitoes, gnats and roaches can be controlled by spraying walls, ceilings, window ledges and other resting places of insects in barns and other buildings with a solution of VET'S MANGE & LICE

8-22-67

8881-4