

FEED TAG FACSIMILE
JIM DANDY
BAR-FLY
MEDICATED BLOCK

(For Ruminants Only)
 33 1/3 LBS. NET WT.

Prevents the breeding of horn flies and face flies in the manure of treated cattle; aids in the prevention of stomach worms, hookworms and nodular worms.

ACTIVE INGREDIENT:

Phenothiazine (Biologically Assayed) 1.25%
 INERT INGREDIENTS:* 98.75%
 Total 100.00%

*Refers only to ingredients which are neither insecticides or drugs.

WARNING: This feed should be used only in accordance with directions on this label.

INGREDIENTS:

Cottonseed meal, meat and bone meal, dehydrated alfalfa meal, dried corn fermentation solubles, grain sorghum meal, yeast culture, cane molasses, urea, salt, monocalcium phosphate, disodium phosphate, low fluorine rock phosphate, magnesium carbonate, cobalt sulfate, cobalt carbonate, ethylene diamine dihydriodide, copper carbonate, zinc sulfate, magnesium sulfate, magnesium oxide, manganese sulfate, iron oxide, sodium bicarbonate, bentonite, guar meal, sulfur, Vitamin B₁₂ supplement, niacin, choline chloride, calcium pantothenate, riboflavin supplement, anise oil, Vitamin A palmitate and irradiated dried yeast (source of Vitamin D₂).

Guaranteed by

WESTERN GRAIN COMPANY
 BIRMINGHAM, ALABAMA 35201

GUARANTEED ANALYSIS:

Crude Protein, not less than	18.00%
(This includes not more than 6.00% equivalent protein from non-protein nitrogen)	
Crude Fat, not less than	1.00%
Crude Fiber, not more than	8.00%
Nitrogen-free extract, not less than	23.00%
Calcium (Ca), not more than	7.00%
Calcium (Ca), not less than	6.00%
Phosphorus (P), not less than	1.50%
Iodine (I), not less than	0.02%
Salt (NaCl), not more than	14.50%
Salt (NaCl), not less than	12.50%
Magnesium (Mg), not less than	1.25%
Manganese (Mn), not less than	0.012%
Cobalt (Co), not less than	0.002%
Vitamin D ₂ , not less than	

30,000 USP units per lb.

FEED TAG FACSIMILE
KATTLE-KING

BAR-FLY
MEDICATED BLOCK

(For Ruminants Only)
 33 1/3 LBS. NET WT.

Prevents the breeding of horn flies and face flies in the manure of treated cattle; aids in the prevention of stomach worms, hookworms and nodular worms.

ACTIVE INGREDIENT:

Phenothiazine (Biologically Assayed) 1.25%
 INERT INGREDIENTS:* 98.75%
 Total 100.00%

*Refers only to ingredients which are neither insecticides or drugs.

WARNING: This feed should be used only in accordance with directions on this label.

INGREDIENTS:

Cottonseed meal, meat and bone meal, dehydrated alfalfa meal, dried corn fermentation solubles, grain sorghum meal, yeast culture, cane molasses, urea, salt, monocalcium phosphate, disodium phosphate, low fluorine rock phosphate, magnesium carbonate, cobalt sulfate, cobalt carbonate, ethylene diamine dihydriodide, copper carbonate, zinc sulfate, magnesium sulfate, magnesium oxide, manganese sulfate, iron oxide, sodium bicarbonate, bentonite, guar meal, sulfur, Vitamin B₁₂ supplement, niacin, choline chloride, calcium pantothenate, riboflavin supplement, anise oil, Vitamin A palmitate and irradiated dried yeast (source of Vitamin D₂).

Guaranteed by

V. M. E. CORP.
 FORT WORTH, TEXAS 76106

GUARANTEED ANALYSIS: -

Crude Protein, not less than	18.00%
(This includes not more than 6.00% equivalent protein from non-protein nitrogen)	
Crude Fat, not less than	1.00%
Crude Fiber, not more than	8.00%
Nitrogen-free extract, not less than	23.00%
Calcium (Ca), not more than	7.00%
Calcium (Ca), not less than	6.00%
Phosphorus (P), not less than	1.50%
Iodine (I), not less than	0.02%
Salt (NaCl), not more than	14.50%
Salt (NaCl), not less than	12.50%
Magnesium (Mg), not less than	1.25%
Manganese (Mn), not less than	0.012%
Cobalt (Co), not less than	0.002%
Vitamin D ₂ , not less than	

ACC'

3-8-68

UNDER THE
 FUNGICIDE AN
 FOR ECONOMIC
 RECORDER NO

88947

Insecticides or drugs.

WARNING: This feed should be used only in accordance with directions on this label.

INGREDIENTS:

Cottonseed meal, meat and bone meal, dehydrated alfalfa meal, dried corn fermentation solubles, grain sorghum meal, yeast culture, cane molasses, urea, salt, monocalcium phosphate, disodium phosphate, low fluorine rock phosphate, magnesium carbonate, cobalt sulfate, cobalt carbonate, ethylene diamine dihydriodide, copper carbonate, zinc sulfate, magnesium sulfate, magnesium oxide, manganese sulfate, iron oxide, sodium bicarbonate, bentonite, guar meal, sulfur, Vitamin B₁₂ supplement, niacin, choline chloride, calcium pantothenate, riboflavin supplement, anise oil, Vitamin A palmitate and irradiated dried yeast (source of Vitamin D₂).

Guaranteed by

WESTERN GRAIN COMPANY

BIRMINGHAM, ALABAMA 35201

GUARANTEED ANALYSIS:

Crude Protein, not less than	18.00%
(This includes not more than 6.00% equivalent protein from non-protein nitrogen)	
Crude Fat, not less than	1.00%
Crude Fiber, not more than	8.00%
Nitrogen-free extract, not less than	23.00%
Calcium (Ca), not more than	7.00%
Calcium (Ca), not less than	6.00%
Phosphorus (P), not less than	1.50%
Iodine (I), not less than	0.02%
Salt (NaCl), not more than	14.50%
Salt (NaCl), not less than	12.50%
Magnesium (Mg), not less than	1.25%
Manganese (Mn), not less than	0.012%
Cobalt (Co), not less than	0.002%
Vitamin D ₂ , not less than	

30,000 USP units per lb.

FEEDING DIRECTIONS

Feed free choice. Intake will be 5 to 6 ounces per head daily depending on bodyweight of animal. Do not offer any other mineral or salt mixtures. The above daily intake will furnish the proper levels of medication.

WARNING: Do not feed this medicated block to lactating dairy animals.

Occasional sensitivity to phenothiazine has been reported in cattle.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Note: For maximum protection against horn flies and face flies, it is recommended that cattle be sprayed once when first starting on a continuous JIM DANDY BAR-FLY Program. JIM DANDY BAR-FLY prevents the breeding of horn flies and face flies in the manure but does not affect any existing adult fly program. Be sure and supply ample blocks at all times -- at least one block for every 10 to 12 head.

11 67

USDA Reg. #8891

WARNING: This feed should be used only in accordance with directions on this label.

INGREDIENTS:

Cottonseed meal, meat and bone meal, dehydrated alfalfa meal, dried corn fermentation solubles, grain sorghum meal, yeast culture, cane molasses, urea, salt, monocalcium phosphate, disodium phosphate, low fluorine rock phosphate, magnesium carbonate, cobalt sulfate, cobalt carbonate, ethylene diamine dihydriodide, copper carbonate, zinc sulfate, magnesium sulfate, magnesium oxide, manganese sulfate, iron oxide, sodium bicarbonate, bentonite, guar meal, sulfur, Vitamin B₁₂ supplement, niacin, choline chloride, calcium pantothenate, riboflavin supplement, anise oil, Vitamin A palmitate and irradiated dried yeast (source of Vitamin D₂).

Guaranteed by

V. M. E. CORP.

FORT WORTH, TEXAS 76106

GUARANTEED ANALYSIS:

Crude Protein, not less than	18.00%
(This includes not more than 6.00% equivalent protein from non-protein nitrogen)	
Crude Fat, not less than	1.00%
Crude Fiber, not more than	8.00%
Nitrogen-free extract, not less than	23.00%
Calcium (Ca), not more than	7.00%
Calcium (Ca), not less than	6.00%
Phosphorus (P), not less than	1.50%
Iodine (I), not less than	0.02%
Salt (NaCl), not more than	14.50%
Salt (NaCl), not less than	12.50%
Magnesium (Mg), not less than	1.25%
Manganese (Mn), not less than	0.012%
Cobalt (Co), not less than	0.002%
Vitamin D ₂ , not less than	

30,000 USP units per lb.

FEEDING DIRECTIONS

Feed free choice. Intake will be 5 to 6 ounces per head daily depending on bodyweight of animal. Do not offer any other mineral or salt mixtures. The above daily intake will furnish the proper levels of medication.

WARNING: Do not feed this medicated block to lactating dairy animals.

Occasional sensitivity to phenothiazine has been reported in cattle.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Note: For maximum protection against horn flies and face flies, it is recommended that cattle be sprayed once when first starting on a continuous KATTLE-KING BAR-FLY Program. KATTLE-KING BAR-FLY prevents the breeding of horn flies and face flies in the manure but does not affect any existing adult fly problem. Be sure and supply ample blocks at all times -- at least one block for every 10 to 12 head.

11 67

USDA Reg. #8891