AGRICULTURAL FUNGICIDE

USDA REG. NO. 8901-2

ACTIVE INGREDIENT

CUPRIC HYDROXIDE _____ 21.0 %

79.0 % INERT INGREDIENTS_____

> TOTAL 100%

(METALLIC COPPER EQUIVALENT - 13.7%)

DO NOT MIX WITH WATER

CAUTION

HARMFUL IF SWALLOWED

KEEP OUT OF REACH OF CHILDREN SEE BACK PANEL FOR ADDITIONAL CAUTIONS

50 GALLONS

Kennecott COPPER CORPORATION



SPECIAL PRODUCTS DIVISION

NEW YORK, NEW YORK 10017

KOCIDE 220 AGRICULTURAL FUNGICIDE

GENERAL INSTRUCTIONS

Use KOCIDE 220 as noted below. KOCIDE 220 is for aircraft or mist sprayers capable of spraying low volumes (1-3 gal/acre). KOCIDE 220 is not intended to be mixed with water, but used as supplied.

In case of skin contact, remove with washing; for eyes, wash with copious amounts of water and get immediate medical attention.

The following specific instructions are based on general applications. However, the recommendation of local Agricultural Experiment Stations should be closely followed as to timing, frequency and number of sprays per season.

SPECIFIC INSTRUCTIONS

CROP


TO CONTROL

RECOMMEN DATIONS

Sugar Beets (Texas-New Mexico)

Cercospora leaf-

Start spray when disease threatens and continue for four to five applications. Spray 10 to 14 days depending on weather conditions and disease severity at one gallon



Avoid contact with skin and eyes. In case of skin contact, remove with washing; for eyes, wash with copious amounts of water and get immediate medical attention.

Keep out of any body of water; fish may be killed.

Kennecott corpus corporation special products division NIW YORK NIW YORK 1001