


LEBANON
LAWN FOOD
10-6-4
WITH
INSECT
CONTROL



LAWN FOOD

WITH INSECT CONTROL

CONTAINS DUREX

Controls Chinch Bugs, Sod Webworms, Brown Mealybugs, Army Worms, Flea Beetles, Cutworms, Sod Webworms, Japanese Beetles, and Hyperaspis Beetles. Controls Japanese Beetles and European Chinch Bugs.

CAUTION: KEEP OUT OF REACH OF CHILDREN See additional caution on back panel.

ACTIVE INGREDIENT:	
Chlorpyrifos [O,O-Diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate]	0.60%
INERT INGREDIENTS:	
	99.40%
TOTAL	100.00%

"Durex" - Trademark of the Dow Chemical Company.
U. S. Patent No. 3,244,586
E.P.A. Reg. No. 961-275

LAWN FOOD

New Weight 20 lbs.
Treats 1,000 sq. ft.
LEBANON LAWN FOOD
WITH INSECT CONTROL

Guaranteed Minimum Fertilizer Analysis

TOTAL NITROGEN IN	6.0%
AVAILABLE PHOSPHORIC ACID P ₂ O ₅	6.0%
SOLUBLE POTASH K ₂ O	4.0%
Potential Acidity 1,000 lb. Calcium Carbonate per ton	

LEBANON CHEMICAL CORPORATION
LEBANON, PA. 17042

Feeds your lawn the essential plant food elements required for good growth, while controlling the injurious lawn and lawn pests: Chinch Bugs, Sod Webworms, Japanese Beetles, Army Worms, Flea Beetles, Cutworms, Sod Webworms, Japanese Beetles, and Hyperaspis Beetles. Also for control of larvae of Japanese Beetles and European Chinch Bugs.

INSECT CONTROL

Army Worm:	1 lb. per 1,000 sq. ft.	Chinch Bug:	1 lb. per 1,000 sq. ft.
Japanese Beetle:	1 lb. per 1,000 sq. ft.	Hyperaspis Beetle:	1 lb. per 1,000 sq. ft.
Cutworm:	1 lb. per 1,000 sq. ft.	Sod Webworm:	1 lb. per 1,000 sq. ft.

NOTE: Above rates are approximate. Check your records on 100 sq. ft. and adjust to apply 1 lb. per 1,000 sq. ft. Rate of water, moisture, and a water conditioner can vary the results suggested.

DIRECTIONS

For control of all insects named above, except Hyperaspis Beetles and larvae of Japanese Beetles and European Chinch Bugs, apply at the rate of 1 lb. per 1,000 sq. ft. to lawns or lawn areas. Treat when insects first appear or before they appear.

For control of Japanese Beetles and European Chinch Bugs, apply at the rate of 1 lb. per 1,000 sq. ft. to lawns or lawn areas. Treat when insects first appear or before they appear. Apply in mid-April and again in mid-May. Consult your local Agricultural Extension Service for best time to apply in your area.

For control of larvae of Japanese Beetles and European Chinch Bugs, apply at the rate of 1 lb. per 1,000 sq. ft. to lawns or lawn areas. Treat when insects first appear or before they appear. Apply in mid-April and again in mid-May. Consult your local Agricultural Extension Service for best time to apply in your area.

Immediately after application sprinkle the treated area with 30 gallons of water per 1,000 sq. ft. to wash the granules into the infested area.

CAUTION

Keep out of reach of children. May be harmful if swallowed. Do not take internally. Avoid eye contact. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before re-use.

Keep away from food, feedstuffs and domestic water supplies. NOT TO BE USED: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift over areas treated.

Do not re-use empty container. Dispose of empty container and wastes by burning or by burying in non-crop lands away from domestic water supplies.

WARRANTY

Lebanon Chemical Corporation warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither the warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

KOCLIDE

3% DUST

AND SULFUR

AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS

CUPRIC HYDROXIDE* 2.5%

SULFUR 87.5%

INERT INGREDIENTS 10.0%

TOTAL 100%

(METALLIC COPPER EQUIVALENT - 1.6%)

*MANUFACTURED UNDER U.S. PATENT NO. RE. 24,328

CAUTION: KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

50 LBS. PER

USDA REG. NO. 80014

**KOCIDE 3% DUST AND SULFUR LABEL
BACK PANEL**

**KOCIDE 3% DUST
AND SULFUR**

AGRICULTURAL FUNGICIDE

GENERAL INSTRUCTIONS

KOCIDE 3% Dust (and Sulfur) should be used as described below.

The specific instructions are based upon general applications. Recommendations of State Agricultural Experiment Station or State Extension Service Specialist should be followed as to frequency and number of applications per season.

SPECIFIC INSTRUCTIONS

KOCIDE 3% Dust (and Sulfur) is especially formulated for control of *Cercospora* Leaf Spot on peanuts. Begin dusting 40-45 days after planting or when disease symptoms appear. Use at 15-25 pounds per acre. Continue applications at 10-14 day intervals.

Do Not Reuse Container.

Destroy When Empty

CAUTION --- Avoid contact with skin and eyes. In case of skin contact, remove with washing; for eyes, wash with copious amounts of water and get immediate medical attention.

Keep out of any body of water; fish may be killed.

Do not contaminate water by cleaning of equipment or by disposal of waste.

Kennecott COPPER CORPORATION SPECIAL PRODUCTS DIVISION

ACCEPTED
7/29/71
IN FULLY
FIVE LBS
ED IN 44 A
TO ATTACHED COMMENTS.
KOCIDE
DUST
SUBJECT

7/21/71