

BACK LABEL

K-COP AGRICULTURAL FUNGICIDE

GENERAL INSTRUCTIONS

Use K-COP as directed below. Depending on the equipment used and the specific crop, the volume applied per acre will differ. For high volume sprays, use from 25 to 100 gallons per acre (GPA). For concentrate ground sprays, apply from 20 to 50 GPA. For aerial spraying, 3 to 15 GPA are commonly used.

Observe all cautions and limitations on label of all products used as mixtures. Since the number of conditions of pesticide mixtures is very large, it is advisable to test mixtures first and to apply mixtures as soon as possible.

The following specific instructions are based on general applications. The recommendations of local Agricultural Experiment Stations should be closely followed as to timing, frequency and number of sprays per season.

SPECIFIC INSTRUCTIONS

BEANS, Bacterial Blight (Halo & Common). Use 0.33 to 0.75 gallon per acre. Make first application when plants are 3 to 5 inches high before disease symptoms appear. Continue application on a 7-10 day schedule, depending on local conditions.

CANTALOUPE, HONEYDEWS, MUSKMELON, WATERMELON, Downy Mildew, Powdery Mildew, Alternaria & Angular Leaf Spot, Scab. Apply at 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis. In areas where Angular Leaf Spot is a problem, applications should be made on a 5-7 day schedule.

CELERY, Early, Late & Bacterial Blight. Apply 0.33 to 0.75 gallon per acre. Begin applications when plants are first established, then every 7 days, depending on disease conditions.

CUCUMBERS, Alternaria & Angular Leaf Spot, Powdery Mildew. Apply at 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis. In areas where Angular Leaf Spot is a problem, applications should be made on a 5-7 day schedule.

PEANUTS, Cercospora Leaf Spot. Begin spraying at 0.33 to 0.75 gallon per acre at first sign of disease and every 10-14 days thereafter.

PEPPERS, Bacterial Spot, Cercospora Leaf Spot. Apply at 0.33 to 0.75 gallon per acre. Begin application before disease appears and continue at 7-10 day intervals.

POTATOES, Early & Late Blight. Apply at 0.33 to 0.75 gallon per acre. Begin applications when plants are 3-4 inches high and continue at 7-10 day intervals.

SOYBEANS, Bacterial Blight (Halo & Common). Use 0.47 to 0.6 gallon per acre. Make first application when plants are 3-5 inches high before disease symptoms appear. Continue applications on a 7-10 day schedule, depending on local conditions.

SQUASH, (summer) & (winter), Alternaria & Angular Leaf Spot, Powdery Mildew. Use 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis. In areas where Angular Leaf Spot is a problem, applications should be made on a 5-7 day schedule.

SUGAR BEETS, Cercospora Leaf Spot. Use 0.33 to 0.75 gallon per acre. Begin when plants are 3-4 inches high and continue on a 10-14 day interval or more frequently if disease conditions are severe.

TOMATO, Bacterial Spot, Early Blight. Use 0.33 to 0.75 gallon per acre. In seed beds, begin applications when seedlings emerge and repeat at 4-5 day intervals. In field, begin applications immediately after transplanting and continue on 7 day intervals or more frequently if disease conditions are severe.

CAUTION--Avoid contact with skin and eyes. In case of skin contact, remove with washing; for eyes, wash with copious amounts of water and get immediate medical attention.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or by disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

NOTICE: KOCIDE CHEMICAL CORPORATION warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the plants and crops specified. THERE ARE NO OTHER WARRANTIES, WHETHER EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

FRONT LABEL

KOCIDE SD

SEED DRESSING

AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT

CUPRIC HYDROXIDE _____ 30%

INERT INGREDIENTS _____ 70%

ACCEPTED

TOTAL 100%

(METALLIC COPPER EQUIVALENT — 19.5%)

MAR 20 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 8901-17


8901-17

**CAUTION: KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED**

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

5 GALLONS

EPA Reg. No. 8901-17



KOCIDE
Chemical

