

K-COP[®]

LIQUID AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT
*Copper as elemental 8
Inert Ingredients 92
100%

*From copper-ammonium complex

CAUTION
KEEP OUT OF REACH
OF CHILDREN
HARMFUL IF SWALLOWED

NET CONTENT:
2½ Gallons



KOCIDE Chemical Corporation
Houston, Texas 77045



BACK LABEL

ACCEPTED

SEP 11 1981

Federal Insecticide, Fungicide, and Rodenticide Act

Registration for the pesticide

K-COP AGRICULTURAL FUNGICIDE

GENERAL INSTRUCTIONS

K-COP should be used as described below. K-COP is adaptable to all types of spray equipment. For dilute, high volume sprays, use from 25 to 100 gallons per acre (GPA). For concentrate ground sprays, apply 20 to 50 GPA. For aerial spraying, 3 to 15 GPA are commonly used. Follow label recommended rates per acre.

Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

The following directions for use are based upon general applications. The recommendations of State Agricultural Experiment Stations should be closely followed as to timing, frequency and number of applications per season.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or Local procedures under the Resource Conservation and Reclamation Act. Dispose of container according to Federal, State or Local procedures under the Resource Conservation and Reclamation Act.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEANS, *Bacteria Blight* (Halo & Common): Use 0.33 to 0.75 gallon per acre. Make first application when plants are 3 to 5 inches high before disease symptoms appear. Continue application on a 7-10 day schedule depending on local conditions.

CANTALOUPE, HONEYDEW'S, MUSKMELON, WATERMELON, *Downy Mildew, Powdery Mildew, Alternaria & Angular Leaf Spot, Scab*: Apply at 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis. In areas where *Angular Leaf Spot* is a problem, applications should be made on a 7 day schedule.

CELERY, *Early, Late & Bacteria Blight*: Apply 0.33 to 0.75 gallon per acre. Begin applications when plants are first established, then every 7 days depending on disease conditions.

CUCUMBERS, *Alternaria & Angular Leaf Spot, Powdery Mildew*: Apply 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis. In areas where *Angular Leaf Spot* is a problem, applications should be made on a 5-7 day schedule.

PEANUTS, *Cercospora Leaf Spot*: Begin spraying at 0.33 to 0.75 gallon per acre at first sign of disease and every 7-14 days thereafter.

PEPPERS, *Bacteria Spot, Cercospora Leaf Spot*: Apply at 0.33 to 0.75 gallon per acre. Begin application before disease appears and continue at 7-10 day intervals.

POTATOES, *Early & Late Blight*: Apply at 0.33 to 0.75 gallon per acre. Begin applications when plants are 3-4 inches high and continue at 7-10 day intervals.

SOYBEANS, *Bacteria Blight, Halo & Common*: Use 0.47 to 0.6 gallon per acre. Make first application when plants are 3-5 inches high before disease symptoms appear. Continue applications on a 7-10 day schedule depending on local conditions.

SQUASH, *Summer & Winter, Alternaria & Angular Leaf Spot, Powdery Mildew*: Use 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis. In areas where *Angular Leaf Spot* is a problem, applications should be made on a 5-7 day schedule.

SUGAR BEETS, *Cercospora Leaf Spot*: Use 0.33 to 0.75 gallon per acre. Begin when plants are 3-4 inches high and continue on a 10-14 day interval or more frequently if disease conditions are severe.

TOMATO, *Bacteria Spot, Early Blight*: Use 0.33 to 0.75 gallon per acre. In seed beds, begin applications when seedlings emerge and repeat at 4-5 day intervals. In field, begin applications immediately after transplanting and continue on 7 day intervals or more frequently if disease conditions are severe.

WALNUTS, *Walnut Blight*: Use 0.33 to 1.0 gallon per acre. Make first application at early pre-bloom. Make second application at late pre-bloom. If disease conditions persist, make additional applications.

CAUTION: Avoid contact with skin and eyes. In case of skin contact, remove with washing; for eyes, wash with copious amounts of water and get immediate medical attention. This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

NOTICE: KOCIDE CHEMICAL CORPORATION warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the plants and crops specified. **THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE.** This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

KOCIDE Chemical Corporation