

BACK LABEL

ACCEPTED

AUG 11 1988

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 596-11

# K-COP AGRICULTURAL FUNGICIDE

## PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Apply this product only as specified on this label.

### GENERAL INSTRUCTIONS

K-COP should be used as described below. K-COP is adaptable to all types of spray equipment. For dilute, high volume sprays, use from 25 to 100 gallons per acre (GPA). For concentrate ground sprays, apply 20 to 50 GPA. For aerial spraying, 3 to 15 GPA are commonly used. Follow label recommended rates per acre.

Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

The following directions for use are based upon general applications. The recommendations of State Agricultural Experiment Stations should be closely followed as to timing, frequency and number of applications per season.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by State or Local authorities.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**BEANS, Bacteria, Blight (HA) & Common:** Use 0.33 to 0.75 gallon per acre. Make first application when plants are 3 to 5 inches high before disease symptoms appear. Continue application on a 7 to 10 day schedule depending on local conditions.  
**CANTALOUPE, HONEYDEWS, MUSKMELON, WATERMELON, Powdery Mildew, Angular Leaf Spot, Scab:** Apply at 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis in areas where Angular Leaf Spot is a problem. Applications should be made on a 5 to 7 day schedule.  
**CELERY, Early & Late Blight:** Apply 0.33 to 0.75 gallon per acre. Begin applications when plants are first established then every 7 days depending on disease conditions.  
**CUCUMBERS, Alternaria & Angular Leaf Spot, Powdery Mildew:** Apply 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis in areas where Angular Leaf Spot is a problem. Applications should be made on a 5 to 7 day schedule.  
**PEANUTS, Cercospora Leaf Spot:** Begin spraying at 0.33 to 0.75 gallon per acre at first sign of disease and every 10 to 14 days thereafter.  
**PEPPERS, Bacterial Spot, Cercospora Leaf Spot:** Apply at 0.33 to 0.75 gallon per acre. Begin application when disease appears and continue at 7 to 10 day intervals.  
**POTATOES, Early & Late Blight:** Apply at 0.33 to 0.75 gallon per acre. Begin applications when plants are 3 to 4 inches high and continue at 7 to 10 day intervals.

**SOYBEANS, Bacteria, Blight (HA) & Common:** Use 0.47 to 0.6 gallon per acre. Make first application when plants are 3 to 5 inches high before disease symptoms appear. Continue application on a 7 to 10 day schedule depending on local conditions.  
**SQUASH, Summer & Winter, Alternaria & Angular Leaf Spot, Powdery Mildew:** Use 0.33 to 0.75 gallon per acre when plants begin to vine. Continue applications on a weekly basis in areas where Angular Leaf Spot is a problem. Applications should be made on a 5 to 7 day schedule.  
**SUGAR BEETS, Cercospora Leaf Spot:** Use 0.33 to 0.75 gallon per acre. Begin when plants are 3 to 4 inches high and continue on a 10 to 14 day interval or more frequently if disease conditions are severe.  
**TOMATOES, Bacterial Spot, Early Blight:** Use 0.33 to 0.75 gallon per acre. In seed beds, begin applications when seedlings emerge and repeat at 4 to 5 day intervals. In field, begin applications immediately after transplanting and continue on 7 day intervals or more frequently if severe conditions are present.  
**WALNUTS, Walnut Blight:** Use 2.5 to 3.0 gallons per acre. Make first application at early pre-bloom. Make second application at late pre-bloom. If disease conditions persist, make additional applications.

**NOTICE:** KOCIDE CHEMICAL CORPORATION warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the plants and crops specified. **THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE.** This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes all risk of any such use.

KOCIDE Chemical Corporation Houston Texas 77045

# K-COP

## LIQUID AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT 8%  
\*Copper as elemental 8%  
INERT INGREDIENTS 92%  
100%

\*From copper-ammonium complex

KEEP OUT OF REACH  
OF CHILDREN  
**CAUTION**  
HARMFUL IF SWALLOWED

EPA Reg No 8901-14

See Back Panel For Additional  
Precautionary Statements.

NET CONTENT:  
2½ Gallons



**KOCIDE Chemical Corporation**  
Houston, Texas 77045