

FRONT LABEL

CORPORATION

Chemical

# Kocide 20/20

ACTIVE INGREDIENT	30.7%
COPPER HYDROXIDE (METALLIC COPPER EQUIVALENT 20%)	
INERT INGREDIENTS	69.3%
(METALLIC ZINC EQUIVALENT 20% DERIVED FROM BASIC ZINC SALTS)	
TOTAL	100%

**CAUTION:** KEEP OUT OF REACH  
OF CHILDREN.  
HARMFUL IF SWALLOWED.  
IF SWALLOWED,  
INDUCE VOMITING.  
GET MEDICAL ATTENTION.

U.S. PATENT NO. 3, 428, 731  
EPA Reg. No. 8901-22

**WETTABLE POWDER  
AGRICULTURAL  
FUNGICIDE  
AND  
ZINC NUTRITIONAL**

**20 LBS. NET**

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

BACK LABEL

**KOCIDE 20/20**  
**GENERAL INSTRUCTIONS**

KOCIDE 20/20 should be used as described below. KOCIDE 20/20 is adaptable to spraying from all types of spray equipment. Depending on the equipment used, the volume applied per acre will differ.

No additional surfactants are needed. Add KOCIDE 20/20 slowly to a spray tank partially filled with water. Spreader-stickers, insecticides, etc. should be added last. KOCIDE 20/20 is compatible with many commercially formulated spreader-stickers, oils and such insecticides as Carbaryl, Parathion, Toxaphene and other fungicides. Observe all cautions and limitations on label of all products used in mixtures. Since the number of combinations of pesticide mixtures is very large, it is advisable to test tank mixtures first and then apply as soon as possible.

The following directions for use are based on general applications. The recommendations of State Agricultural Extension Services should be closely followed as to timing, frequency and number of sprays per season.

**STORAGE AND DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or Local disposal authorities. Before disposal of container, triple rinse (or equivalent) and dispose of in an approved landfill or bury in a safe place. Consult Federal, State or Local disposal authorities for approved alternative procedures.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**CITRUS, Brown Rot & Septoria.** Apply 10 pounds per acre in 500-600 gallons by dilute ground spray or 50-100 gallons per acre as concentrate ground spray. For aerial application, use 10 pounds in 10-20 gallons per acre. Begin applications in fall after first rains and before fruit becomes infected.

**CAUTION:** Avoid contact with skin and eyes. In case of skin contact, remove with washing; for eyes, wash with copious amounts of water and get immediate medical attention. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or by disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

**NOTICE:** KOCIDE CHEMICAL CORPORATION warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions on the plants and crops specified. THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING A WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

BEST DOCUMENT AVAILABLE