




Ring Around Brand

7 1/2 % SEVIN DUST

ACTIVE INGREDIENTS

INGREDIENTS

75% gamma-Hexachlorocyclopentadiene



CAUTION
KEEP OUT OF REACH OF CHILDREN

See additional CAUTIONS Statements on front panel

Keep out of reach of children. Swallowed or absorbed through skin. Do not breathe dust. Avoid contact with eyes. Avoid contact with any insect. Do not use in or near areas where children play. Wash hands thoroughly after use. Do not use in or near areas where children play. Avoid contact with eyes. Avoid contact with skin. Avoid contact with clothing and watering plants.

USE ONLY FOR USE

TOBACCO

CORN

CUCUMBER, MELONS AND SQUASH

CABBAGE, BROCCOLI AND CAULIFLOWER

POTATOES, EGGPLANT AND PEPPERS

TOMATOES

STRAWBERRIES

NOTICE

Seller makes no warranty of any kind, expressed or implied, concerning the use of this product by the Buyer or others. Seller shall not be liable for any claims arising out of the use of such materials, whether used singly or in combination with other substances. Buyer assumes all risk from use or handling of this product. Use only for the purpose and in the manner stated in this label.

NET WEIGHT
30 POUNDS

RING AROUND PROTECTION

PRODUCT NO. 10

8934-7

Change to clean clothing daily. Avoid contamination of feed and foodstuffs. Avoid contamination of feeding and watering receptacles.

DIRECTIONS FOR USE

Apply when the air is calm with a good hand duster or suitable power operated dusting machine. Make thorough coverage of plants with particular attention to under-surfaces of foliage. Apply when the insects first appear, and repeat at regular intervals or as needed to maintain control of the insects on the following crops.

TOBACCO - To control hornworms, budworms, flea beetles, Japanese beetle adults, and grasshoppers, uniformly apply 20 to 30 pounds per acre, depending upon size of plants. Use the higher rate only on large tobacco plants. Heavier application may cause plant injury. Uniformly full coverage of each plant is essential.

To control budworms and flea beetles when using rotary hand duster to treat individual tobacco plants, apply 6 to 8 pounds per acre, with particular attention to the buds if budworms are prevalent. Efficient hand dusting equipment is required. Shaker cans and cloth bags cause excessive deposits of dust.

If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered as soon as the dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

COTTON For early-season control for thrips, cotton fleahopper, and cotton leafworm, apply 7 to 14 pounds per acre. Begin applications when insects are first observed and repeat at 7 to 10-day intervals as necessary.

For mid and late-season control of boll weevil, bollworm, fall armyworm, cotton leaf perforator, leafhoppers, and tarnished plant bug, apply 14 to 28 pounds per acre, depending upon the size of cotton plants and relative abundance of insects, and repeat at 5 to 7-day intervals as necessary. Repeated applications may be required at 4 to 5-day intervals for boll weevil, and at 3 to 5-day intervals for bollworm. Aphids on cotton are usually suppressed by these repeated applications in a regular treatment schedule.

For pink bollworm, apply 20 to 30 pounds per acre, depending upon the size of cotton plants and severity of infestation, beginning at the time the eggs start to hatch and repeated at 5 to 7-day intervals as long as control is necessary.

For stink bugs and salt-marsh caterpillar, apply 28 pounds per acre. For grasshoppers, apply 14 to 20 pounds per acre. Begin applications on appearance of the insects or their feeding damage and repeat as needed. The highest dosage should be used when the insects are approaching maturity or application is made to partially defoliated plants.

Applications to cotton should be made uniformly and thoroughly at the rates described throughout the entire field. Applications may be made after bolls open, but allow 7 days from last application before grazing treated fields or harvesting crop remnants for use as food or bedding for dairy or meat animals.

CORN - To control corn earworm, European corn borer, fall armyworm, flea beetles, Japanese beetle (adults), apply 14 to 28 pounds per acre. For larvae in the whorls (budworms) and foliage feeders, apply to entire plants. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3-day intervals beginning when first silks appear and continuing until they begin to dry. Three or more applications may be required depending upon severity of infestation. Timing and good coverage are essential.

CUCUMBER, MELONS, AND SQUASH - To control pickleworm, melonworm, cucumber beetles, leafhoppers, and squash bug, apply 14 pounds per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of high humidity for an extended period. Do not apply to watermelons.

CABBAGE, BROCCOLI, AND CAULIFLOWER - To control flea beetles, harlequin bug, armyworm, cabbage caterpillar (does not include cabbage looper), apply 14 to 28 pounds per acre. Repeat applications as required. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

BEANS - To control Mexican bean beetle, leafhoppers, corn earworm (larvae), and cucumber beetles, apply 14 to 20 pounds per acre.

POTATOES, EGGPLANT, AND PEPPERS - To control Colorado potato beetle, flea beetles, and leafhoppers, apply 14 pounds per acre to foliage only.

TOMATOES - To control tomato fruitworm, tomato cutworm, stink bug, armyworms, and climbing cutworms, apply 28 pounds per acre. For tomato fruitworm, apply to foliage when fruits of main crop are beginning to set; repeat in 2 and 4 weeks.

STRAWBERRIES - To control spittlebug, strawberry leaf roller, and strawberry weevil, apply 28 pounds per acre. For spittlebug make first application when small spittle musses first appear on or between the young leaves and repeat as needed. For strawberry leaf roller and strawberry weevil, apply to blossom buds and repeat as needed.

DO NOT APPLY TO STRAWBERRIES WITHIN 1 DAY OF HARVEST.

NOTICE

Seller makes no warranty of any kind, expressed or implied, concerning the use of this product by the buyer or others. Seller shall not be liable for any claims arising out of the use of such materials, whether used singly or in combination with other substances. Buyer assumes all risk from use or handling of this product. Use only for the purpose and in the manner stated in this label.

NET WEIGHT
50 POUNDS

MANUFACTURED BY
RING AROUND PRODUCTS, INC.
MONTGOMERY, ALA.
FORMERLY
STEGALL-SYLVEST SEED CO., INC.

PRODUCT NO. 1452

USDA Reg. No. 8934-7