

RING AROUND BRAND
10% CHLORDANE GRANULAR

INGREDIENTS	
ACTIVE INGREDIENTS:	
Technical Chlordane*-----	10.0%
INERT INGREDIENTS -----	90.0%
TOTAL	100.0%

*Equivalent to 6.0% Octachloro-4,7-methanotetrahydroindane and 4.0% related compounds.

CAUTION (in red and 18 pt.type)
KEEP OUT OF REACH OF CHILDREN (in red and 12 pt. type)
See additional CAUTIONS statements on front panel.

DIRECTIONS FOR USE

Preplanting Soil Applications

CUTWORMS-Apply 12½ to 15 lb.per acre (3 oz.per 500 sq.ft.) uniformly over surface of prepared land prior to setting of tomato, cabbage, pepper or tobacco transplants. If desired, place 1 to 1½ teaspoonfuls in a ring around each transplant but not in direct contact with each plant.

WIREWORMS-Apply 35 to 50 lb.per acre (½ to 2/3 lb.per 500 sq.ft.) uniformly over surface of prepared land, disk lightly into top 3 or 4 inches of soil ahead of planting of potatoes, corn, sweet corn, tobacco or strawberries.

CABBAGE MAGGOT-Apply 35 to 50 lb.per acre (½ to 2/3 lb.per 500 sq.ft.) work into top 3 inches of soil just before setting of cabbage, broccoli, or cauliflower transplants.

SEED CORN MAGGOT-Apply 12½ to 15 lb.per acre (3 oz.per 500 sq.ft.) work into top 3 inches of seed bed ahead of seeding of peas, beans, lima beans, sweet corn.

WHITE GRUBS-Uniformly apply 75 to 100 lb.per acre (1 lb.per 500 sq.ft.) of prepared soil and disk into top 4 to 6 inches of soil.

FOR USE ON LAWNS

CAUTION (in red and 18 pt.type)
Keep children and pets off treated areas until insecticide is washed into the soil and the grass is thoroughly dry.
DO NOT USE on newly seeded lawn.

ACCEPTED
4-26-67
8934-46

ANTS, CHINCH BUGS, CHIGGERS, SODWORMS, CRICKETS, CUTWORMS and EAWIGS-Apply 12½ to 15 lbs.per acre (3 oz.per 500 sq.ft.of area) directly into the grass. Repeat as necessary.

WHITE GRUBS and JAPANESE BEETLE GRUBS-Apply 75 lb.per acre (1 lb.per 500 sq.ft.of area) in early spring or late fall.

CRAB GRASS-Apply 1½ lb.per 100 sq.ft. or 15 lb.per 1000 sq.ft. in early spring before crab grass seed germinate. The time crab grass seed germinates varies with the seasonal temperature but in most areas, the application may be made between March 1 and April 15. Repeat annually if necessary.

CAUTION (in red and 18 pt.type)

Avoid inhalation of vapor or dust. Avoid contamination of feed or foodstuffs.

TOXIC to Fish and Wildlife. Do not contaminate any body of water nor apply when conditions favor runoff from the area treated.

BIRDS feeding on treated areas may be killed.

DO NOT graze or feed forage to dairy animals or animals being finished for slaughter.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of smoke).

NOTICE

Seller makes no warranty of any kind, expressed or implied, concerning the use of this product by the buyer or others. Seller shall not be liable for any claims arising out of the use of such materials, whether used singly or in combination with other substances. Buyer assumes all risk from use or handling of this product. Use only for the purpose and in the manner stated in this label.

NET WEIGHT
POUNDS

STEGALL - SYLVEST SEED CO., INC.
Montgomery, Ala.
USDA Reg.No.

PRODUCT NO. 2000

ACCEPTED
4-26-67
8934-46