

ACCEPTED

MAY 18 1967

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 29,344 SUBJECT TO ATTACHED COMMENTS.

CHLORDANE-8 EMULSIFIABLE CONCENTRATE

DIRECTIONS FOR USE

Dilute the concentrate in required volume of water by stirring or agitation; agitate during application. If allowed to stand, re-agitate before use.

USE ON SOIL

WHITE GRUBS, JAPANESE BEETLE LARVAE, AND GREEN JUNE BEETLE LARVAE—Use 2 to 3 quarts of concentrate (4 to 6 lb. chlordane) in sufficient water to spray 1 acre (4 tablespoonfuls per 1,000 sq. ft.) in turf or grassland (not used for hay or grazing). Spray directly into the grass—early spring or late fall. Repeat annually as necessary. DO NOT APPLY TO GRASS TO BE FED TO DAIRY ANIMALS OR ANIMALS BEING FINISHED FOR SLAUGHTER. DO NOT PASTURE SUCH ANIMALS ON TREATED TURF OR GRASSLAND. DO NOT APPLY TO FORAGE TO BE SOLD COMMERCIALY.

For strawberries, apply to the soil surface and work into top 3 inches of soil before setting plants.

WIREWORMS—Use 2 to 3 quarts of concentrate (4 to 6 lb. chlordane) in sufficient water to spray 1 acre (4 tablespoonfuls per 1,000 sq. ft.). Work into top 3 inches of soil prior to the planting of the following crops: broccoli, cabbage, cantaloupe, cauliflower, collard, eggplant, lettuce, onion, peas, peppers, potatoes, strawberries, tomatoes, turnips, and watermelon.

CUTWORMS—Use 1 to 1 1/2 pint of concentrate (1 to 1 1/2 lb. chlordane) in sufficient water to spray 1 acre (1/2 to 1 tablespoonfuls per 1,000 sq. ft.). Spray uniformly around base of tomato, cabbage, and pepper plants at time of planting.

ROOT MAGGOTS—Use 1/2 pint of concentrate for each 50 gallons of water (1 1/2 tablespoonfuls per gallon) as a transplanter solution on broccoli, cabbage, cauliflower, collard, and radish. Apply 1/4 cup of this solution around plants at time of transplanting.

USE ON PLANTS

STRAWBERRIES—To control spittlebug, crown borer, strawberry weevil, use 1 pint of concentrate (1 lb. chlordane) in sufficient water to spray 1 acre (1 tablespoonful per 1,000 sq. ft.). For spittlebug and crown borer, apply in early spring before blossoms appear. For strawberry weevil, apply spray to flower buds just before they open. DO NOT APPLY TO STRAWBERRIES AFTER FRUITS START TO FORM.

POTATOES—To control Colorado potato beetle, flea beetle, and blister beetle, use 1 to 1 1/2 pint (1 to 1 1/2 lb. chlordane) in sufficient water to spray 1 acre. Apply spray to foliage when insects first appear and repeat at 7 to 10-day intervals or as necessary.

CHIGGERS, SOD WEBWORM, AND ANTS on lawns or turf (not used for hay or grazing) use 1 to 1 1/2 pint (1 to 1 1/2 lb. chlordane) in sufficient water to spray grass on 1 acre (1 1/2 tablespoonfuls in water for each 1,000 sq. ft.). Apply when insects are first observed and repeat as necessary.

DO NOT FEED TREATED PLANTS OR ENSILAGE MADE FROM TREATED PLANTS TO POULTRY OR ANIMALS. DO NOT APPLY TO FOOD CROPS AFTER EDIBLE PARTS BEGIN TO FORM.



Ring Around Brand

CHLORDANE-8 EMULSIFIABLE CONCENTRATE

INGREDIENTS

Table listing ingredients: ACTIVE INGREDIENTS (Technical chlordane* 72%, Petroleum distillate 21%), INERT INGREDIENTS (7%), Total 100%.

* Equivalent to 43.2% octachloro-4,7-methanotetrahydroindane and 28.8% related compounds.

CONTAINS 8 POUNDS CHLORDANE PER GALLON

CAUTION

KEEP OUT OF REACH OF CHILDREN

See additional CAUTIONS statements on side panel

MANUFACTURED BY

STEGALL - SYLVEST SEED CO., INC.

Montgomery, Ala.

NET VOLUME U. S. GALLONS

USDA Reg. No. Applied For

PROD. 152

USE ON CRABGRASS

To control crabgrass, use 7 1/2 gallons of concentrate in sufficient water to spray 1 acre (2 quarts per 3,000 sq. ft.) of grass area (mowed or grazed) as golf fairways, parks, estates, and ball parks in early spring before crabgrass seeds germinate or sprout, with a second application between March 1 and April 15. Repeat annually if necessary. KEEP CHILDREN AND DOMESTIC ANIMALS OFF TREATED AREAS. INSECTICIDE HAS BEEN WASHED INTO THE SOIL BY RAINFALL AND THOROUGH SPRINKLING, AND THE GRASS HAS COMPLETED ITS GROWTH.

USE AGAINST TERMITES

TERMITES—Remove all wood and other cellulose materials from the soil. Prepare a water emulsion containing 1 percent (1/2 gallon of this concentrate in 50 gallons of water) and apply as follows:

(a) Under attached porches, platforms, and utility entrances, apply 1 gallon of water mixture per 10 square feet of soil surface.

(b) Along inside foundation walls and around piers, apply 1 gallon of water mixture per 5 linear feet.

(c) Along outside foundation walls, dig a trench 8 to 10 inches deep, but not below the footing; pour a water mixture evenly into the trench throughout its length, and apply additional water mixture. Repeat this process until the trench is filled. If foundation wall is shallow (15 inches or less) apply 2 gallons of the mixture per 5 linear feet; if deep (more than 15 inches) apply 4 gallons per 5 linear feet.

(d) Unexposed soil under terraces, patios, sidewalks, and porches, apply treatment close to foundation wall by drilling, or through cracks, so that soil along foundation wall will receive 1 gallon of water mixture per 5 linear feet.

(e) For soil treatment prior to pouring concrete slabs, apply 1 gallon of the water mixture per 10 square feet. Where slabs join or at service entrances, apply 2 gallons per 5 linear feet.

Inspect semi-annually and retreat if necessary.

USE FOR FLY REDUCTION

FLIES—To control adult houseflies and maggots, use 1 gallon of concentrate mixed in 5 gallons of water to spray outside of buildings, porches, and garbage cans and dumps. A minimum of 1 pint of spray should be used to cover 100 square feet of surface.

NECESSARY. DO NOT APPLY TO ANIMALS. DO NOT APPLY TO DAIRY BARN, MILK ROOM, OR OTHER AREAS WHERE FEED IS PROCESSED, HANDLED, OR STORED.

CAUTION -> 18

Harmful if swallowed. Contact with skin causes toxic reaction. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuff.

Do not use, pour, spill or store near heat or open flame.

TOXIC to Fish & Wildlife. Do not contaminate any body of water or other conditions favor runoff into the area treated.

NOTICE

Stegall makes no warranty of any kind, expressed or implied, and is not liable for any claims arising out of the use of such material whether used singly or in combination with other substances. It assumes all risk from use or handling of this product. Use only in the manner stated in this label.



Ring Around Brand

CHLORDANE-8
EMULSIFIABLE CONCENTRATE

INGREDIENTS

ACTIVE INGREDIENTS:	
Technical chlordane*	72%
Petroleum distillate	21%
INERT INGREDIENTS	7%
Total	
	100%

* Equivalent to 43.2% octachloro-4,7-methanotetrahydroindane and 28.8% related compounds.

CONTAINS 8 POUNDS CHLORDANE PER GALLON

CAUTION

KEEP OUT OF REACH OF CHILDREN

See additional CAUTIONS statements on side panel

MANUFACTURED BY

STEGALL - SYLVEST SEED CO., INC.
Montgomery, Ala.

NET VOLUME
U. S. GALLONS

USDA Reg. No. ~~Applied For~~

PROD. 152

USE ON CRABGRASS

To control crabgrass, use 7½ gallons of concentrate in sufficient water to spray 1 acre (2 quarts per 3,000 sq. ft.) of grass area (not used for hay or grazing) as golf fairways, parks, estates, and ball parks. Apply in early spring before crabgrass seeds germinate or sprout, which in most areas is between March 1 and April 15. Repeat annually if necessary. KEEP CHILDREN AND DOMESTIC ANIMALS OFF TREATED AREAS UNTIL INSECTICIDE HAS BEEN WASHED INTO THE SOIL BY RAINFALL OR THOROUGH SPRINKLING, AND THE GRASS HAS COMPLETELY DRIED.

USE AGAINST TERMITES

TERMITES—Remove all wood and other cellulose material in contact with the soil. Prepare a water emulsion containing 1 percent of chlordane by mixing ½ gallon of this concentrate in 50 gallons of water (1 pint in 12½ gallons) and apply as follows:

- (a) Under attached porches, platforms, and utility entrances, apply 1 gallon of water mixture per 10 square feet of soil surface.
 - (b) Along inside foundation walls and around piers, apply 2 gallons of water mixture per 5 linear feet.
 - (c) Along outside foundation walls, dig a trench 8 to 10 inches wide and 30 inches deep, but not below the footing; pour a portion of the water mixture evenly into the trench throughout its length, replace part of the soil, and apply additional water mixture. Repeat this process until trench is filled. If foundation wall is shallow (15 inches or less), apply no less than 2 gallons of the mixture per 5 linear feet; if foundations are deep (more than 15 inches) apply 4 gallons per 5 linear feet.
 - (d) Unexposed soil under terraces, patios, sidewalks, and driveways, apply treatment close to foundation wall by drilling, or through shrinkage cracks, so that soil along foundation wall will receive 2 gallons of water mixture per 5 linear feet.
 - (e) For soil treatment prior to pouring concrete slabs, apply 1 gallon of the water mixture per 10 square feet. Where slabs join inside walls or service entrances, apply 2 gallons per 5 linear feet.
- Inspect semi-annually and retreat if necessary.

USE FOR FLY REDUCTION

FLIES—To control adult houseflies and maggots, use 1 pint of concentrate mixed in 5 gallons of water to spray outside of buildings, manure piles, and garbage cans and dumps. A minimum of 1 pint of finished spray should be used to cover 100 square feet of surface. Repeat as necessary. DO NOT APPLY TO ANIMALS. DO NOT APPLY INSIDE DAIRY BARN, MILK ROOMS, OR OTHER AREAS WHERE FOOD IS PROCESSED, HANDLED, OR STORED.

CAUTION → 18 point

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

Do not use, pour, spill or store near heat or open flame.

TOXIC to Fish & Wildlife. Do not contaminate any body of water nor apply when conditions favor runoff from the area treated.

NOTICE

Seller makes no warranty of any kind, expressed or implied. Seller shall not be liable for any claims arising out of the use of such materials, whether used singly or in combination with other substances. Buyer assumes all risk from use or handling of this product. Use only for the purposes and in the manner stated in this label.