



ACCEPTED

MAY 20 1957

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON CONTROL
ED UNDER NO. 2034-1A
TO ATTACHED COMMISSION

Ring Around Brand

5% CHLORDANE DUST

INGREDIENTS

ACTIVE INGREDIENTS:	
*Technical Chlordane	5.0%
INERT INGREDIENTS	95.0%
TOTAL	100.0%

*Equivalent to 3% Octochloro-4,7-methanoindane and 2% related compounds.

WARNING

Red 18 point

KEEP OUT OF REACH OF CHILDREN *Red 12 point*

See additional CAUTIONS statements on front panel.

DIRECTIONS

FOR USE ON VEGETABLES: Cabbage—Imported Cabbage Worm, Cabbage Looper, Diamond Back Moth: Use at the rate of 2 oz. per 50 feet of row. Do not dust after edible parts start to form.
CUCURBITS—Squash Bug: Use at the rate of 3 to 4 oz. per 50 feet of row. Chlordane is injurious to certain varieties of cucurbits. Do not apply this material to susceptible varieties. Do not apply after edible parts start to form.
POTATOES—Colorado Potato Beetle and Flea Beetles: Use 2 oz. per 50 feet of row.
TOMATOES—Hornworm: Use 2 to 4 oz. per 50 feet of row. Do not use after fruit begins to form.

LAWNS

Apply the indicated amount of 5% Dust to 1000 square feet.
ARMYWORM, FALL ARMYWORM, CRICKETS, CHIGGERS, TICKS, ANTS, AND GRASSHOPPERS: Use 1 pound. Do not sprinkle for 48 hours.
MOLE CRICKETS: Sprinkle thoroughly before application. Apply 2 pounds.
EARWIGS: Apply 1 pound. Treat around foundations, along fences, around refuse and trash and lawn. Do not sprinkle for 48 hours.
CHINCH BUGS, FALSE CHINCH BUGS, AND CUTWORMS: Apply 2 pounds and sprinkle thoroughly after application.
SOD WEBWORMS or LAWN MOTHS: Sprinkle before application. Apply 2 1/2 pounds, then wait 48 hours before watering again.

SOIL INSECTS

ANTS, WHITE-FRINGED BEETLE LARVAE, JAPANESE BEETLE LARVAE, MOLE CRICKETS, WHITE GRUBS and WIREWORMS: On infested sites apply 1 pound per 1000 square feet. For control of these pests on vegetables such as beans, cabbage,

KEEP OUT OF REACH OF CHILDREN → *Rel 12 point*
See additional CAUTIONS statements on front panel.

DIRECTIONS

FOR USE ON VEGETABLES: Cabbage—Imported Cabbage Worm, Cabbage Looper, Diamond Back Moth: Use at the rate of 2 oz. per 50 feet of row. Do not dust after edible parts start to form.

CUCURBITS—Squash Bug: Use at the rate of 3 to 4 oz. per 50 feet of row. Chlorpyrifos is injurious to certain varieties of cucurbits. Do not apply this material to susceptible varieties. Do not apply after edible parts start to form.

POTATOES—Colorado Potato Beetle and Flea Beetles: Use 2 oz. per 50 feet of row.

TOMATOES—Hornworm: Use 2 to 4 oz. per 50 feet of row. Do not use after fruit begins to form.

LAWNS

Apply the indicated amount of 5% dust to 1000 square feet.
ARMYWORM, FALL ARMYWORM, CRICKETS, CHIGGERS, TICKS, ANTS, AND GERMAN COCKROACHS: Use 1 pound. Do not sprinkle for 48 hours.

MOLE CRICKETS: Sprinkle thoroughly before application. Apply 2 pounds.

LAWN GRUBS: Apply 1 pound. Treat around foundations, along fences, around refuse and trash and lawn. Do not sprinkle for 48 hours.

CHINCH BUGS, FALSE CHINCH BUGS, AND CUTWORMS: Apply 2 pounds and sprinkle thoroughly after application.

SOIL WORMS or LAWN MOTHS: Sprinkle before application. Apply 2 1/2 pounds, then wait 48 hours before watering again.

SOIL INSECTS

ANTS, WHITE-FRINGED BEETLE LARVAE, JAPANESE BEETLE LARVAE, MOLE CRICKETS, WHITE GRUBS and WIREWORMS: On infested sites apply 1 pound per 1000 square feet. For control of these pests on vegetables such as beans, cabbage, potatoes, sweet potatoes, corn and turnips, apply to the soil surface before planting and work into the upper 2 or 3 inches of soil. On flowers, shrubs and lawns, apply either before planting as above or to existing plantings, working the material into the upper few inches of soil.

HOUSEHOLD

ANTS: Dust areas from which ants emerge such as cracks in floors, walls, around drain pipes, etc. Dust on areas where ants crawl such as window sills, drainboards, and along floors. Repeat applications as necessary.

ROACHES: Dust into cracks and crevices and all points where roaches gain entrance to the house. Dust near sinks, drainboards, behind wall cabinets, beneath refrigerators and other similar places where roaches hide in the day. Repeat applications as necessary.

CAUTION → *18 point*

Avoid breathing dust or vapor. Do not get on skin, in eyes or on clothing. Wash thoroughly after using. Do not contaminate feed, foodstuffs or drinking water.

TOXIC to Fish and Wildlife. Do not contaminate any body of water nor apply when conditions favor runoff from the crop area treated. Do not feed treated forage to dairy animals or animals being finished for slaughter.

NOTICE

Seller makes no warranty of any kind, express or implied, concerning the use of this product by the buyer or others. Seller shall not be liable for any claims arising out of the use of such materials, whether used singly or in combination with other substances. Buyer assumes all risk from use or handling of this product. Use only for the purpose and in the manner stated in this label.

Manufactured By
STEGALL - SYLVEST SEED CO., INC.

NET WEIGHT
4 POUNDS

Montgomery, Ala.
USDA Reg. No.

8004-S. P.