



ACCEPTED

MAY 3 1 1967

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER
 ED UNDER NO. 38-56 SUBJECT
 TO ATTACHED COMMENTS.

Ring Around Brand MALATHION WETTABLE POWDER

INGREDIENTS	
ACTIVE INGREDIENTS:	
Malathion*	25%
INERT INGREDIENTS	75%
	Total
	100%

*O,O-dimethyl dithiophosphate of diethylmercaptosuccinate

CAUTION
KEEP OUT OF REACH OF CHILDREN
 See additional CAUTIONS Statements on front panel.

DIRECTIONS FOR USE

To prepare sprays, fill tank about one-quarter full with water, add the correct amount of powder, and agitate thoroughly to obtain a good suspension. Then add the remainder of the water and continue agitation or stirring. Some agitation is required to maintain the suspension while spraying.

A NUMBER OF INSECTS ON THE CROPS INDICATED MAY BE CONTROLLED BY PROPER AND TIMELY SPRAYS. DOSAGES STATED BELOW ARE IN POUND OF POWDER PER 100 GALLONS OF WATER, UNLESS SPECIFIED OTHERWISE.

VEGETABLES

BEANS — Mexican bean beetle, leafhoppers, mites, use 2 pounds; repeat as needed. Do not apply within 3 days of harvest.

BEETS AND SPINACH — Leafminers, use 2 pounds; apply spray before miners have entered leaves. Do not apply within 7 days of harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, AND RELATED CROPS — Aphids; use 2 pounds; if heavy populations are established, use 4 pounds. Repeat as necessary. For cabbage worms use 6 pounds; repeat as necessary. Do not apply within 7 days of harvest.

CUCUMBERS, SQUASH, PUMPKIN, AND MUSKMELON — Aphids; use 2 pounds. DO NOT apply sprays to wet plants. Do not apply within 3 days of harvest.

EGG PLANT — Aphids, use 2 pounds, repeat as necessary; not within 3 days of harvest.

ONIONS — Thrips; use 2 pounds per 100 gallons when first noticed, or 4 pounds in sufficient water to cover 1 acre, repeat in one week and again if they reappear. Make thorough applications, but not within 3 days of harvest.

PEAS AND PEPPERS — Aphids; use 2 pounds. Repeat as necessary, but not within 3 days of harvest.

POTATOES — Colorado potato beetle, flea beetle; use 2 pounds; repeat as necessary. Aphids; use

DIRECTIONS FOR USE

To prepare sprays, fill tank about one-quarter full with water, add the correct amount of powder and agitate thoroughly to obtain a good suspension. Then add the remainder of the water and continue agitation or stirring. Some agitation is required to maintain the suspension while spraying.

A NUMBER OF INSECTS ON THE CROPS INDICATED MAY BE CONTROLLED BY PROPER AND TIMELY SPRAYS. DOSAGES STATED BELOW ARE IN POUND OF POWDER PER 100 GALLONS OF WATER UNLESS SPECIFIED OTHERWISE.

VEGETABLES

BEANS — Mexican bean beetle, leafhoppers, mites; use 2 pounds per 100 gallons of water. Do not apply within 3 days of harvest.

BEETS AND SPINACH — Leafminers; use 2 pounds; apply spray before miners have started feeding. Do not apply within 7 days of harvest.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, AND RELATED CROPS — Aphids; use 2 pounds, if heavy populations are established, use 4 pounds. Repeat as necessary. For cabbage worms; use 6 pounds; repeat as necessary. Do not apply within 7 days of harvest.

CUCUMBERS, SQUASH, PUMPKIN, AND MUSKMELON — Aphids; use 2 pounds. DO NOT apply sprays to wet plants. Do not apply within 3 days of harvest.

EGG PLANT — Aphids; use 2 pounds, repeat as necessary, not within 3 days of harvest.

ONIONS — Thrips; use 2 pounds per 100 gallons when first noticed, or 4 pounds in sufficient water to cover 1 acre, repeat in one week and again if they reappear. Make thorough applications, but not within 3 days of harvest.

PEAS AND PEPPERS — Aphids; use 2 pounds. Repeat as necessary, but not within 3 days of harvest.

POTATOES — Colorado potato beetle, flea beetle; use 2 pounds; repeat as necessary. Aphids; use 2 pounds for light populations and 4 pounds for heavy populations. Repeat as necessary, but not within 3 days of harvest.

TOMATO — Aphids; use 2 pounds, repeat as necessary, but not within 5 days of harvest.

CAUTION: DO NOT apply malathion with Bordeaux mixture or fixed copper. DO NOT apply to vegetable crops later than the times indicated.

MEASUREMENTS FOR SMALL EQUIPMENT

To convert dosages expressed in pounds per 100 gallons of water, use the following quantities for 1 gallon:

POUNDS PER 100 GALLONS

1 Pound
2 Pounds
3 Pounds
4 Pounds

QUANTITY PER GALLON

$\frac{3}{4}$ Tablespoonsful
 $1\frac{1}{2}$ Tablespoonsful
 $2\frac{1}{4}$ Tablespoonsful
3 Tablespoonsful

CAUTION

HARMFUL IF SWALLOWED. Avoid breathing dust or spray mist. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed or foodstuffs. Keep out of reach of children.

Fish may be killed if their waters are contaminated with this product. DO NOT apply when weather conditions favor drift of spray or dust from crop area treated.

NOTICE

Seller makes no warranty of any kind, expressed or implied, concerning the use of this product by the buyer or others. Seller shall not be liable for any claims arising out of the use of such materials, whether used singly or in combination with other substances. Buyer assumes all risk from use or handling of this product. Use only for the purpose and in the manner stated in this label.

NET WEIGHT

STEGALL - SYLVEST SEED CO., INC.
Montgomery, Ala.

PRODUCT NO. 3001