



Ring Around Brand

GAMMA KIL - BENZENE HEXACHLORIDE
(Wettable)

ACTIVE INGREDIENTS:

Gamma isomer of Benzene Hexachloride(BHC)-- 6.0%

Other isomer of Benzene Hexachloride(BHC)--30.0%

INERT INGREDIENTS -----64.0%

TOTAL 100.0

WARNING

KEEP OUT OF REACH OF CHILDREN

See additional CAUTIONS statements on front panel.

DIRECTIONS

DOGS

SARCOPTIC MANGE and FLEAS: Dip animals in a suspension, kept well mixed, using 1/2 pound to 15 gallons of water (1 cupful to 5 gallons). Make two treatments 1 week apart. Do not apply to young animals under 3 months of age or to emaciated, sick or lactating animals.

RESIDUAL SPRAYS

DO NOT use in or around dairy barns.

STABLE FLY, MOSQUITOES, ROACHES, ANTS: For external surfaces only, of barns, hog pens, stables and other building, use 3 1/2 pounds in 10 gallons of water and apply at the rate of 1 gallon to 1000 square feet of surface. Thoroughly cover resting surfaces of flies and mosquitoes. Apply to cracks, crevices and other hiding places for control of roaches and ants. Repeat applications as necessary. Do not contaminate feed or drinking water.

VEGETABLES

CABBAGE: ~~Apply at the rate of 4 pounds per 100 gallons of water.~~ Cabbage Aphids and Thrips—Use 4 pounds per 100 gallons of water. Do not apply more than 200 gallons per acre. Do not apply to ~~vegetables~~ after seedling stage.

PEPPERS: Aphids and Serpentine Leaf Miner—Use 4 pounds per 100 gallons of water. Do not apply more than 150 gallons per acre. Do not apply after edible parts begin to form.

DO NOT use on crops in rotation with root crops.

TICKS and CHIGGERS: Apply at the rate of 33 pounds per acre in sufficient water for coverage. Make application to weeds and underbrush harboring pests. Do not use on food crops or grazing areas. Repeat application if necessary.

CAUTION

HARMFUL IF SWALLOWED. May be absorbed through

