

ACCEPTED
 11/25/68
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND FUNGICIDE ACT
 FOR ECONOMIC PEST CONTROL
 REGISTERED UNDER NO. 14307-2
 SUBJECT TO ATTACHED COMMENTS.



Ring Around Brand

10% HEPTACHLOR GRANULAR

INGREDIENTS

ACTIVE INGREDIENTS:	
Heptachlor*	10.0%
Related compounds	3.8%
INERT INGREDIENTS	86.2%
<hr/>	
Total	100.0%

*Heptachlorotetrahydro-4, 7-methanoindene

WARNING

KEEP OUT OF REACH OF CHILDREN

See additional CAUTIONS statements on front panel

DIRECTIONS FOR USE

PRE-PLANTING SOIL APPLICATIONS

CORN, WHEAT, OATS, RYE, TOBACCO—To control wireworms, apply 20 to 30 pounds per acre (2 to 3 pounds heptachlor) as a uniform broadcast treatment of the soil prior to planting, and work into the top few inches of surface.

To control rootworms on corn, apply 10 pounds per acre (1 pound heptachlor) as a uniform broadcast treatment of the soil prior to planting and work into the top few inches of surface.

NON-CROP APPLICATIONS

LAWNS OR TURF AND VEGETATION IN NON-CROP AREAS—To control ants, chiggers, ticks, sod webworms, and cutworms, in grass and low-growing brush in areas where no crop is growing, apply 20 pounds per acre (4 ounces per 500 sq. ft.) directly into the grass or soil. Repeat as necessary.

To control white grubs and Japanese beetle larvae in turf, apply 30 to 37.5 pounds uniformly per acre (6.5 ounces per 500 sq. ft.) directly into the grass in early spring or late fall. Repeat treatment as new infestations develop, or as necessary.

CAUTION

DO NOT PERMIT CHILDREN AND PETS TO GO ON TREATED AREAS UNTIL THE INSECTICIDE HAS BEEN WASHED OFF THE GRASS AND INTO THE SOIL AND THE GRASS HAS COMPLETELY DRIED. DO NOT TREAT WHEN THE SOIL IS WET. APPLY BEFORE RAIN OR SPRINKLING.

non Crop Areas

Ornamental Turf

WARNING

KEEP OUT OF REACH OF CHILDREN

See additional CAUTIONS statements on front panel

DIRECTIONS FOR USE

PRE-PLANTING SOIL APPLICATIONS

CORN, WHEAT, OATS, RYE, TOBACCO—To control wireworms, apply 20 to 30 pounds per acre (2 to 3 pounds heptachlor) as a uniform broadcast treatment of the soil prior to planting, and work into the top few inches of surface.

To control rootworms on corn, apply 10 pounds per acre (1 pound heptachlor) as a uniform broadcast treatment of the soil prior to planting and work into the top few inches of surface.

NON-CROP APPLICATIONS

LAWNS OR TURF AND VEGETATION IN NON-CROP AREAS—To control ants, chiggers, ticks, sod webworms, and cutworms, in grass and low-growing brush in areas where no crop is growing, apply 20 pounds per acre (4 ounces per 500 sq. ft.) directly into the grass or soil. Repeat as necessary.

To control white grubs and Japanese beetle larvae in turf, apply 30 to 37.5 pounds uniformly per acre (6.5 ounces per 500 sq. ft.) directly into the grass in early spring or late fall. Repeat treatment as new infestations develop, or as necessary.

CAUTION

DO NOT PERMIT CHILDREN AND PETS TO GO ON TREATED AREAS UNTIL THE INSECTICIDE HAS BEEN WASHED OFF THE GRASS AND INTO THE SOIL AND THE GRASS HAS COMPLETELY DRIED. DO NOT TREAT WHEN THE SOIL IS WET. APPLY BEFORE RAIN OR SPRINKLING.

DO NOT graze or feed turf to livestock,

DO NOT apply to pasture or range lands.

This product is TOXIC to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

WARNING

MAY BE FATAL IF SWALLOWED! Do not breathe dust or vapor.

Do not get on skin, in eyes or on clothing. In case of contact wash immediately with soap and water. Do not contaminate feed and foodstuffs.

NOTICE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NET WEIGHT
50 POUNDS

STEGALL - SYLVEST SEED CO., INC.
Montgomery, Ala.

PRODUCT NO. 103045

USDA Reg. No. 8834-32

non Crop Areas
Ornamental Turf