

33-1/3 LBS. NET



(BAR-FLY)

BEEF CATTLE PASTURE FEED BLOCK MEDICATED

Prevents the breeding of horn flies and face flies in the manure of treated cattle, when fed as directed. Aids in prevention of stomach-worms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.), nodular-worms (*Oesophagostomum* spp.), and large-mouth bowllworms (*Chabertia* spp.); hookworms (*Bunostomum* spp.) in cattle. Aids in the prevention of cattle foot rot and soft tissue lumpy jaw. A nutritional source of iodine when fed to cattle as directed.

PESTICIDALLY ACTIVE DRUG INGREDIENT:

Phenothiazine*	1.15%
INERT ** INGREDIENT	98.85%
	100.00%

* Biologically assayed.
** Refers only to ingredients which are neither insecticides nor drugs.

(OTHER) ACTIVE DRUG INGREDIENT:

Ethylene Diamine Dihydroiodide	0.022%
--------------------------------	--------

GUARANTEED ANALYSIS

CRUDE PROTEIN, not less than	14.00%
CRUDE FAT, not less than	1.00%
CRUDE FIBER, not more than	8.00%
CALCIUM (Ca), not less than	2.00%
CALCIUM (Ca), not more than	3.00%
PHOSPHORUS (P), not less than	1.00%
IODINE (I), not less than	0.0176%
SALT (NaCl), not less than	12.00%
SALT (NaCl), not more than	14.00%
VITAMIN A, not less than	10,000 USP Units/Lb.
VITAMIN D ₂ , not less than	10,000 USP Units/Lb.

INGREDIENTS: Soybean Meal, Cottonseed Meal, Ground Yellow Corn, Ground Wheat Middlings, Molasses, Molasses, Dried Cane Molasses, Wheat Mill Feed, Calcium Sulfate, Calcium Carbonate, Alfalfa, Dicalcium Phosphate, Sulfate, Vitamin A Palmitate, Inactivated Yeast, Source of Vitamin D₂, Vitamin F Supplement, Ethylene Diamine Dihydroiodide, Nicotinic Acid, Calcium Pantothenate, Choline Chloride, Manganese Oxide, Magnesium Oxide, Copper Oxide, Zinc Oxide, Cobalt Carbonate, Iron Oxide and Anise Oil.

Directions: Feed 1/2 pound of (Bar-Fly) Beef Cattle Pasture Feed Block Medicated per day for each 1,000 pounds of body weight. Feed continuously at the rate of one pound of salt per 100 pounds of body weight. Feed carefully to avoid any loss of product. Feed one pound per day of this block for each 10 animals. Apply to all animals in the herd. For details of use, see the label on the product.

WARNING:

KEEP OUT OF REACH OF CHILDREN. DO NOT FEED TO PASTURE GRAZING DAIRY ANIMALS. OCCASIONAL FEEDING TO DAIRY ANIMALS HAS BEEN REPORTED IN CATTLE.

Distributed By:

SUPREMA MANUFACTURING, INC., Kansas City, Mo.
816-231-1111 EPA Reg. No. 149-100-01