

**BEEF CATTLE PASTURE
FEED BLOCK**
Medicated



Prevents the breeding of horn flies and face flies in the manure of treated cattle, when fed as directed; Aids in prevention of stomachworms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.); nodularworms (*Oesophagostomum* spp.); and large-mouth bowelworms (*Chabertia* spp.); hookworms (*Bunostomum* spp.) in cattle; Aids in the prevention of cattle foot rot and soft tissue lumpy jaw. A nutritional source of iodine when fed to cattle as directed.

PESTICIDALLY ACTIVE DRUG INGREDIENT:

Phenothiazine*	1.15%
INERT** INGREDIENT	98.85%
Total	100.00%

* Biologically assayed.

**Refers only to ingredients which are neither insecticides nor drugs.

(OTHER) ACTIVE DRUG INGREDIENT:

Ethylene Diamine Dihydroiodide	0.022%
--------------------------------	--------

GUARANTEED ANALYSIS:

CRUDE PROTEIN, not less than	14.00%
CRUDE FAT, not less than	1.00%
CRUDE FIBER, not more than	8.00%
CALCIUM (Ca), not less than	3.00%
CALCIUM (Ca), not more than	4.00%
PHOSPHORUS (P), not less than	2.00%
IBUFENE (I), not less than	0.0176%
SALT (NaCl), not less than	12.00%
SALT (NaCl), not more than	14.00%
VITAMIN A, not less than 40,000 USP UNITS PER POUND.	
VITAMIN B2, not less than 10,000 USP UNITS PER POUND.	

INGREDIENTS: Soybean Meal, Cottonseed Meal, Ground Yellow Corn, Ground Milo, Alfalfa Meal, Wheat Middlings, Cane Molasses, Dried Cane Molasses, Soybean Mill Feed, Dicalcium Phosphate, Calcium Carbonate, Salt, Lignin Sulfonate, Smectite Vermiculite, Vitamin A Palmitate, Irradiated Dried Yeast (Source of Vitamin D2), Vitamin E Supplement, Ethylene Diamine Dihydroiodide, Niacin, Calcium Pantothenate, Choline Chloride, Iron Oxide, Manganese Oxide, Magnesium Oxide, Copper Oxide, Iron Carbonate, Cobalt Carbonate, Zinc Oxide and Anise Oil.

Distributed By

HALES & HUNTER CO., General Office Chicago, Illinois 60606 **AF**

**PIONEER® BEEF CATTLE
PASTURE FEED BLOCK**
Medicated

NET WEIGHT 33 1/3 LBS.

FEEDING DIRECTIONS

1. Feed 1/2 pound of PIONEER Beef Cattle Pasture Blocks Medicated per day for each 1,000 pounds of body weight.
2. Self feed continuously as the sole source of salt.
3. Observe cattle consumption rate carefully to assure consumption at the directed rate per animal per day.
4. Use 1 block for each 10 animals. Maintain block supply at all times for adequate control.
5. Locate blocks near the animals' water supply or feeding area.
6. Phenothiazine is not effective against existing adult flies. Spray cattle at least once in early summer and feed PIONEER Beef Cattle Pasture Block Medicated on a year-around basis.

WARNING

KEEP OUT OF REACH OF CHILDREN. DO NOT FEED TO LACTATING DAIRY ANIMALS. Occasional Sensitivity To Phenothiazine Has Been Reported In Cattle.

EPA Reg. No. 8939-10