

Blair's **CERTIFIED BRAND** NET WT 25 1/3 Lbs.
BAR-FLY BEEF CATTLE
PASTURE FEED BLOCK

MEDICATED

Prevents the breeding of horn flies and face flies in the pasture of treated cattle, when fed as directed. Aids in prevention of stomach worms (Haemonchus, Ostertagia, Trichostrongylus spp.); nodular worms (Oesophagostomum spp.); and large-mouth bowelworms (Chabertia spp.); hookworms (Bunostomum spp.) in cattle; Aids in the prevention of cattle foot rot and soft tissue lumpy jaw; A nutritional source of iodine when fed to cattle as directed.

PESTICIDALLY ACTIVE DRUG INGREDIENT:

Phenothiazine*	1.15%
INERT ** INGREDIENT.	98.85%
	100.00%

* Biologically assayed.
 ** Refers only to ingredients which are neither insecticides nor drugs.

(OTHER) ACTIVE DRUG INGREDIENT:

Ethylene Diamine Dihydroiodide	0.022%
--	--------

GUARANTEED ANALYSIS

CRUDE PROTEIN, not less than	14.00%
CRUDE FAT, not less than	1.00%
CRUDE FIBER, not more than	8.00%
CALCIUM (Ca), not less than	2.00%
CALCIUM (Ca), not more than	3.00%
PHOSPHORUS (P), not less than	1.00%
IODINE (I), not less than	0.0176%
SALT (NaCl), not less than	12.00%
SALT (NaCl), not more than	14.00%
VITAMIN A, not less than	80,000 USP Units/Lb.
VITAMIN B2, not less than	10,000 USP Units/Lb.

INGREDIENTS: Soybean Meal, Cottonseed Meal, Ground Yellow Corn, Ground Milk, Wheat Middlings, Alfalfa Meal, Cane Molasses, Dried Cane Molasses, Soybean Mill Feed, Dicalcium Phosphate, Calcium Carbonate, Salt, Magnesium Sulfate, Smeectite Vermiculite, Vitamin A Palmitate, Irradiated Yeast (Source of Vitamin B2), Vitamin B Supplement, Ethylene Diamine Dihydroiodide, Nickel, Calcium Phosphate, Choline Chloride, Iron Oxide, Manganous Oxide, Magnesium Oxide, Copper Oxide, Iron Carbonate, Zinc Carbonate, Zinc Oxide and Arise 211.

FEEDING DIRECTION: Feed 1/2 lb. of Bar Fly Beef Cattle Pasture Feed Block Medicated per day for each 1,000 lbs. of live weight. Let feed continuously as the sole source of feed for the cattle. Consumption rate carefully to assure they are consuming the product at the intended rate per day. Use 1 block for 100 head of animals. Maintain block supply at all times for adequate supply. Locate block near the animals' water supply or loafing area. Phenothiazine is effective against existing fly flies. Since it is not a larvicide, use in early summer and feed Bar Fly Beef Cattle Pasture Feed Block continuously on a year round basis. Since face flies migrate from farm to farm, it is essential that the larvicide be used to keep fly out of an area, to assist in reduction of adult fly population. Consult veterinarian before using in severely debilitated animals.

WARNING: KEEP OUT OF REACH OF CHILDREN. DO NOT FEED TO LACTATING DAIRY ANIMALS. Occasional Sensitivity To Phenothiazine has been Reported In Cattle.

MANUFACTURED FOR
THE BLAIR MILLING & ELEVATOR CO.
 ARDEN, MISSOURI 64601 EPA Reg. No. 100-100-01