

40 LBS. NET



BAR-FLY BLOCK No. 185 MEDICATED

Prevents the breeding of horn flies and face flies in the manure of treated cattle when used as directed. Also prevents the invasion of stomachworms (Haemonchus, Ostertagia, and Trichostrongylus), nodularworms (Oesophagostomum spp.), and lungworms (Dictyocaulus) (Chabertia spp.); hookworms (Bunostomum spp.); and prevents the prevention of cattle foot rot and soft foot. Also a new and useful nutritional source of iodine when fed to cattle as directed.

PESTICIDALLY ACTIVE DRUG INGREDIENT:

Phenothiazine*	2.00%
INERT ** INGREDIENTS:	98.00%
TOTAL	100.00%

* Biologically assayed.
** Refers only to ingredients which are neither insecticides nor drugs.

(OTHER) ACTIVE DRUG INGREDIENT:

Ethylene Diamine Dithiodiiodide	0.002%
---------------------------------	--------

GUARANTEED ANALYSIS

CRUDE PROTEIN, not less than	5.00%
CRUDE FAT, not less than	.50%
CRUDE FIBER, not more than	6.00%
PHOSPHORUS (P), not less than	1.00%
CALCIUM (Ca), not less than	7.00%
CALCIUM (Ca), not more than	8.00%
SALT (NaCl), not less than	14.00%
SALT (NaCl), not more than	22.00%
IODINE (I), not less than	0.035%
VITAMIN A, not less than	40,000 IU Units Per Lb.
VITAMIN D ₃ , not less than	20,000 IU Units Per Lb.

INGREDIENTS: Cane Molasses, Ground Molasses, Alfalfa Meal, Wheat Middlings, Cottonseed Meal, Lignin Sulfonate, Sweet or Vermilionite, Salt, Dicalcium Phosphate, Calcium Carbonate, Vitamin A Palmitate, Ferrous Sulfate, Ferrous Gluconate, Ferrous Fumarate, Ferrous Citrate, Ferrous Oxide, Manganese Oxide, Iron Carbonate, Copper Oxide, Cobalt Carbonate, Magnesium Oxide and Zinc Oxide.

DIRECTIONS: Feed 1/4 pound (1/2 ounce) of Bar-Fly Block No. 185 Medicated daily for each 1,000 lbs. of body weight. For calves, feed 1/4 pound per day for each 300 lbs. animal. Provide the animal with a clean water supply and a loafing area. If feed is not available, the animal should be given water and a loafing area. The animal should be kept in a clean, dry, well-ventilated area. The animal should be kept in a clean, dry, well-ventilated area. The animal should be kept in a clean, dry, well-ventilated area.

WARNING: KEEP OUT OF REACH OF CHILDREN. DO NOT FEED TO LACTATING DAIRY ANIMALS. Occasional Sensitivity to Phenothiazine Has Been Reported In Cattle

Manufactured by
M. W. M. Co.
St. Louis, Mo.