

33 1/3 LBS. NET



# BAR-FLY SWEET BLOCK MEDICATED

Prevents the breeding of horn flies and face flies in the manure of treated cattle when used as directed. Aids in the prevention of stomachworms (Haemonchus, Ostertagia, and Trichostrongylus spp.) nodularworms (Oesophagostomum spp.); and large-mouth bowelworm (Chabertia spp.); hookworms (Bunostomum spp.) in cattle; Aids in the prevention of cattle foot rot and soft tissue lumpy jaw in cattle; nutritional source of iodine when fed to cattle as directed.

**PESTICIDALLY ACTIVE DRUG INGREDIENT:**

Phenothiazine\* . . . . . 2.00%

INERT \*\* INGREDIENT . . . . . 98.00%

TOTAL 100.00%

\* Biologically assayed.

\*\* Refers only to ingredients which are neither insecticides nor drugs.

**(OTHER) ACTIVE DRUG INGREDIENT:**

Ethylene Diamine Dihydriodide. . . . . 0.044%

**GUARANTEED ANALYSIS**

CRUDE PROTEIN, not less than. . . . .	5.00%
CRUDE FAT, not less than. . . . .	.50%
CRUDE FIBER, not more than. . . . .	12.50%
PHOSPHORUS (P), not less than. . . . .	1.50%
CALCIUM (Ca), not less than. . . . .	3.00%
CALCIUM (Ca), not more than. . . . .	4.00%
SALT (NaCl), not less than. . . . .	19.00%
SALT (NaCl), not more than. . . . .	22.00%
IODINE (I), not less than. . . . .	0.035%
TOTAL SUGARS (as invert), not less than. . . . .	24.00%
VITAMIN A, not less than. . . . .	100,000 USP UNITS PER LB.
VITAMIN D2, not less than. . . . .	10,000 USP UNITS PER LB.

**INGREDIENTS:** Cane Molasses, Dried Cane Molasses, Soybean Mill Feed, Ground Milo, Cottonseed Meal, Soybean Meal, Animal Fat, BHA (a preservative), Salt, Dicalcium Phosphate, Vitamin A Palmitate, Irradiated Dried Yeast (Source of Vitamin D2), Ethylene Diamine Dihydriodide, Smectite Vermiculite, Manganous Oxide, Iron Carbonate, Copper Oxide, Cobalt Carbonate, Magnesium Oxide and Zinc Oxide.

**FEEDING DIRECTIONS:** Self feed 1/2 ounce of Bar-Fly Sweet Block Medicated per day for each 100 pounds of body weight. This is equivalent to 1/4 pound per day for each 800 pound animal. Locate blocks near the animals' water supply or loafing area. Self feed continuously as the sole source of salt. Observe cattle consumption rate carefully to assure consumption is at the directed rate per animal per day. Use 1 block for each 10 animals. Maintain block supply at all times for adequate control. Phenothiazine is not effective against existing adult flies. Spray cattle at least once in early summer and feed Bar-Fly Sweet Blocks until frost. For internal parasite control feed Bar-Fly Sweet Blocks on a year round basis.

**WARNING: KEEP OUT OF REACH OF CHILDREN. DO NOT FEED TO LACTATING DAIRY ANIMALS. Occasional Sensitivity To Phenothiazine Has Been Reported In Cattle.**

CODE 182

Manufactured By: EPA Reg. No. 9208-1  
SUPREMA MILLS, INC.  
Kansas City, Missouri 64132

