


ACCEPTED
FEB 5 1969
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 21422 SUBJECT
TO ATTACHED COMMENTS.



SUP RO CHLORDANE

10% GRANULAR INSECTICIDE

INGREDIENTS

ACTIVE INGREDIENTS:

*Technical chlordane 10%

INERT INGREDIENTS:

Perlite 90%
TOTAL 100%

*Equivalent to 6% Octachloro-4, 7-Methanotetrahydroindane and 4% related compounds.

CAUTION

INGREDIENTS

ACTIVE INGREDIENTS:

*Technical chlordane 10%

INERT INGREDIENTS:

Perfls 90%
TOTAL 100%

*Equivalent to 6% Octachloro-4, 7-Methanotetrahydroindane and 4% related compounds.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See back panel for antidote and other precautionary statements.

MANUFACTURED BY

LITE WEIGHT PRODUCTS, INC.

707 Funston Road

Kansas City, Kansas 66115

U. S. D. A. REGISTRATION NO. 8940-2

NET WEIGHT 5 LBS.

DIRECTIONS

INSECT CONTROL ON HOME LAWNS: For the control of Sod Webworms, White Grubs, Ants, Armyworms, Box-elder Bugs, Chiggers, Chinch Bugs, Crickets, Rose Chafer Grubs, Spiders, Ticks, Wireworms. On home lawns apply 1¼ - 2 lbs. of SUP-RO Chlordane 10% Granular for each 1,000 sq. ft. to cover the surface uniformly using a lawn fertilizer spreader. Treatment should be applied in the spring, or when insects observed to be troublesome. Immediately after treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce hazard to children and pets. Do not treat when soil is wet. Apply before rain or sprinkling.

INSECT CONTROL ON TURF, NURSERY AND GREENHOUSE SOILS: For the control of the above listed insects in Turf. (such as Golf Courses, Athletic Fields, Large Estate Nurseries, Cemeteries and Park Sod), and in Nursery and Greenhouse soils where ornamental plants are grown, use at the rate of 60 lbs. of SUP-RO Chlordane 10% granular per acre (1¼ - 2 lbs. to each 1,000 sq. ft.). For Cutworms, Earthworms, Cicada Killer Wasps, 3 times above rate. To cover surface uniformly use a lawn fertilizer spreader. Treatment should be applied in the spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling.

CAUTION: Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust and spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep children or pets off treated area until material has been washed into the soil by watering or rain, and grass is thoroughly dry.

ANTIDOTE: External—In case of spillage on skin, wash promptly with soap and water.
Internal—Emetic of 1 tablespoon mustard in tumbler of water. Call a Physician.

.....

This product is toxic to fish and wildlife.
Birds feeding on treated areas may be harmed.
Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

MANUFACTURED BY
LITE WEIGHT PRODUCTS, INC.
707 Funston Road
Kansas City, Kansas 66115

ACCEPTED
2/5/69
UNCL. THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR ECONOMIC PEST CONTROL REGISTERED
ED UNDER NO. 84002 SUBJECT
TO ATTACHED COMMENTS.