



50 LBS. NET WT.

THE MACK LAWN 1000 12-4-4

is a highly effective combination of chamical and restoral facilities with immediate. When combined an excenterability product will food your grows it control white grade and the before control and testing.







CHARANTEED ANALYSIS

	11.5% 3%				
Water inschille Nivegen Analishio Phespheric Acid (PiOs) Sahida Patash (KiO)					4/
Primary Flust Foods durined from					
Subhate of Ammeric, Common	ion Phosphots	, Sullub	d No.	A Magnesia, To	~ 4
Bland Mool, Sono Mool, and Cal	beend Mani				
				. ,	
Magnesium					
Magnesium					15
Magnesium Suffer (combined) Iron			-		. 151 . 1.
Magnesium Suffer (combined) Iron			-		. 151 . 1.

Secondary Plant Feed: & Trace Stamons derived from:
Sulphate of Patesh Magnesia, tran Sulphate, Managenese Sulphate, Zinc Sulphate & EDTA
tran Chalate. Sulfur and Magnesium derived from Sulfate of Patesh Magnesia and Ammanium Sulfate.

Natural Plant Feed derived from Bane Meel, Stand Meel, Collegeed Meel & Tung Meel.

Marium Soffete.

Natural Plant Food derived from Bane Meel, Blood Meel, Coffeeneed Meel & Tung Meel.

ACTIVE INGREDIENTS
Technical Chlordane*

INGREDIENTS
TOTAL

105.00%

Marinton to 21% (Secretary & Zandarana-Audaindase and 14% Belated Companie

DIRECTIONS

When to Apply

TURE MAGIC LAWFE FOOD containing chlorders is effective enytime during the year, however best results can be expected when it is applied in early spring. This will prevent domage from grubs during summer mentles.

How Much to Appl

Apply of the rate of 2π per 100 sq. ft. Contents of this bag covers 2500 sq. ft. or on ero 30 s 30 ft.

How to Appl

To obtain best results TURF MAGIC LAWN FOOD certaining chlordene should be evenly applied with no slops or misses. It is recommended that the area be covered trice. Apply one-half the application, then apply the remaining half at right engles to the first to source therough coverage.

Spreader Settings	
Terl Magic	
Control	
Soon	 4.4
Cpclane	\$4

TAYLOR EVANS SEED CO. — TULIA, TEXAS 79088

bolowed law food well and and welled.



ACCEPTED

NOV 1 3 1972

INDIPATION AND DESCRIPTION OF ACTION OF ACTION

Lami Follo

contonuit 34/15/16/2011

DO NOT REUSE EMPTY CONTAINER.
DESTROY IT BY BURYING WITH WASTE
OR BURNING. STAY AWAY FROM SMOKE
OR FUMES.

E.P.A. Registration No. 8952.4

. W. T. A. C. S.


	CAUTIN
\$1,450 CM MAC COMPANY OF WAR \$1,450 CM MAC COMPANY OF WAR \$1,550 CM COMPANY OF WAR \$1,500 CM CM COMPANY OF WAR \$1,500 CM	THE STATE OF THE S
THIS PRODUCT IS TORIC TO FISH AND A LOU ON TREATE! AFEAS MAY BE A LIED MEEP DU BATER AFFLY I'M SECOND ONLY AS SPEL F DO NOT AFFLY MHERE RUNCHES SUMELY! CONTAIN MAYE MATER BY CLEANING OF EQUIP OF MAGNES.	(10.00 N
ACTIVE INCREDIENTS Technical Chiorages	351
MERT INCHEDENTS	99 55 V
*Epispert to 21% Octor is 8 ** restaucts after conduct and -1	n bou sown


50 LBS. NET WT.

TURF MAGIC LAWN FOOD 12-4-4 centaining chlordane

is a highly effective combination of chamical and natural fartilizer with insecticide. When applied as recommended, this product will feed your grass & central white grabs and the following and self-inserts.

Relatic Garden Bootle (Larves)





GUARANTEED ANALYSIS

Tatal Nitrogen	(N) 12.00%
Ammeniace	d Nitrogen 11.5%
Water inco	hèle Nitregen
Available Phas	pheric Acid (P ₁ O ₁)
Soluble Putesh	(KO) 4.00%
Primary Plant Feet	ls derived from:
Sulphate of A	ramonia, Diammonium Phoephote, Sulfate of Petash Magnesia, Tung Moel, one Mool, and Cottonwood Mool.
	2.50%
Sulfur (combin	ed)
	1.00%
	3
Sulphate of F tran Cholate. manium Sulfat	- -
Natural Plant Fe	ed derived from: Bone Meal, Blood Meal, Cottonseed Meal & Tung Meal.
ACTIVE INGREDIE	
Technical Chlo	rdone*
INERT INGREDIEN	TS
TOTAL	

DIRECTIONS

When to Apply

TURF MAGIC LAWN FOOD containing chlordane is effective anytime during the year, however best results can be expected when it is applied in early spring. This will prevent damage from grubs during summer months.

*Equivalent to .21% Octachlare-4, 7-methanatetrahydroindane and .14% Related Compounds.

How Much to Apply

Apply of the rate of 2= per 100 sq. ft. Contents of this bog covers 2500 sq. ft. or on orea

ACCEPTED

NOV131972

FUNGICISE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER-FD UNDER NO.8952-1 SUBJECT TO ATTACHED COMMENTS.



							• • • • • •			1.00%
Moreonies		• • • • •	• • • • •						• • • • •	
20cc	• • • • • • •		. 	.	• • • • •		• • • • • •	• • • • • •	• • • •	.88%
Secondary Plant Fo Sulphate of Po	eds & Tre	co Blor	nonts	derived	frem:		`1			
Sulphoto of Po tran Cholato. S monium Sulfato.	ultur and	nesia, Magn	iren : stium	Sulphata derived	, Mang from	anoso Sulfat	of Pote	, Zinc Su sah Magu	nosia (nd Am
Natural Flori Foo	d derived	from:	Bone	Meal,	Blood	Moal,	Cattonee	ed Meal	& Tun	g Mod
Natural Flant Fact ACTIVE INGREDIEN Technical Chlori	ats.									
ACTIVE INGREDIEN	rrs deno*					. 	• • • • • •		* • • • •	
ACTIVE INGREDIENT	175 Jano* S					. 				

DIRECTIONS

When to Apply

TURF MAGIC LAWN FOOD containing chlordene is effective enytime during the year, however bost results can be expected when it is applied in early spring. This will prevent demage from grubs during summer months.

New Much to Apply

Apply at the rate of $2 = por^2 100$ sq. ft. Contents of this bag covers 2500 sq. ft. or on order 50 = 50 ft.

How to Apply

To obtain best results TURF MAGIC LAWN FOOD containing chlordane should be evenly applied with no skips or misses. It is recommended that the area be covered twice. Apply ane-half the application, then apply the remaining half at right angles to the first to assure thorough coverage. Water thoroughly. Avaid run-aff. Birds feeding on treated areas may be harmed.

proader S	ettic	Y	}												
Turf Ma	gic			. .											8
Central	٠.				 		 				•	•			•
Scott			. <i>.</i>		 		 •							-	9-1/2
Cyclone															

TAYLOR EVANS SEED CO. — TULIA, TEXAS 50 LBS. NET WT. 79088

NOV131972

UNDER THE ELDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER-ED UNDER NO. \$252-1 SUBJECT TO ATTACHED COMMENTS.



DO NOT REUSE EMPTY CONT DESTROY IT BY BURYING WITH OR BURNING. STAY AWAY FROM OR FUMES.

C.P. A. Regulated

FOR ECONOMIC POISON REGISTER-ED UNDER NO. \$952-1 SUPJECT TO ATTACHED COMMENTS.

Lawn Food

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

E.P. A. Registration No. 8952-4

centerative chevilence

CAUTION

PECH C REPORT THE PLAN RETURNED OF CONTROL CON

THIS PRODUCT IS TOXIC TO FISH AND MILDLIFE BIFDS FEEDING ON TREATER APEAS MAY BE KILLED MEEPING OF ANY BODY OF WATER APPLY THIS PRODUCT ONLY AS SPECIFED ON THIS LASEL DO NOT APPLY WHERE RUNOFF IS LIKELY TO DOCUR DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES.

ACTIVE INGREDIENTS:

Technical Chlorophe*

Xs.

MERT INCREDIENTS

99 65%

100.000

IVIA.

W 27

"Equipment to 21% Octobrishs 4 7 methodologically branching and 14% floated Compounds

MACROLIA WEEDS PLUS FEEDS FERTILIZER 10-5-5

plus catalysts

U.S.D.A. Reg. No. 9123-2

Active Ingredients: Technical chlordane*. Silvex (2- (2, 4, 5-trichlorophenoxy) propionic soid 0.15 % *Remivalent to 0.36 % Chlordane and 0.24 % related compounds CAUTION: KEEP OUT OF THE REACH OF CHILDREN. See back panel for additional cautions. Mfg. by Magnolia Fertiliser Co., Seattle, Wa. Net wt. 40 lbs. back Catalyst A-10 makes fertilizer work Catalyst C makes weed killer more better, faster. Helps grass roots effective. Helps weed killer to cover, adhere to pick up fertilizer more easily for and penetrate the weed leaf for botter quicker greening. Helps prevent forweed destruction. tiliser runoff that starve a lawns high Woods plus Foods will not burn when spots, & overgeeds low spots; for more applied according to directions. even greening and growth. DIRECTIONS: Apply when weeds are young and actively growing. 1. Before application sprinkle lawn using a fine misty spray. 2. Apply Woods plus Foods evenly with fertilizer spreader. See spreader chart below. Ampid overlapping. ACCE To be party atter for 24 hours after application unless temperature remains ever 70 F. Watering should then be done 12 hours after application. Water the Watering should then be done 12 hours after application. Water thereughly. MAF-7-1468 Amount to apply AL HEFCTICIDE UNDER 3: spreader settings SENSONE ACT FUES CON

CAUTION: Avoid contact with skin, eyes & clothing. Avoid prolonged inhalation of dust. Avoid contamination of feed & foodstuffs. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not store near seeds, fertilizers, insecticides or fungicides. Avoid contact with desirable plants & with bentgrass, clover and dichondra, & with grass seedlings on newly seeded launs as possible injury may result. Avoid dust drifting to & application to shrubs & crnementals. Wash hands thoroughly after use. Do not contaminate ornamental fish pends, demestic or irrigation water.

Modern

4

Do not reuse container - destroy when empty. Clean & wash fertilizer spreader thoroughly after use.

Controls these weeds:	plantain buttercup pig veed	docks clovers sorrel	chickweed dandelion mustard	lambaquarter sow thistle

Plant food analysis:

POINCE HEGISTERED

40 1be WEEDS - MEEDS

7 per 5000 sq. ft.

FOR LC CHOM

917

DHOFK IN

Total Nitrogen (N) 10 % Available Phosphoric acid (P₂O₂) . 5 % Soluble Potash (K₂O) 5 % Magnesium (Mg) from dolomite . . 2 %

Ingredients: Urea Formaldehyde, tankage, ammonium phosphate, urea, ammonium sulfate, muriate potash, manures, calcium 2 magnesium carbonates, vermipulite, and herbicides & insecticides as shown em front.

Lawn Beauty

5

Scott

CAUTION: COMBUSTIBLE MIXTURE

Long range jet spray kills wasps, hornets, bees, yellow jackets. Shoots a lethal jet spray into nests, crevices, holes, etc. from 12 feet or more.

DIRECTIONS FOR USE: Remove protective cap. Point nozzle towards nest, crevice or hole where infestation or insects are known to be inside. Press valve all the way down until area is saturated with this insecticide. Also spray around this area until completely wet with blitz jet spray. This treated surface will have a residual killing action and prevent reinfestation. In the event the insecticide film becomes ineffective due to rain or other causes, repeat treatment. For outdoor use only.

CAUTION: Keep out of reach of children. Do not apply on persons. In case of skin contact wash with soap and water. Repeated or prolonged contact with skin can cause toxic symptoms. Harmful is swallowed.

Do not spray on pets or other warm blooded animals. Do not apply on vegetation. Do not spray on asphalt tile. Contents of this container are under pressure. Do not puncture container or dispose of same in fire. Container should not be stored in sunlight or where temperatures exceed 120 F. Do not use near fire or open flame.

ACTIVE INGREDIENTS: Pyrenthrins 0.15% technical piperonyl butoxide *0.30%, n-octyl bicycloheptene Dicarboximide 0.53%, technical chlordane **2.00%, petroleum distillate 62.02%.


INERT INGREDIENTS: 35.00%

*Equivalent to 0.24% (butylcarbityl) (6-propylpiperonyl) ether and 0.6% other related compounds.

**Equivalent to 1.2% octachloro-4, 7-methanotetrahy-droindane and 0.8% of related compounds.

U.S.D.A, Reg. No. 9143-7

manufactured by
CHEMSCOPE CORPORATION
155 Howell Street
Dallas, Texas



Wasp and Hornet Long Range Jet Spray

KILLS . . . HORNETS • YELLOW JACKETS WASPS • BEES

Shoots A Lethal Jet Into Nests, Crevices, Holes, etc., From 12 Feet Or More Away

Net Weight 1 pound 2 %.

CAUTION:

Keep out of reach of children. Contents under pressure. See other cautions on back panel.