UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C. 20460

PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

October 25, 2005

Marianne Radtke Regulatory Affairs Manager Applied Biochemists W175 N11163 Stonewood Drive Suite 234 Germantown, WI 53022

Subject:

Cutrine Plus

EPA Registration No. 8959-10

Application Date: September 17, 2005 EPA Received Date: September 27, 2005

Dear Ms. Radtke:

This acknowledges receipt of your notification, submitted under the provision of PR Notice 98-10, FIFRA section 3(c)9.

Proposed Notification

- Alternate Brand Name "Cutrine Plus Algaecide"
- Revised Directions for Use

General Comments

Based on a review of the material submitted, the following comments apply:

The notification application is acceptable. A copy has been inserted in your file for future reference.

Should you have any questions or comments concerning this letter, please contact me at 703-308-6422.

Sincerely.

Adam Heyward

Product Manager 34

Regulatory Management Branch II Antimicrobials Division (7510 C)

	•	
	CONCURRENCES	
YMBOL) 75 /0C		
JRNAME Day		
ATE \$ 1,-75.55		
² PA Form 1320-1A (1/90)	Printed on Recycled Pages	OFFICIAL EN E CODY

United States Environmental Protection Agency Washington, DC 20460 Registral Amends Amends Control Control					tion	OPP Identifier Number		
	Applicat	<u>ion for Pesticide -</u>	Section	1				
1. Company/Product Number	2. EPA Produ	_		3. Pro	oposed Classification			
8959-10	Adam Hey	/ward		[None Restricted			
4. Company/Product (Name) Cutrine Plus	PM# 34							
5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3)								
Applied Biochemists		1	(b)(i), my product is similar or identical in composition and labeling to:					
W175 N11163 Stonewood Dr. S	te 234	EPA Reg. (No					
Germantown, WI 53022								
Gheck if this is a new addr	9S\$	Product Na	ame					
		Section - II						
Amendment - Explain below. Resubmission in response to Agenc	v letter dated	Ager	printed labe ncy letter dat Too* Applica		to			
✓ Notification - Explain below.	, , , , , , , , , , , , , , , , , , , ,		ır - Explain be					
Explanation: Use additional page(s) if ne	seesen. (For secti	on Land Section II)						
This is a notification of an additional brand no attached label. This notification is consisten made to the labeling or the confidential state statement to EPA. I further understand that it violation of FIFRA and I may be subject to er	t with the provisions ment of formula of thi if this notification is no	of PR Notice 98-10 and EPA is product. I understand that of consistent with the terms of	A regulations a it is a violatio of PR Notice	at 40 CFR 152 in of 18 U.S.C 98-10 and 40	2.46, and n . Sec. 1001	o other changes have been I to willfully make any false		
		Section - III						
1. Material This Product Will Be Packaged I	ភ:			1	·			
Child-Resistant Packaging Unit Packagin	ng	Water Soluble Package	ing	2. Type of	-			
Yes		Yes	✓ Plastic					
✓ No ✓ No		V No	Glass					
* Cartification must Unit Packagin	No. per ng wgt. container		o. per entainer		Paper Other (S	pecify)		
3. Location of Net Contents Information	etail Container	5. Lo	cation of Lab	el Directio	ua			
Label Container 1, 2-1/2, 5, 25 gallon								
5. Manner in Which Label is Affixed to Product Screenprinted Lithograph Paper glued Stenciled								
	Otton.	Section - IV						
1. Contact Point (Complete items directly L	selow for identificati	· · · · · · · · · · · · · · · · · · ·	acted if nec	essan to on	ress this	ennlication l		
Name		Title	20100,					
Marianne Radtke	Regulatory Affairs Mar	tory Affairs Manager Telephone No. (Include Area Code) 262 2565 4449 ext 328						
I certify that the statements I have m I acknowledge that any knowlinglly fa both under applicable law.		d all attachments thereto a	-		•	6. Late Application Received (Stamped)		
2. Signature	3. Title							
Regulatory Affairs Manager								
4. Typed Name		5. Date						
Marianne Radtke	Sept. 17	⁷ , 2005						

CUTRINE - PLUS

ALGAECIDE

Pat. No. 3.930.834

EPA Reg. No. 8959-10

EPA Est. No. 42291-GA-1

FOR USE IN LAKES - POTABLE WATER RESERVOIRS FARMS, FISH AND INDUSTRIAL PONDS, FISH HATCHERIES AND RACEWAYS, CROP AND NON-CROP IRRIGATION CONVEYANCE SYSTEMS, DITCHES, CANALS AND LATERALS

ACTIVE INGREDIENTS:	*n.00
COPPER AS ELEMENTAL	9.0%
INERT INGREDIENTS:	91.0%
TOTAL	100.0%

CUTRINE-PLUS contains 0.909 lbs. of elemental copper per gallon. *From mixed Copper-Ethanolamine complexes

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID				
If inhaled:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to- mouth if possible. Call a poison control center or doctor for further treatment advice. 			
lf on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. 			
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 			
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. 			

See Additional Precautions on Back Panel

____GALLONS

MANUFACTURED BY:



GERMANTOWN, WISCONSIN 53022 1-800-558-5106

GENERAL INFORMATION



CUTRINE-PLUS, under field conditions, is effective in controlling a broad range of algae including: Chara, Spirogyra, Cladophora, Vaucheria, Ulothrix, Microcystis and Oscillatoria. CUTRINE-PLUS has also been proven effective in controlling the rooted aquatic plant. The ethanolamines in CUTRINE-PLUS prevent the precipitation of copper with carbonates and bicarbonates in the water. Waters treated with CUTRINE-PLUS may be used for swimming, fishing, drinking, livestock watering or imigating turf, ornamental plants or crops immediately after treatment.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SURFACE SPRAY/INJECTION

ALGAECIDE APPLICATION

For effective control, proper chemical concentration should be maintained for a minimum of three hours contact time. The application rates in the chart are based on static or minimal flow situations. Where significant dilution or loss of water from unregulated inflows or outflows occur (raceways) within a three hour period, chemical may have to be metered in.

- Identify the algae growth present as one of the following types: Planktonic (suspended),
 Filamentous (mat forming), or Chara/Nitella.
- Determine the surface acreage (1 acre=43,560 sq. ft.) and average depth of infested area.
- Refer to the chart below to determine gallons of CUTRINE-PLUS to apply per surface acre.

Application Rates Gallons Per Surface Acre

ALGAE TYPE	PPM COPPER	DEPTH IN FEET					
ALGAE TIPE	GOTTER	1	2	3	4		
Planktonic	0.2 - 0.6	0.6 – 1.8	1.2 – 3.6	1.8 - 5.4	2.4 – 7.2		
Filamentous	0.2 - 0.8	0.6 - 2.4	1.2 – 4.8	1.8 - 7.2	2.4 - 9.6		
Chara/Nitella	0.4 - 1.0	1.2 – 3.0	2.4 – 6.0	3.6 - 9.0	4.8 - 12.0		

- Under conditions of heavy infestation, treat only 1/3 to 1/2 of the water body at a time to avoid fish suffocation caused by oxygen depletion from decaying algae.
- Before applying, dilute the required amount of CUTRINE-PLUS with enough water to ensure even distribution with the type of equipment being used (e.g. 9 parts water to 1 part Cutrine-Plus used in a hand-pump sprayer). Faster results will be seen when applied under calm and sunny conditions when water temperature is a least 60°F. Break up floating algae mats before spraying or while application is being made. Use hand or power sprayer adjusted to rain-sized droplets. Spray shoreline areas first to avoid trapping fish.

CUTRINE-PLUS Granular Algaecide may be used as an alternative in low volume flow situations, spot treatments or treatment of bottom-growing algae in deep water.

HERBICIDE APPLICATION (For Hydrilla Control) CUTRINE-PLUS:

Control of Hydrilla verticillata can be obtained from copper concentrations of 0.4 to 1.0 ppm resulting from CUTRINE-PLUS treatment. Choose the application rate based upon stage and density of Hydrilla growth and respective water depth from the chart below.

Application Rates
Gallons/Surface Acre*

Growth/Stage Relative Density	PPM Copper	DEPTH IN FEET					
		1	2	3	4	5	6
Early Season Low Density	0.4 0.5	1.2 1.5	2.4 3.0	3.6 4.5	4.8 6.0	6.0 7.5	7.2 9.0
***	0.6	1,8	3.6	5.4	7.2	9.0	10.8
Mid-Season Moderate Density	0.7	2.1	4.2	6.3	8.4	10.5	12.6
	0.8	2.4	4.8	7.3	9.6	12.0	14.4
Late Season High Density	0.9 1.0	2.7 3.0	5.4 6.0	8.1 9.0	10.8 12.0	13.5 15.0	16.2 18.0

^{*} Application rates for depths greater than six feet may be obtained by adding the rates given for the appropriate combination of depths.

Application rates should not result in excess of 1.0 ppm copper concentration within treated water.

CUTRINE®PLUS: REWARD®TANK MIX

On waters where enforcement of use restrictions for recreational, domestic and irrigation uses are acceptable, the following mixture can be used as an alternative Hydrilla control method.

Tank mix 3 gallons of CUTRINE-PLUS with 2 gallons of REWARD[®]. Apply mixture at the rate of 5½ gallons per surface acre. Dilute with at least 9 parts water and apply as a surface spray or underwater injection. Observe all cautions and restrictions on the labels of both products used in this mixture.

PERMITS:

Some states may require permits for the application of this product to public waters. Check with your local authorities.

DRIP SYSTEM APPLICATION

FOR USE IN POTABLE WATER AND IRRIGATION CONVEYANCE SYSTEMS

- CUTRINE PLUS should be applied as soon as algae or Hydrilla begins to interfere noticeably with normal delivery of water (clogging of lateral headgates, suction screens, weed screens and siphon tubes). Delaying treatment could perpetuate the problem causing massing and compacting of plants. Heavy infestations and tow flow conditions increasing water flow rate during application may be necessary.
- Prior to treatment it is important to accurately determine water flow rates. In the absence of weirs, orifices, or similar devices which give accurate water flow measurements, volume of flow may be estimated by the following formula:

Average Width (feet) x Average Depth (feet) x Velocity* (feet/second) x 0.9 = Cubic Feed per Second (C.F.S.)

*Velocity is the time it takes a floating object to travel a given distance. Dividing the distance traveled (feet) by the time (seconds) will yield velocity (feet/second). This measurement should be repeated at least three times at the intended application site and then averaged.

 After accurately determining the water flow rate in C.F.S. or gallons/minute, find the corresponding CUTRINE-PLUS drip rate on the chart below.

WAT FLOW I			DRIP RATE*	3
C.F.S.	Gal/Min	Qts/Hr.	MI/Min.	FL.Oz/Min.
1	450	1	16	0.5
2	900	2	32	1.1
3	1350	3	47	1.6
4	1800	4	63	2.1
5	2250	5	79	2.7

^{*}REWARD* is a trademark of Syngenta Crop Protection, Inc.

- Calculate the amount of CUTRINE-PLUS needed to maintain the drip rate for a period of 3 hours by multiplying Qts./Hr. x 3; ml/Min. x 180; or Fl. Oz./Min. x 180. Dosage will maintain 1.0 ppm Copper concentration in the treated water for the 3 hour period. Introduction of the chemical should be made in the channel at weirs or other turbulence-creating structures to promote the dispersion of chemical.
- Pour the required amount of CUTRINE-PLUS into a drum or tank equipped with a brass needle valve and
 constructed to maintain a constant drip rate. Use a stop watch and appropriate measuring container to set the
 desired drip rate. Readiust accordingly if flow rate changes during the 3 hour treatment period.
- Distance of control obtained down the waterway will vary depending upon density of vegetation growth. Periodic maintenance treatments may be required to maintain seasonal control.

GENERAL TREATMENT NOTES

The following suggestions apply to the use of **CUTRINE-PLUS** as an algaecide or herbicide in all approved use sites: **For optimum effectiveness...**

- · Apply early in the day under calm, sunny conditions when water temperatures are at least 60°F.
- Treat when growth first begins to appear or create a nuisance, if possible.
- Apply in a manner that will ensure even distribution of the chemical within the treatment area.
- Re-treat areas if regrowth begins to appear and seasonal control is desired. Allow one to two weeks between consecutive treatments.
- Allow seven to ten days to observe the effects of treatment (bleaching and breaking apart of plant material).

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

CORROSIVE. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling this product. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep container closed when not in use. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Reseal container and offer for recondition or triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Consult Federal, State or local authorities for approved alternative procedures.

ENVIRONMENTAL HAZARDS:

This product may be toxic to trout and other species of fish. Fish toxicity is dependent upon the hardness of water. **Do not** use in water containing trout if the carbonate hardness of water does not exceed 50 ppm. **Do not** use in waters containing Koi and hybrid goldfish. This product is not intended for use in small volume, garden pond systems.

NOTICE

Neither the manufacturer nor the seller makes any warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes risk of use of this material when such use is contrary to label instructions. Read and follow the label directions carefully.

@ Applied Biochemists 2002 091405

APPLIED BIOCHEMISTS

W175 N11163 Stonewood Drive Suite 234
Regulatory Services
Germantown, WI 53022-4799
262 255-4619 Fax
262 255-4449 ext. 328

E-mail -mradtke@advantistech.com

September 19, 2005

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7510C)
U.S. Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Avenue, N.W.
Washington, D.C. 20460-0001

Re: EPA Reg. No. 8959-10

Cutrine Plus Algaecide

Attached is an application for notification of an alternate brand name and additional of a statement to clarify mixing directions. As stated on the form no other changes have been made to any part of the product labeling.

Enclosed you will find

Form 8570-1 for Notification 1 label copy with change highlighted

Sincerely,

(

APPLIED BIOCHEMISTS

Marianne Radtke

Regulatory Affairs Manager