



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

8959-56

Date of Issuance:

FEB 24 2010

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional
~~Conditional~~ Expires 2/28/2015

Name of Pesticide Product:

NAVITROL

Name and Address of Registrant (include ZIP Code):

APPLIED BIOCHEMISTS
W175 N11163 Stonewood Drive Suite 234
Germantown, WI 53022

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for registration review of your product under FIFRA section 4.

2. Make the following label changes before you release the product for shipment:

a. Revise the EPA Registration Number to read "EPA Reg. No. 8959-56".

b. Add the batch number to label on all non-refillable containers.

c. The 100% repack product you reference does not have a label for aquatic uses. Therefore, remove all references to aquatic uses:

- On page 1, under the label name, NAVITROL, remove "and Aquatic Herbicide". In the first paragraph under label name, change to "FOR CONTROL OF SELECTIVE WOODY PLANTS,

VINES, AND ANNUAL AND PERENNIAL BROADLEAF WEEDS IN GRASS PASTURES AND RANGELAND, FORESTS, AND NON-CROP AREAS WHICH ARE INDUSTRIAL, MANUFACTURING, AND STORAGE SITES, RIGHTS-OF-WAY WHICH ARE ELECTRICAL POWER LINES, COMMUNICATION LINES, PIPELINES, ROADSIDES, RAILROAD BEDS, AND NON-IRRIGATION DITCH BANKS, AND FENCE ROWS NOT ADJACENT TO CROP FIELDS EXCEPT GRASS PASTURES AND RANGELAND, AND AROUND FARM BUILDINGS."

- On page 3, under "ENVIRONMENTAL HAZARDS", remove paragraph starting with "For aquatic uses,". Remove "For terrestrial uses" from beginning of second paragraph, and change "do not apply directly to water, etc." to "Do not apply directly to bodies of water, etc." to the beginning of second paragraph.

- On page 4, under "GENERAL INFORMATION", remove "aquatic plants" from first sentence, and remove the sixth bullet starting with "In aquatic sites such as ponds, etc."

- On page 5, change "GENERAL USE PRECAUTIONS" to "USE RESTRICTIONS AND PRECAUTIONS" and change any reference to "GENERAL USE PRECAUTIONS" on label to "USE RESTRICTIONS AND PRECAUTIONS section". Remove the fourth bullet starting with "It is permissible to treat non-irrigation ditch banks, etc.", because it is redundant, and there are potential aquatic uses included in this bullet. Change the 7th bullet to "Do not apply through a mist blower."

- Remove all content on pages 16 to 24. These are wetland and direct application use directions pertaining to water bodies, which are not approved for this label.

3. On page 3, change the bullet "Protective eyewear" to "Protective eyewear with side-shields or full face shield."

4. On page 12, under "FOLIAGE TREATMENT (RIGHTS-OF-WAY)", correct typo "2,4-d" to "2,4-D".

5. On pages 25 and 26, in the "STORAGE AND DISPOSAL" section, change the third to the last sentence in the >5 gallon, >5 gallon, and Refillable paragraphs to "Then offer for recycling or reconditioning if available, or if allowed by state and local authorities, puncture and dispose of in an approved landfill or by burning."

6. On page 26, under "WARRANTY DISCLAIMER" statement, to the next to the last sentence of paragraph, add "TO THE EXTENT PERMITTED BY LAW" before "...ASSUMED BY BUYER OR USER."

7. On page 26, under "LIMITATION OF LIABILITY" statement, add "TO THE EXTENT PERMITTED BY LAW" to the beginning of the second sentence starting with "MANUFACTURER OR SELLER, etc.", and to the beginning of the last sentence.

8. Submit one copy of the final printed label for the record before you release the product for shipment.


If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

For additional assistance with this matter, please contact Phil Errico at 703-305-6663/
errico.philip@epa.gov .

A stamped copy of the label with comments is enclosed for your records.

Enclosure: Accepted label with comments for product with EPA Reg. No. 8959-56.

Signature of Approving Official:



James A. Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505P)

Date:

FEB 24 2010

EPA Form 8570-6

NAVITROL

Landscape and Aquatic Herbicide

EPA Reg. No. 8959-XX

EPA Est. No. 42291-GA-1

FOR CONTROL OF WOODY PLANTS, AQUATIC PLANTS, VINES; AND ANNUAL AND PERENNIAL BROADLEAF WEEDS IN FORESTS, NON-CROP AREAS INCLUDING INDUSTRIAL MANUFACTURING AND STORAGE SITES, RIGHTS-OF-WAY SUCH AS ELECTRICAL POWER LINES, COMMUNICATION LINES, PIPELINES, ROADSIDES, AND RAILROADS, FENCE ROWS, IN NON-IRRIGATION DITCH BANKS, AQUATIC SITES SUCH AS PONDS, LAKES, RESERVIORS, NON-IRRIGATION CANALS, AND DITCHES WHICH HAVE LITTLE OR NO CONTINUOUS OUTFLOW AND AROUND FARM BUILDINGS.

Active Ingredients:

*Triclopyr: (3,5,6-trichloro-2-pyridinyl)Oxyacetic acid, triethylamine salt ...44.4%
OTHER INGREDIENTS:55.6%
TOTAL.....100.0%

*Contains 3 pounds of triclopyr acid equivalent per gallon (31.8%)

KEEP OUT OF REACH OF CHILDREN DANGER – PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.
For Medical Emergencies Only,

See additional precautions on Back Panel

NET CONTENTS: _____ Gallons (_____ Liters)

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

FEB 24 2010

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

8959-56

MANUFACTURED FOR:



W175 N11163 Stonewood Drive Suite 234
 GERMANTOWN, WISCONSIN 53022
 1-800-558-5106
www.appliedbiochemists.com

FIRST AID	
IF IN EYES	* Hold eye open and rise slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. * Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	* Take off contaminated clothing. * Rinse skin immediately with plenty of water for 25 to 20 minutes. * Call a poison control center or doctor for treatment advise.
IF SWALLOWED	* Call a poison control center or doctor immediately for treatment advise. * Have person sip a glass of water if able to swallow. * Do not induce vomiting unless told to do so by the poison control center or doctor. * Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. If a medical emergency arises contact Arch Chemicals Emergency Action Network in the US call 1-800-654-6911 or outside the US call 423-780-2970. For help with a spill, leak, fire or exposure involving this material call CHEMTREC 1-800-424-9300.	
NOTE TO APPLICATOR Allergic skin reaction is not expected from exposure to spray solution of this product when used as directed.	
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage.	

**PRECAUTIONARY STATEMENTS
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 DANGER- PELIGRO**

Corrosive. Causes Irreversible Eye Damage. Harmful if absorbed through skin or swallowed. Do not get in eyes or on skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers who handle this pesticides must wear:

- Long-sleeved shirt and long pants

- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves (>14mils) such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of glove before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For aquatic uses, under certain conditions, treatment of aquatic weeds can result in oxygen depletion or loss to decomposition of dead plants, which may contribute to fish suffocation. This loss can cause fish suffocation. Therefore, to minimize this hazard, do not treat more than one-third to one-half of the water area in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow fish to move into untreated areas. Consult with the State agency for fish and game before applying to public water to determine if a permit is needed.

For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE. Do not use or store the product near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves (>14mils) such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber.
- Shoes plus socks
- Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For applications to non-cropland area, do not enter or allow others to enter the treated area until sprays have dried.

GENERAL INFORMATION

This product is used to control unwanted woody plants; aquatic plants; and annual perennial broadleaf weeds

- In forest

- In non-crop area including industrial manufacturing and storage sites
- In rights-of-way such as electrical power lines, communication lines, pipelines, roadsides, and railroads
- In fence rows
- In non-irrigation ditch banks
- In aquatic sites such as ponds, lakes, reservoirs, non-irrigation canals, marshes, wetlands and ditches which have little or no continuous outflow.
- Around farm buildings

This product's use on these sites may include application to grazes areas as well as for the establishment and maintenance of wildlife openings.

GENERAL USE PRECAUTIONS

- Do not apply this product through any type of irrigation system.
- Do not apply to ditches or canals used to transport irrigation water. It is permissible to treat non-irrigation ditch banks.
- Do not apply where runoff or irrigation water may flow onto agricultural fields as injury to crops may result.
- It is permissible to treat non—irrigation ditch banks, seasonally dry wetlands (such as flood plains, Deltas, marshes, swamps, or bogs) and transitional areas between upland and lowland sites.
- Do not apply directly to un-impounded rivers or streams. Do not apply to salt water bays or estuaries.
- When making application to banks or shorelines of moving water sites, minimize the overspray to open water.
- Application through a mist blower is not recommended.
- Do not make direct applications or allow spray mists to drift onto cotton; grapes; soybeans; tobacco; vegetable crops; flowers; ornamental shrubs or trees; or other desirable broadleaf plants.
- For range and pasture sites, including rights-of-way, fence rows, or any area where grazing and harvesting is allowed, do not apply more than 2 pounds a.e. of triclopyr (2/3 gallon of this product) per acre per year.
- For forestry uses, do not apply more than 6 pounds a.e. triclopyr (2 gallons of this product) per acre per year.
- For all terrestrial uses other than rangeland, pasture, forestry sites, and grazes area, a maximum of 9 pounds a.e. of triclopyr (3 gallons of this product) per acre per year may be applied.
- All livestock, except lactating dairy animals, can graze at any time.
- Lactating dairy animals cannot graze forage until the next growing season after application.
- For all livestock, wait 14 days after application before harvesting hay.
- Grazes areas of non-cropland and forestry sites may be spot treated if they comprise no more than 10% of the total grazable area.
- Withdraw livestock from grazing treated grass or consumption of treated hay at least 3 days before slaughter.

- Arizona: This product has not been approved for use on plants grown for commercial production, specifically forests grown for commercial timber production, or on designated grazing areas.
- Obtain required permits: Consult with appropriate state or local water authorities before applying this product to public waters. State or local public agencies may require permits.

APPLICATION DIRECTIONS

Rates

This table assists in determining proper volumes of this product in the spray tank to avoid exceeding the maximum use rates using carrying spray volumes.

Maximum Rate of This Product			
Spray Volume (gal/acre)	Rangeland and Pasture Sites ¹ (gal/100gal of spray)	Forestry Sites ² (gal/100 gal of spray)	Other Non-Cropland Sites ³ (gal/100 gal of spray)
400	Do not use	0.5	0.75
300	Do not use	0.67	1
200	Do not use	1	1.5
100	0.67	2	3
50	1.33	4	6
40	1.67	5	7.5
30	2.33	6.65	10
20	3.33	10	15
10	6.67	20	30

- ¹ For range and pasture sites, including rights-of ways, fence rows, or any area where grazing and harvesting is allowed, do not apply more than 2 lb a.e. of triclopyr (2/3 gallon of this product) per acre per year.
- ² For forestry uses, do not apply more than 6lb a.e. triclopyr (2 gallons of this product) per acre per year.
- ³ For all terrestrial uses other than rangeland, pasture, forestry sites, and grazes area, a maximum of 9lb a.e of triclopyr (3 gallons of Triclopyr) per acre per year.
- ⁴

SPRAY ADDITIVES

All surfactants and drift control agents must be approved for food or feed use when used on food or feed sites.

Surfactants- When using surfactants, follow the use directions and precautions listed on the surfactant manufacturer's label. Use the higher recommended concentration of surfactant in the spray mixture when applying lower sprayer volumes per acre.

Drift Control Agents- Agriculturally approved spray thickening drift control agents or high viscosity invert systems may be used with this product. When using these agents, follow all use directions and precautions on the product label. Do not use a thickening agent with the Microfoil boom, Thru-Valve boom or other systems that cannot accommodate thick sprays.

TANK MIXES

Always refer to labels of other pesticide products for mixing directions and precautions which may differ from those outlined here. Use in accordance with most restrictive of label limitations and precautions. No label dosage rates may be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

Tank Mixing Recommendations:

- Fill spray tank $\frac{1}{2}$ full with water.
- Add spray thickening agent (if used).
- Add additional herbicide (if used).
- Add this product.
- Add surfactant (if used).
- Fill remainder of spray tank.

If combined with emulsifiable concentrate herbicides, moderate continuous adequate agitation is required.

SPRAY DRIFT MANAGEMENT

Avoid injurious drift. Application should only be made when there is little or no hazard from spray drift. Very small quantities of spray may seriously injure susceptible plants.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all of these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications.

- The distance of the outer most operating nozzles on the boom must not exceed $\frac{3}{4}$ the length of the rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed.

Aerial Drift Reduction Advisory

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity and Temperature Inversions).

Controlling Droplet Size:

- **Volume-** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure-** Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles-** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation-** Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type-** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between winds speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence

wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas:

The pesticide should only be applied when potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

APPLICATION EQUIPMENT AND TECHNIQUES

Broadcast Applications

Aerial Application – When making aerial applications on right-of-way or other areas near susceptible crops, apply through a Microfoil (1) or the Thru-valve(1) boom or use an agriculturally approved drift control agent. Other drift reducing systems or thickened sprays prepared by using high viscosity inverting systems may be used if they are made as drift-free as are mixtures containing agriculturally approved thickening agents or applications made with the Microfoil or Thru-Valve boom. Keep spray pressures low enough to provide coarse spray droplets. Spray boom should be no longer than $\frac{3}{4}$ of the rotor length. Spray only when the wind velocity is low (follow state regulations). Avoid application during air inversions. If a spray thickening agent is used, follow all use recommendations and precautions on the product label.

- (1) Note: Reference within this label to equipment produced by or available from other parties is provided without consideration for use by the reader at its discretion and subject to the reader's independent circumstances, evaluation and expertise. Such reference by Applied Biochemists is not

intended as an endorsement of such equipment, shall not constitute a warranty (express or implied) of such equipment, and is not intended to imply that other equipment is not available and equally suitable. Any discussion of methods of use of such EQUIPMENT does not imply that the reader should use the equipment other than is advised in directions available from the equipment's manufacturer. The reader is responsible for exercising their own judgment and expertise, or consulting with sources other than Applied BioChemists, in selecting and determining how to use its equipment.

Ground Application – To aid in reducing spray drift, this product should be applied in thickened (high viscosity) spray mixtures using an agriculturally approved drift control additive, high viscosity invert system, or equivalent as directed by the manufacturer. Use of low pressure nozzles; and operating nozzles in the lower end of the manufacturer's recommendations is advised. To minimize drift, keep the spray boom as low as possible, apply in > 20 gallons of spray volume per acre, spray when wind velocities are low; or use an approved drift control agent.

In Hand Gun Applications, select the minimum spray pressure that will provide adequate plant coverage (without forming a mist). Do not apply with nozzles that produce a fine droplet spray.

High Volume Leaf-Stem Treatment: To minimize spray drift, do not use pressure exceeding 50PSI as the spray nozzle and keep sprays no higher than brush tops. An agriculturally approved thickening agent may be used to reduce spray drift.

APPROVED USES

Refer to Table 1 and 2 which follow, for a list of wood plants and broadleaf weeds that are controlled by this product.

Apply this product at rates of 0.25 to 3 gallons per acre for the control of broadleaf weeds and woody plants. Apply in enough water to provide uniform and complete coverage of the plants to be controlled. Use only water suitable for spraying. Use of an agriculturally nonionic surfactant is recommended for all foliar applications. For best results make application when woody plants and weeds are actively growing.

Use higher rates within the range when brush averages 15feet or more in height or when brush covers >60% of the area to be treated, Re-sprouting may occur the year following treatment if lower rates are used on hard-to-control species when easy to control brush species dominate, rates below those recommended may be effective. Consult State or local Extension personnel for information.

For hard-to-control species such as ash, black gum, choke cherry, elm, maples, oak, pines or winged elm; during late summer applications when plants are mature; or during drought conditions; use higher rates of this product alone or use in combination with Tordon 101 Mixture. If lower rates are used on hard-to-control species, re-sprouting may occur in the year following treatment. When applying this product in a tank mix with 2,4-D 3.8lb amine, or low volatile ester herbicides, use higher rates of this product for satisfactory brush control.

When tank mixing, refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

FOLIAGE APPLICATIONS WITH GROUND EQUIPMENT

High Volume Foliage Applications

For control of woody plants, apply this product at 1 to 3 gallons per 100 gallons of spray solution. Make applications in 100 to 400 gallons of total spray per acre depending on size and density of woody plants. Coverage should be thorough to wet all leaves, stems, and root collars.

Tank Mixing: 1 to 4 quarts of this product may be tank mixed with 1 to 2 quarts of 2,4-D 3.8 lb amine, or low volatile ester or Tordon 101 Mixture, diluted to make 100 gallons of spray solution. Make applications in 100 to 400 gallons of total spray per acre depending on size and density of woody plants. When tank mixing, refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

Low Volume Foliage Applications

For control of woody plants, mix up to 5 gallons of this product in 10 to 100 gallons of spray solution. Adjust the spray concentration of this product and total spray volume per acre to match the size and density of target woody plants and kinds of spray equipment used. With low volume sprays, use sufficient spray volume to obtain uniform coverage of target plants including the surfaces of all foliage, stems and root collars. For best results, a surfactant should be added to all spray mixtures. See the **Spray Additives** section of **Application Directions**.

Match equipment and delivery rate of spray nozzles to height and density of woody plants. When treating tall, dense brush, a truck mounted spray gun with spray tips that deliver up to 2 gallons per minute at 40 to 60 PSI may be required. Backpack or other types of specialized spray equipment with spray tips that deliver less than 1 gallon of spray per minute may be appropriate for short, low to moderate density brush.

Tank Mixing: Up to 3 gallons of this product may be applied tank mix combinations with 2 to 4 quarts of Tordon K or 1 to 2 gallons Tordon 101 Mixture as a low volume foliar spray. These applications should be made in 10 to 100 gallons of spray solution. When tank mixing, refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved uses, and a list of weeds and woody plants.

BROADCAST APPLICATION WITH GROUND EQUIPMENT

Use equipment that will assure thorough and uniform coverage at spray volumes applied. To improve spray coverage, add an agriculturally approved nonionic surfactant. See the **Spray Additives** sections of **Application Directions**. See **Maximum Application Rates Table** in the **Application Directions** for relationship between mixing rate, spray volume, and maximum application rate.

Woody Plant Control

Foliage Treatment: Apply 2 to 3 gallons of this product in 20 to 100 gallons of spray solution per acre.

Tank Mixing: This product at 2 to 4 quarts per acre may be tank mixed with 1 to 2 gallons of 2,4-D 3.8 lb amine, or low volatile esters or Tordon 101 Mixture, in 20 to 100 gallons of spray solution per acre. When tank mixing, refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

Broadleaf Weed Control

Apply 1.3 to 6 quarts of this product in 20 to 100 gallons of spray solution per acre. Apply any time during the growing season.

Tank Mixing: This product at 1.3 to 4 quarts per acre may be tank mixed with 2 to 4 quarts of Tordon K; Tordon 101 Mixture, or 2,4-D 3.8 lb amine or low volatile ester to improve the spectrum of activity. When tank mixing, refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

AERIAL APPLICATION (HELICOPTER ONLY)

Aerial sprays should be applied using suitable drift control. See the Spray Drift Management section for drift control advice. Add an agriculturally approved non-ionic surfactant. See the **Spray Additives** section of **Application Equipment and Techniques** sections. See **Maximum Application Rates Table** in the **Application Directions** for relationship between mixing rate, spray volume, and maximum application rate.

FOLIAGE TREATMENT (RIGHTS-OF WAY)

Apply 2/3 gallon of this product per acre alone or in a tank mix with 1 to 1.5 gallons per acre of this product with 1 to 2 gallons of 2,4-d 3.8 lb amine, or low

volatile esters; or Tordon 101 Mixture. Apply in total spray volume of 10 to 30 gallons per acre.

When tank mixing, refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

FOREST MANAGEMENT APPLICATIONS

For broadcast applications, apply the recommended rate of this product in 10 to 25 gallons per acre by air on in 10 to 100 gallons per acre by ground. Use sufficient spray volumes to provide thorough plant coverage. To improve spray coverage at volumes less than 50 gallons per acre, add an agriculturally approved nonionic surfactant. See the **Spray Additives** section of **Application Directions**. Use application systems designed to prevent spray drift to off-target sites. Nozzles or additives used for drift minimization that produce larger droplets may require higher spray volumes to provide brush control. See the **Application Equipment and Techniques** section.

Forest Site Preparation (Not for Conifer Release)

To control susceptible woody plants and broadleaf weeds, apply up to gallons per acre of this product in a total spray solution of 10 to 30 gallons per acres. This product may be applied at a rate of 1 to 1.5 gallons per acre in a tank mix combination with 1 to 2 gallons of Tordon 101 Mixture or 2,4-D 3.8 lb volatile ester to broaden the spectrum of woody plants and broadleaf weeds controlled. Use of nonionic agricultural surfactant is recommended for all foliar applications. See the **Spray Additives** section of **Application Directions**.

Refer to the individual product labels for precautionary statement, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

Conifer Plant Back Interval: Conifer injury may occur if conifers are planted sooner than 1 month after product treatments at rates $<1\frac{1}{3}$ gallon per acre or if conifers are planted sooner than 2 months after treatment with rates of $1\frac{1}{3}$ to 3 gallons per acre. When herbicide tank mixtures are used for forest site preparation, use the longest plant back waiting period recommended on any tank mix partner.

Directed Spray Applications for Conifer Release

To release conifers from competing hardwoods such as red maple, sugar maple, striped maple, sweetgum, red and white oaks, ash, hickory, alder, birch, aspen, pin cherry, mix 1 to 2 gallons of this product in enough water to make 100 gallons of spray mixture. To improve spray coverage, add an agriculturally approved nonionic surfactant. See the **Spray Additives** section of **Application Directions**.

Direct the spray onto foliage of competitive hardwoods using knapsack or backpack sprayers with flat fan nozzles or equivalent. Make applications any

time after hardwoods have reached full leaf size, but before autumn coloration. The majority of treated hardwoods should be less than 6 feet in height to ensure adequate spray coverage. Care should be taken to direct the spray solution away from conifer foliage, particularly foliage of desirable pines.

Conifer Release Applications: Spray may cause temporary damage and growth suppression of conifers where direct contact occurs; however, injured conifers should recover and grow normally. **Over-the-top spray applications can kill pines.**

Broadcast Application for Conifer Release in the Northeastern United States

To release spruce, fir, red pine, and white pine from competing hardwoods such as red maple, sugar maple, striped maple, alder, birch (white, yellow and grey), aspen, ash, pin cherry, and *Rubus* spp. And perennial and annual broadleaf weeds, apply this product at 2 to 4 quarts per acre alone or in a tank mix with 2,4-D amine, or 2,4-D low volatile ester. Apply no more than 4 pounds acid equivalent per acre from the combined products. Make applications in late summer or early fall after conifers have formed their over-wintering buds; and hardwoods are in full leaf prior to autumn coloration.

Broadcast Applications for Douglas Fir Release in the Pacific Northwest and California

To release Douglas fir from competing vegetation such as broadleaf weeds, alder, blackberry or Scotch broom, apply this product at 1-1/3 to 2 quarts per acre alone or in combination with 4lb per acre of atrazine. Add a nonionic surfactant to the spray solution. See the **Spray Additives** section of **Application Directions**. Applications should be made in early spring after hardwoods begin growth and before Douglas fir bud break ("early foliar" hardwood stage). Applications can also be made in late summer, after Douglas fir seasonal growth has "hardened off" (winter bud set). Make applications while hardwoods are still actively growing. When treating after Douglas fir winter bud set, apply prior to onset of hardwood autumn coloration.

Note: Treatments applied during active Douglas fir shoot growth) after spring bud break and prior to winter bud set), may cause injury to Douglas fir trees.

Cut Surface Treatments

To control hardwood unwanted species such as elm, maple, oak; and conifers in right-of-way and other non-crop areas, apply this product, either undiluted or diluted in a 1:1 ratio with water by one of the following methods:

Tree Injector Method: Inject ½ milliliter (ml) of undiluted product or 1ml of diluted (1:1) solution through the bark at intervals of 3-4 inches between injection wounds. The tree injections should completely surround the tree at any convenient height.

Note: Worker Protection Standard **Agricultural Use Requirements** reentry restrictions do not apply for this application method. Refer to the **Non-Agricultural Use Requirements** box.

Hack and Squirt Method: Use a hatchet or similar equipment to make cuts in the bark at intervals of 3-4 inches at a convenient height around the circumference of the tree trunk. Spray ½ milliliter (ml) of undiluted product or 1 ml of the diluted (1:1) solution into each cut.

Frill or Girdle Method: Make a single girdle though the bark completely around the tree at a convenient height. Wet the cut surface with undiluted product or diluted (1:1) solution.

Both the **Hack and Squirt Method**; and the **Frill or Girdle Method** may be successfully used during any season except during periods of heavy sap flow of certain species such as maples.

Stump Treatment: Spray or paint undiluted product on to the freshly cut surfaces of cut stumps and stubs. The cambium area next to the bark is most vital area to wet.

TABLE 1
WOODY PLANTS CONTROLLED BY THIS PRODUCT

Alder	Chinquapin	Maleleuca	Sweetbay Magnolia
Arrowwood	Choke Cherry	Manles	Sweet Gum
Ash	Cottonwood	Mulberry	Sycamore
Aspen	Crataegus	Oaks	Tan Oak
Bear Clover	Dogwood	Persimmon	Thimbleberry
Beech	Douglas fir	Pine	Tulip Poplar
Birch	Elderberry	Poison Ivy	Wax Myrtle
Blackberry	Elm	Poison Oak	Western Hemlock
Black gum	Gallberry	Poplar	Wild Rose
Brazilian Pepper	Hazel	Salmonberry	Willow
Cascara	Hornbeam	Salt-bush (Baccharis)	Winged elm
Ceanothus	Kudzu ¹	Sassafras	
Cherry	Locust	Scotch Broom	
Chinese Tallow	Madrone	Sumac	

¹For complete control, retreatment may be necessary.

TABLE 2
ANNUAL AND PERENNIAL BROADLEAF WEEDS CONTROLLED BY THIS PRODUCT

Bindweed	Dandelion	Plantain	Tropical Sodaapple
Burdock	Elephant Ear	Purple Loosestrife	Vetch
Canada Thistle	Field Bindweed	Ragweed	Wild Lettuce
Chickory	Lambsquarter	Smartweed	
Curly Dock	Ligodium	Tansy Ragwort	

WETLAND SITES IN PRODUCTION FORESTS AND INDUSTRIAL NON-CROP AREAS

This product may be used in wetlands within forests; wildlife habitat restoration, wildlife management areas, and industrial non-crop sites; as well as areas adjacent to or surrounding domestic water supply reservoirs, lakes and ponds to control target vegetation in and around standing water sites, such as flood plains, delta, marshes, wetlands, swamps, bogs and transitional areas between upland and lowland sites, and the banks of ponds and lakes and transition areas between upland and lowland sites.

For control of woody plants and broadleaf weeds in these sites, follow use directions and application methods on this label for **Forest Management Application**. Refer to Tables 1 and 2 for lists of woody plants and broadleaf weeds that are controlled by this product.

General Use Precautions for Wetland Sites

- Refer to the **General Use Precautions** section for additional precautions.
- Minimize overspray to open water when treating target vegetation in and around non-flowing, quiescent or transient water. When making applications to control unwanted plants on banks or shorelines of flowing water, minimize over spray to open water.
- **Obtain Required Permits:** Before applying this product in and around public water, consult appropriate local public water control authorities. Permits may be required to treat such areas.
- **Recreational Use of Water in the Treatment Area:** There are no restrictions on water use in the treatment area for recreational purposes, including swimming and fishing.
- **Livestock Use of Water from Treatment Area:** There are no restrictions on consumption of water from treated areas by livestock.

Purple Loosestrife (*Lythrum salicaria*)

Purple loosestrife can be controlled with broadcast foliar application of this product at a minimum of 6 to 8 quarts per acre. Apply when purple loosestrife is at the bud or mid-flowering stage of growth. Follow-up applications for control of regrowth should be made the following year to achieve increased control of this weed species. For all applications, add a nonionic surfactant labeled for aquatics to the spray mixture. Follow all directions and use precautions on the surfactant label.

Thorough wetting of the foliage and stems is necessary to achieve satisfactory control. A minimum spray volume of 50 gallons per acre is recommended for ground broadcast applications.

For back pack applications, a spray solution of 1 to 1.5% of this product (5 to 7.6 fluid ounces of this product per 4 gallons of water) should be used. All purple loosestrife plants should be thoroughly wetted.

Aerial application by helicopter may be needed when treating restoration sites that are inaccessible, remote, difficult to traverse, isolated, or otherwise unsuited to ground application, or in circumstances where invasive exotic weeds dominate native plant populations over extensive areas and efforts to restore native plant diversity are being conducted. By air, apply in a minimum spray volume of 30 gallons per acre using Thru-Valve or Microfoil boom only.

Terrestrial Sites Associated with Wetland Areas

Refer to Tables 1 and 2 for a list of woody plants and broadleaf weeds that are controlled by this product.

Apply this product at rates of 0.25 to 2 gallons per acre for the control of broadleaf weeds and woody plants. Apply in enough water to provide uniform and complete coverage of the plants to be controlled. Use only water suitable for spraying. Use of an agriculturally approved nonionic surfactant is recommended for all foliar applications. Refer to **Spray Additives** section of **Application Directions** section. Refer to **Tank Mixing Recommendations** section for the order of addition of surfactants. For best results make application when woody plants and weeds are actively growing.

Use higher rates within the range when brush averages 15 feet or more in height or when brush covers > 60% of the area to be treated. Re-sprouting may occur the year following treatment if lower rates are used on hard-to-control species.

For hard-to-control species such as ash, black gum, choke cherry, maples, or oaks; during late summer applications when plants are mature; or during drought conditions; use higher rates of this product alone or use in combination with 2,4-D approved for aquatic use, generally the higher rates should be used for satisfactory brush control. When tank mixing, refer to the individual product labels for precautionary statements, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

General Use Precautions for Wetland Sites

- Refer to the General Use Precautions section for additional precautions.
- If applied to areas where livestock will graze, including right-of-way or fence rows do not apply more than 2/3 gallon of this product per acre per year.
- For forestry uses, DO NOT apply more than 2 gallons of this product per acre per year.

High Volume Foliage Applications

For control of woody plants, apply this product at 1 to 2 gallons per 100 gallons of spray solution. Make application in 100 to 400 gallons of total spray per acre depending on size and density of woody plants. Coverage should be thorough to wet all leaves, stems and root collars.

Tank Mixing: 1 to 4 quarts of this product may be tank mixed with 1 to 2 quarts of 2,4-D 3.8 lb amine diluted to make 100 gallons of spray solution. Make applications in 100 to 400 gallons of total spray per acre depending on size and density of woody plants. When tank mixing, refer to the individual product labels for precautionary statement, restrictions, recommended rates, approved uses, and a list of weeds and woody plants controlled.

Low Volume Foliage Applications

For control of woody plants, mix up to 5 gallons of this product in 10 to 100 gallons of spray solution. Adjust the spray concentration of this product and total spray volume per acre to match the size and density of target woody plants and kinds of spray equipment used. With low volume sprays, use sufficient spray volume to obtain uniform coverage of target plants including the surfaces of all foliage, stems, and root collars. For best results, a labeled aquatic surfactant should be added to all spray mixtures.

Match equipment and delivery rate of spray nozzles to height and density of woody plants. When treating tall, dense brush, a truck mounted spray gun with spray tips that deliver up to 2 gallons per minute at 40 to 60 PSI may be required. Backpack or other types of specialized spray equipment with spray tips that deliver less than 1 gallon of spray per minute may be appropriate for short, low to moderate density brush.

Cut Surface Treatments (Woody Plants)

To control unwanted trees and other listed woody plants in Table 1, apply this product, either undiluted or diluted in a 1:1 ration with water by one of the following methods:

Tree Injector Method: Inject ½ milliliter (ml) of undiluted product or 1ml of the diluted (1:1) solution through the bark at intervals of 3-4 inches between injection wounds. The tree injections should completely surround the tree at any convenient height.

Note: Worker Protection Standard **AGRICULTURAL USE REQUIREMENTS** reentry restrictions do not apply for this application method. Refer to the **NONAGRICULTURAL USE REQUIREMENTS** box.

Hack and Squirt Method: Use a hatchet or similar equipment to make cuts in the bark at intervals of 3-4 inches at a convenient height around the

circumference of the tree trunk. Spray ½ milliliter (ml) of undiluted product or 1 ml of the diluted (1:1) solution into each cut.

Frill or Girdle Method: Make a single girdle though the bark completely around the tree at a convenient height. Wet the cut surface with undiluted product or diluted (1:1) solution.

Both the **Hack and Squirt Method**; and the **Frill or Girdle Method** may be successfully used during any season except during periods of heavy sap flow of certain species such as maples.

Stump Treatment: Spray or paint undiluted product on to the freshly cut surfaces of cut stumps and stubs. The cambium area next to the bark is most vital area to wet.

AQUATIC SITES – Other than Rice

This product can be used to control emersed, submersed, and floating aquatic plants in aquatic sites such as ponds, lakes, reservoirs, non-irrigation canals, and ditches (with little or no continuous outflow), marshes, and wetlands. This product can also be used to control broadleaf and woody vegetation on banks and shores within or adjacent to these and other aquatic sites.

AQUATIC WEEDS CONTROLLED BY THIS PRODUCT

Alligatorweed	Milfoil species	Purple loosestrife
American lotus	Nuphar	Waterhyacinth
American frogbit	(spatterdock)	Waterlilly
Aquatic sodapple	Parrotfeather ¹	Waterprimrose
Eurasian watermilfoil	Pickerelweed	
	Pennywort	

¹Retreatment may be needed to achieve desired level of control

GENERAL USE PRECAUTIONS FOR AQUATIC SITES

- Refer to the **General Use Precautions** section for additional precautions.
- **Obtain Required Permits:** Before applying this product to public waters, consult with appropriate state or local water authorities. State or local agencies may require permits.
- Do not use treated water for irrigation for 120 days following application. As an alternative to waiting 120 days, treated water may be used for irrigation once the triclopyr level in the intake water is determined to be non-detectable by laboratory analysis (immunoassay). There is no restriction on use of water from the treatment area to irrigate established grasses.

- **Recreational Use of Water in the Treatment Area-** There are no restrictions on water use in the treatment area for recreational purposes, including swimming and fishing.
- **Livestock Use of Water from Treatment Area-** There are no restrictions on consumption of water from treated areas by livestock.

FLOATING AND EMERGED AQUATIC WEEDS

Surface Application: Use a spray boom, handgun or other similar suitable equipment mounted on a boat vehicle. Thorough wetting of foliage is essential for maximum effectiveness. Use 20 to 200 gallons per acre of spray mixture. Special precautions such as the use of low spray pressure, large droplet producing nozzles or addition of a labeled thickening agent may minimize spray drift in areas near sensitive crops.

Aerial Application (Helicopter only): Apply using a Microfoil or Thru-Valve boom, or a drift control additive in the spray solution. Apply in a minimum of 10 gallons of total spray solution per acre. Do not apply when weather conditions favor drift to sensitive areas. See the **Spray Drift Management** section for drift control advice.

Apply 0.5 to 2 gallons of this product per acre as a foliar application for control of waterhyacinth, alligatorweed (see specific directions below), and other susceptible emerged and floating herbaceous weeds and woody plants. Make applications using surface or aerial equipment. Use higher rates in the rate range when plants are mature, when the weed mass is dense, or for difficult to control species. Repeat treatments may be necessary to control regrowth and weeds which escaped spray, but do not apply more than 2 gallons of this product per acre per annual growing season. Make application when plants are actively growing.

Use of nonionic surfactant in the spray solution is recommended to improve control. Follow all directions and use precautions on the aquatic surfactant label.

FLOATING AND EMERGED WEED CONTROL – THIS PRODUCT RATES

Weed Species	Scientific Name	Gallons Per Acre	Application Timing and Remarks
Waterhyacinth	Eichhornia crassipes	0.5 – 2	Apply when plants are actively growing. Use the higher rate when the weed mass is dense. Thoroughly wet all foliage. Repeat treatments may be needed to control regrowth or escaped plants.
Alligatorweed	Alternanthera philoxeroides	0.75 – 2	Thoroughly wet all foliage. Weeds growing outside the margins of a body of water can be controlled. Alligatorweed growing in water will be only partially controlled. Top growth above water will be controlled, but plants will likely regrow from underwater tissue. Use a nonionic aquatic surfactant for best results.

POTABLE WATER INTAKE SETBACKS FOR CONTROL OF FLOATING AND EMERGED WEEDS – LAKES, RESERVOIRS, OR PONDS

Minimum setback distances from functioning potable water intakes for human consumption for the application of this product must be observed when controlling floating and emerged weeds in lakes, reservoirs or ponds. These setback restrictions do not apply to terrestrial application made adjacent to potable water intakes. Existing potable water intakes which are no longer in use are not considered to be functioning and these setback restrictions do not apply. Examples of this would be potable water intakes replaced by potable water wells or connections to a municipal water system.

The following table provides minimum setback distances based on the product rate and the area treated for floating and emerged weed control.

POTABLE WATER INTAKE SETBACK DISTANCES FOR APPLICATION OF THIS PRODUCT FOR CONTROL OF FLOATING AND EMERGED WEEDS IN LAKES, RESERVOIRS, OR PONDS

Minimum Setback Distances (feet)				
	Product Rate (quarts/acre)			
Area Treated (acres)	2	4	6	8
<4	0	200	400	500
>4 to 8	0	200	700	900
>8 to 16	0	200	700	1000
>16	0	200	900	1300

This product can be applied around functioning potable water intakes or closer than these setback distances as long as the intake is turned off until the level of triclopyr in the intake water is determined to be less than or equal to 0.4 parts per million (ppm) as determined by laboratory analysis or immunoassay.

SUBMERGED WEEDS – CONTROL OF EURASIAN WATERMILFOIL AND OTHER SUSCEPTIBLE SPECIES

Subsurface Application: This product can be applied directly into the water through boat-mounted distribution systems. Subsurface application may be desirable near areas of susceptible crops or other desirable broadleaf plants to avoid spray drift. Refer to Table to determine the desired amount.

Surface Application: This product can be applied either as a concentrate or as a spray solution diluted in water. Use a minimum spray volume of 5 gallons per acre. Do not apply when weather conditions favor drift to sensitive areas. See the **Spray Drift Management** section for drift control advice.

Apply 0.75 to 2.5 PPM ace (ae) of this product for control of Eurasian watermilfoil (*Myriophyllum spicatum*) and other susceptible submerged weeds in ponds, lakes, reservoirs, and in non-irrigation canals or ditches that have little or no continuous outflow. Make applications using surface or subsurface application. Use higher rates within the rate range in areas of greater water exchange. Repeat treatments may be necessary, but do not apply more than 2.5 PPM acid equivalent of this product per acre per annual growing season. Refer to following table to determine the desired amount.

THIS PRODUCT'S RATES FOR CONTROL OF SUBMERGED WEEDS IN PONDS, LAKES, RESERVOIRS, AND IN NON-IRRIGATION CANALS OR DITCHES					
Concentration of Triclopyr Acid Equivalent in Water (PPM ae)					
Navitrol gallons per surface area at specified depth					
Water Depth (feet)	0.75 ppm	1 ppm	1.5 ppm	2 ppm	2.5 ppm
1	0.7	0.9	1.4	1.8	2.3
2	1.4	1.8	3.3	3.6	4.6
3	2.1	2.9	4.1	5.4	6.8
4	2.7	3.6	5.4	7.2	9.1
5	3.4	4.5	6.8	9	11.3
6	4.1	5.4	8.1	10.9	13.6
7	4.8	6.3	9.5	12.7	15.8
8	5.5	7.2	10.9	14.5	18.1
9	6.1	8.1	12.2	16.3	20.4
10	6.8	9	13.6	18.1	22.6
15	10.2	13.6	20.4	27.2	33.9
20	13.6	18.1	27.2	36.2	45.3

POTABLE WATER INTAKE SETBACKS FOR CONTROL OF SUBMERGED WEEDS – LAKES, RESERVOIRS, OR PONDS

Minimum setback distance from functioning potable water intakes for human consumption for the application of this product must be observed when controlling submerged weeds in lakes, reservoirs or ponds. These setback restrictions do not apply to terrestrial applications made adjacent to potable water intakes. Existing potable water intakes which are no longer in use are not considered to be functioning and these setback restrictions do not apply. Examples of this would be potable water intakes replaced by potable water wells or connections to a municipal water system.

The following table provides the minimum setback distances based on the product rate and the area treated for submerged weed control.

Minimum Setback Distances (feet)					
	Concentration of This Product Acid Equivalent in Water (PPM ae)				
Area Treated (acres)	0.75 ppm	1 ppm	1.5 ppm	2 ppm	2.5 ppm
<4	300	400	600	800	1000
>4 – 8	420	560	840	1120	1400
>8 – 16	600	800	1200	1600	2000
>16 – 32	780	1040	1560	2080	2600
>32 acres, calculate the minimum setback distance using formula	Setback (ft) = $\frac{[800 \times \ln(\text{acres}) - 160]}{3.33}$	Setback (ft) = $\frac{[800 \times \ln(\text{acres}) - 160]}{2.5}$	Setback (ft) = $\frac{[800 \times \ln(\text{acres}) - 160]}{1.67}$	Setback (ft) = $\frac{[800 \times \ln(\text{acres}) - 160]}{1.25}$	Setback (ft) = $\frac{[800 \times \ln(\text{acres}) - 160]}{1}$

Example Calculations:

To apply this product at 2.5 PPM ae to 50 acres

$$\begin{aligned}
 \text{Setback in feet} &= [800 \times \ln(50 \text{ acres})] - 160 \\
 &= [800 \times 3.912] - 160 \\
 &= 2970 \text{ feet}
 \end{aligned}$$

To apply this product at 0.75 PPM ae to 50 acres

$$\begin{aligned}
 \text{Setback in feet} &= \frac{[800 \times \ln(50 \text{ acres})] - 160}{3.33} \\
 &= \frac{[800 \times 3.912] - 160}{3.33} \\
 &= 892 \text{ feet}
 \end{aligned}$$

This product can be applied around functioning potable water intakes or closer than these setback distances as long as the intake is tuned off until the level of triclopyr in the intake water is determined to be less than or equal to 0.4 parts per million (ppm) as determined by laboratory analysis or immunoassay.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store above 28F or agitate before use. Keep container closed when not in use. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Do not reuse or refill container. Do not contaminate feed, feedstuffs, or drinking water. Do not store or transport near feed or food.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

(For <5 gallon non-refillable containers only):

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse OR REFILL THIS product container. TRIPLE RINSE CONTAINER (OR EQUIVALENT) PROMPTLY AFTER EMPTYING. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank **AND DRAIN FOR 10 SECONDS AFTER THE FLOW BEGINS TO DRIP.** Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning if available or puncture and dispose of in approved landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Consult Federal, State or local authorities for approved alternative procedures.

(For >5 gallon non-refillable containers only):

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse OR REFILL THIS PRODUCT container. TRIPLE RINSE CONTAINER (OR EQUIVALENT) PROMPTLY AFTER EMPTYING. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ with water and recap. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. **TURN THE CONTAINER OVER ONTO ITS OTHER END AND TIP IT BACK AND FORTH SEVERAL TIMES.** Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning if available or puncture and dispose of in approved landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Consult Federal, State or local authorities for approved alternative procedures.

(For 275 Gallon refillable container only):

CONTAINER DISPOSAL: Refillable container. **REFILL THIS CONTAINER WITH PESTICIDE ONLY. DO NOT REUSE THIS CONTAINER FOR ANY OTHER PURPOSE.** Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat rinsing procedure two more times. Then offer for recycling or reconditioning if available or puncture and dispose of in approved landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Consult Federal, State or local authorities for approved alternative procedures.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. **THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT PERMITTED BY LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO CROP INJURY OR INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.**

LIMITATION OF LIABILITY

TO THE EXTENT PERMITTED BY LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF

INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS.