

U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg.

8959-57

Date of Issuance:

JAN 20 2011

NOTICE OF PESTICIDE:

x Registration
Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

AB Fluridone Aquatic Herbicide

Name and Address of Registrant (include ZIP Code):

Applied Biochemists W175 N11163 Stonewood Drive, Suite 234 Germantown, WI 53022-4799

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product when the Agency requires all registrants of similar products to submit data.
- 2. Make the following label changes listed below before you release the product for shipment:
 - a. Add the phrase, "EPA Reg. No. 8959-57".

Signature of Approving Official:

Product Manager (23)

Herbicide Branch

Registration Division (7505P)

Date:

JAN 20 2011

page 2 EPA Reg. No. 8959-57

b. In the Environmental Hazard section add the following statement as the first sentence:

Do not apply to water except as specified on the label.

- c. Delete the term "General" in heading on page 2 of the labeling. This term implies that the directions and restrictions do not all have to be followed as specified on the label. We have no objection to the phrase "Product Information".
- d. On page 5 in the directions for use with "other aquatic herbicides and algaecides" add a statement similar to "labeled for this use".
- e. According to PR Notice 82-2, the Tank Mix directions must include statements similar to the following:

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

ACCEPTED

with COMMENTS

AB FLURIDONE AQUATIC HERBICIDE

A herbicide for management of aquatic vegetation in fresh water ponds, lakes, reservoirs, potable water sources, drainage canals and irrigation canals.

Active Ingredient:

Fluridone: 1-methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-

pyridinone			In EPA Letter Dated: JAN 20 2011
Total			Under the Federal Insticide,
Contains 4 pounds active ingredient per gallon.			as amended, for the pesticide registered under EPA Reg. No.
	KEEP OUT OF REACH OF	CHILDREN	8959-57
	CAUTION/PRECA	UCION	0139-57
Si usted no entiende	la etiqueta, busqua a alguien para que se la ex this label, find someone to explain		ou do not understand
	FIRST AID		
If in eyes:	Hold eye open and rinse slowly and gently w		
	 Remove contact lenses, if present, after the Call a poison control center or doctor for treat 		ng eye.
If on skin or	Take off contaminated clothing.	inient advice.	
clothing:	Rinse skin immediately with plenty of water f	or 15-20 minutes.	
	Call a poison control center or doctor for treat		
If swallowed:	Immediately call a poison control center or d		
	Have person sip a glass of water if able to sy		
	Do not induce vomiting unless told to do so to		tor.
If inhaled	Do not give anything by mouth to an unconst	cious person.	
ii iiiilaieu	 Move person to fresh air. If person is not breathing, call 911 or an amb 	vulance then give artificial respirat	ion preferably
	mouth to mouth if possible.	uiance, men give annicial respirat	ion, preferably
	Call a poison control center or doctor for further	her treatment advice.	
	HOT LINE NUMBE		
medical emerge	t container or label with you when calling a poison oncy arises contact Arch Chemicals Emergency A call 423-780-2970. For help with a spill, leak, fire or	ction Network in the US call 1-8	300-654-6911 or call CHEMTREC
			(((((((((((((((((((
Manufactured for:			1766
Applied Biochemists		ς ι ι · · · · · · · · · · · · · · · · ·	· ·
An Arch Chemicals		(€ € € € € € ~ € €
	ewood Drive Suite 234	₩, €, ₹, €, €, €, €, €, €, €, €, €, €, €, €, €,	* (v
Germantown WI 53		. (()	(
800-558-5106	•	\$ 6 C	
EPA Reg. No. 895	9-57	EPA Est. No.	. (() () ()
	Net Contents: Gallor	ns () Liters	

Product of China - Formulated and Packaged in the United States with US and imported ingredients

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, absorbed through skin, or if inhaled. Causes moderate eye irritation. Avoid breathing of spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Follow use directions carefully so as to minimize adverse effects on nontarget organisms. Do not contaminate water when disposing of equipment washwaters. Trees, turf, and shrubs growing in water treated with this product may occasionally develop chlorosis. Do not apply in tidewater/brackish water. Lowest rates should be used in shallow areas where the water depth is considerably less than the average depth of the entire treatment site, for example, shallow shoreline areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

SHAKE WELL BEFORE USING.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. Do not store near feed or foodstuffs. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be used according to label directions or disposed of at an approved waste disposal facility.

container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ½ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

GENERAL INFORMATION

This product is a selective translocated aquatic herbicide. Applied to freshwater ponds, lakes, reservoirs, drainage canals and irrigation canals, This product helps manage undesirable aquatic weeds. Susceptible aquatic vascular plants absorb this product through the shoots and roots. For effective control, contact of this product with the target plants must be maintained for at least 45 days. Effective control is reduced if conditions exist that dilute the concentration of this product in the water.

The mode of action of this product involves inhibition of carotene synthesis in the target weeds. Lack of carotene in plants causes the chlorophyll to break down when the plants are exposed to sunlight. New shoot growth on target weeds begins to turn chlorotic (white) or pink in color within 7 to 10 days of exposure to this product. Ideally, 30 to 90 days of continuous exposure to this product will provide optimum control of target weeds. Some plant species may not be controlled by this product under all conditions. Factors affecting herbicide performance include growth stage of the target weed, the time of year when this product is applied, and dilution or movement of treated water.

Optimum results are achieved when this product is applied before weeds begin to actively grow. For mature plants, the higher application rates will be required and effects due to this product treatment will take longer to observe.

A suitable analysis of the water to determine the concentration of this product is highly recommended. The most common method of water analysis for measuring fluridone concentrations which is recommended is the Enzyme-Linked Immunoassay (ELISA Test). Contact Applied Biochemists for information on this test when using this product in treatment programs.

Application rates are shown in fluid ounces or quarts of this product to achieve a desired concentration of the active ingredient in parts per billion by weight (ppbw).

PRECAUTIONS

- **Permits:** Consult with appropriate state or local water authorities before applying this product since state or local agencies may require permits to use this product.
- Chemigation: Do not apply this product through any type of irrigation system.
- Hydroponic Farming: Do not use this product treated water for hydroponic farming.
- Greenhouse and Nursery Plants: Do not use water which has been treated with this product to irrigate greenhouse
 or nursery plants unless chemical assays of the water indicate fluridone residues are less than one parts per billion
 (ppb).
- Maximum Use Rates: Do not apply more than a total of 90 ppb in ponds and 150 ppb in lakes and reservoirs per annual growth cycle. These maximum concentrations are the amounts of fluridone calculated as the target application rate, NOT the concentration determined by analysis of fluridone in the treated water.
- Waiting Periods: If application rates are 150 ppb or less, treated water may be used immediately with no waiting period for drinking (potable) water (including watering livestock and pets), fishing or swimming. See specific restrictions below for Potable Water Intakes and Irrigation.
- Potable Water Intakes: In lakes and reservoirs or other sources of potable water, <u>DO NOT APPLY</u> this product at application rates greater than 20 ppb within 1/4 mile (1320 feet) of any functioning potable water intake. If rates are between 6 and 20 ppb, This product <u>MAY BE APPLIED</u> where functioning potable water intakes are present.
 Note: Existing potable water intakes which are no longer in use, such as those replaced by potable water wells or connections to a municipal water system, are not considered to be functioning potable water.
- Irrigation: Irrigation using water treated with this product may injure the irrigated vegetation. Instruct those who use
 treated water to follow the recommended waiting periods listed in the table below and to assay the water for fluridone
 residues. For crops grown on low organic and sandy soils and irrigated with treated water, the potential for crop
 injury is greater than for crops grown on heavier soils.

If a shorter waiting period is desired for irrigation of crops using treated water, use a suitable analysis (ELISA or other methods) to measure the concentration of fluridone in the treated water. If the concentration of fluridone is less than 10 ppb, established tree crops, established row crops or turf can be irrigated with this product treated water.

If the concentrations of fluridone are greater than 5 ppb, tobacco, tomatoes, peppers or other plants within the Solanaceae Family and newly seeded crops or newly seeded grasses such as over-seeded golf course greens should NOT be irrigated with treated water. Rotation Crops: Do not plant members of the Solanaceae family on land that has been previously irrigated with water containing more than 5 ppb of fluridone. Consult an aquatic specialist prior to commencing irrigation of such sites.

	Number of Days to Wait after This product Application Before Irrigating with treated Water			
Product Application Sites ^a	Established Tree Crops	Established Row Crops/Turf/Plants	Newly Seeded Crops/Seedbed or Areas To be Planted Including Overseeded Golf Course Greens	
Ponds and Static Canals	7	30	Assay required	
Canals	7	14	Assay required	
Lakes and Reservoirs	7	14	Assay required	

^aPonds: For this product labeling purposes, a pond is defined as a body of water 10 acres or less in size

Lakes or Reservoirs: For this product's labeling purposes, a lake or reservoir is defined as greater than 10 acres in size. When only one-half or more of the lake or reservoir is treated, follow the Pond and Static Canal precautions.

AQUATIC PLANT INFORMATION

Depending on the use rate, water movement, application timing, weed growth stage and application method, this product will control, partially control, or will not control certain aquatic plant species. The table below categorizes the species when this product is applied under ideal application conditions at higher to maximum label rates. When lower rates are used, certain species in the controlled or partially controlled categories will show increased tolerance to this product. Aquatic plants not listed may also be controlled, partially controlled, or be tolerant to this product.

Before applying this product, identify the aquatic plants to determine their susceptibility to this product.

Vascular Aquatic Plants Controlled by this Product

Floating Plants	Emersed Plants	Submersed Plants	Shoreline Grasses
common duckweed	spatterdock (Nuphar	bladderwort (Utricularia	paragrass (Urochloa
(Lemna minor)	luteum)	spp.)	mutica)
	water-lily (Nymphaea	common coontail	
	spp.)	(Ceratophyllum	ļ
		demersum)	
		common elodea (Elodea	
		canadensis)	
		egeria, Brazilian elodea	
		(Egeria densa)	
		fanwort, cabomba	
		(Cabomba caroliniana)	
·		hydrilla (<i>Hydrilla</i>	
		verticillata)	
		nalad (Najas spp.)	
		pondweed (Potamogeton	
		spp.,	
		except Illinois pondweed)	
		watermilfoil	
		(Myriophyllum spp.,	
		except variable-leaf]
		milfoil)	

Vascular Aquatic Plants Partially Controlled by this Product

Floating Plants	Emersed Plants	Submersed Plants	Shoreline Grasses
common watermeal (Wolffia columbiana) [†]	alligatorweed (Alternanthera philoxeroides)	Illinois pondweed (Potamogeton illinoensis)	barnyardgrass (Echinochloa crusgalli)
	American lotus (Nelumbo lutea)	limnophila (Limnophila sessiliflora)	giant cutgrass (Zizaniopsis miliacea)
	cattail (Typha spp.)	tapegrass,	reed canarygrass (Philaris arundinaceae)
	creeping waterprimrose (Ludwigia peploides)	American eelgrass (Vallisneria americana)	southern watergrass (Hydrochloa caroliniensis)
	parrotfeather (Myriophyllum aquaticum)	watermilfoil-variable-leaf milfoil (Myriophyllum heterophyllum)	torpedograss (Panicum repens)
	smartweed (Polygonum spp.)		
	spikerush (<i>Eleocharis</i> spp.)		
	waterpurslane (Ludwigia palustris)		
† Th:	watershield (Brasenia schreberi)		

This product when used at the maximum use rate only provides partial control of this species.

Vascular Aquatic Plants Not Controlled by this Product

Floating Plants	Emersed Plants	Submersed Plants	Shoreline Grasses
waterlettuce (Pistia	American frogbit		maidencane (Panicum
stratiotes)	(Limnobium spongia)		hemitomon)
	arrowhead (Sagittaria		
	spp.)		
	bacopa (Bacopa spp.)		
	big floatingheart,		
	banana lily (Nymphoides aquatica)		
	bulrush (Scirpus spp.)		-
· · · · · · · · · · · · · · · · · · ·	floating waterhyacinth		
	(Eichhornia crassipes)		
	pickerelweed, lanceleaf		•
	(Pontederia spp.)	l.	
	rush (Juncus spp.)		
	water pennywort		
	(Hydrocotyle umbellata)	<u> </u>	

*Note: This product does not control algae (Chara, Nitella, and single-cellular, colonial and filamentous species).

PREPARATION OF THIS PRODUCT'S SPRAY SOLUTIONS

Determine the amount of area (acres) to be treated. Water depths in the treatment sites should also be known so that the correct application rate is selected.

Use the steps below to prepare spray mixtures of this product:

- 1. Be sure to shake well the containers of this product before adding the product to the spray tank during mixing and loading operations.
- 2. Add ½ to ¾ the required amount of water to the spray tank. Begin agitation of the spray mixture and continue agitation during the mixing operations.
- 3. Add the required amount of this product to the spray tank during the remainder of the mixing operation.
- 4. Continue agitation of the spray mixture during the herbicide application operation.

Make surface or subsurface applications using conventional spray equipment. Use weighted trailing hoses to apply this product near the surface of the hydrosoil. Make applications with a spray volume of 5 to 100 gallons per acre. A metering system which mixes concentrated product with water and then introduces this slurry into the suction side of the application equipment may also be used.

Note: This product is not corrosive to application equipment.

Tank Mix Information

Tank mixes of this product with other aquatic herbicides and algaecides may provide greater efficacy and broader weed control or plant selectivity. Refer to the label for the herbicide or algaecide used as a tank mix with this product for use directions, precautions, and restrictions.

DIRECTIONS FOR APPLICATION - PONDS

For additional application rate calculations, refer to the section How To Calculate Application Rates at the end of this label.

Average Water Depth of Treatment Site in Feet	Fluid Ounces (or Qts.) of This Product per Treated Acre To Achieve Desired Herbicide Concentration:		Application Directions
<u> </u>	45 ppb	90 ppb	
1	3.8 fl. oz.	7.7 fl. oz.	Apply this product to the entire surface area of the pond.
	(0.12 qts.)	(0.24 qts.)	
2	7.7	15.7	Single Applications: Use the amount of this product listed to give 45 to 90 ppb fluridone in
· · · · · · · · · · · · · · · · · · ·	(0.24)	(0.49)	treated water. Higher rates should be used for
3	11.8	23.4	dense weed infestations, for difficult-to-control species, and for smaller ponds (less than 5
	(0.37)	(0.73)	acres in size and average water depths of less
4	15.7	31.4	than 4 feet).
	(0.49)	(0.98)	·
5	19.5	39.0	Split or Multiple applications: Use when dilution of the treated water is likely to occur.
	(0.61)	(1.22)	
6	23.4	46.7	Do not exceed 90 ppb per annual growth cycle.
	(0.73)	(1.46)	
7	27.2	54.4	
	(0.85)	(1.70)	
8	31.4	62.4	
	(0.98)	(1.95)	
9	35.2	70.1	
	(1.1)	(2.19)	
10	39.0	78.1	
	(1.22)	(2.44)	

DIRECTIONS FOR APPLICATION – LAKES AND RESERVOIRS

This product may be used for treatment of both whole lakes and reservoirs and partial areas of lakes or reservoirs (bays and coves). Target weeds in partial lake and reservoir treatments which are at least 5 acres in size are more effectively treated with this product than smaller size areas. Smaller treatment areas (less than 5 acres) or narrow strips such as boat trails or shorelines may not produce satisfactory results as this product may be diluted with untreated water. Due to a number of environmental factors, rate ranges are provided. Select the rates and application methods based on the specific goals of the aquatic plant management program at each different site.

Whole Lake or Reservoir Treatments (Limited or No Water Discharge)

Single Application to Whole Lakes or Reservoirs: Apply this product at an application rate of between 10 and 90 ppb. Consult the table below for the amount of this product required to achieve these concentrations in the treated water. Rates should be based on the goals of the aquatic plant management program.

If control of **Eurasian watermilfoil and curlyleaf pondweed** is desired or for greater plant selectivity, use an application rate lower in the range. For other plant species, contact an aquatic specialist to help determine when to choose lower application rate.

The higher rates within the rate range can be used when dense weed infestations are present or when treating hard-to-control weed species. Additional applications may be required to control more difficult-to-control species or when dilution of the treatment concentration has occurred such as from a heavy rainfall. If multiple applications are made, do not exceed 150 ppb (the sum of all applications) per annual growth cycle. Read the directions below on Split or Multiple Applications. For additional application rate calculations, refer to the section How To Calculate Application Rates at the end of this label.

Rates For Single Application of this Product

Rates For Single Application of this Product		
Average Water Depth at Treatment Site in Feet	Fluid Ounces (or Qts.) of This Product per Treated Acre To Achieve Desired Herbicide Concentration:	
	10 ppb	90 ppb
1	1.0 fl. oz.	7.7 fl. oz.
2	(0.03 qts.) 106	(0.24 qts.)
2	(0.05)	15.7 (0.49)
3	2.6	23.4
· ·	(0.08)	(0.73)
4	3.2	31.4
	(0.11)	(0.98)
5	4.5	39.0
	(0.14)	(1.22)
6	5.1	46.7
	(0.16)	(1.46)
7	6.1	54.4
	(0.19)	(1.70)
8	7.0	62.4
	(0.22)	(1.95)
9	7,6	70.1
	(0.24)	(2.19)
10	8.6	78.1
	(0.27)	(2.44)
11	9.6	86.0
12	(0.30) 10.2	(2.68) 93.8
12	(0.32)	(2.93)
13	11.2	101.4
19	(0.35)	(3.17)
14	12.1	109.4
• • •	(0.38)	(3.42)
15	13.1	117.1
	(0.41)	(3.66)
16	13.8	124.8
	(0.43)	(3.90)
17	14.7	132.2
	(0.46)	(4.15)
18	15.7	140.5
	(0.49)	(4.39)
19	16.3	148.2
·	(0.51)	(4.63)
20	17.3	156.2
	(0.54)	(4.88)

Split or Multiple Applications to Whole Lakes or Reservoirs: If the goal of the aquatic plant management program is to use the lowest effective rate and to maintain a low herbicide concentration for sufficient time to ensure efficacy and enhanced selectivity, split or multiple application programs are appropriate. However, water analyses using ELISA (or other analyses) must be carried out to ensure that the water is treated at an initial application rate of 6 to 50 ppb. Continue split applications to maintain a sufficient concentration of fluridone for a minimum of 45 days or longer. As with single applications, to control Eurasian watermilfoil and curlyleaf pondweed and to provide greater plant selectivity, use an application rate lower in the rate range. For other weed species, contact an aquatic specialist to help determine when to choose lower application rate.

A single application at no more than 20 ppb may be made to lakes or reservoirs comaining functional potable water intakes within ¼ mile of these functioning potable water intakes. Do not apply more than 150 ppb (sum of all applications) per annual growth cycle.

Partial Lake or Reservoir Treatments

If the chance of dilution of this product with untreated water is expected in partial lake or reservoir treatments, using split or multiple applications may extend the herbicide contact time with the target weeds. Use higher application rates and more frequent applications if the likelihood of untreated water diluting this product concentration in the treatment area is anticipated.

Refer to the table below for additional application instructions and for use rates. For additional application rate calculations, refer to the section How To Calculate Application Rates at the end of this label.

Partial Lake or Reservoir Treatment Site	Rates and Instructions
Treatment Areas Greater Than 1/4 Mile from a Functioning Potable Water Intake	Single applications: Apply this product at 30 to 150 ppb. Split or multiple applications: Do not exceed 150 ppb (total of all applications) per annual growth cycle. If split applications are made, maintain a sufficient concentration in the target area for a period of 45 days or longer. Use the ELISA or other analyses to ensure that the desired concentration of fluridone is maintained over time.
Treatment Areas Within 1/4 Mile of a Functioning Potable Water Intake	One may apply a concentration of greater than 20 ppb if the application is made at least ¼ mile or more from the functioning potable water intake. Application rates of less than 20 ppb may be made with ¼ mile of the potable water intake but use ELISA or other methods to verify that the fluridone concentration do not exceed 150 ppb at the potable water intake.

DIRECTIONS FOR APPLICATION - DRAINAGE CANALS AND IRRIGATION CANALS

For additional application rate calculations, refer to the section How To Calculate Application Rates at the end of this label.

Application Site	Rates and Instructions
Static Canals	1 – 2 quarts per treated acre
Moving Water Canals	Optimum performance will be achieved when water flow is restricted or reduced.
	For slowly moving bodies of water, apply using techniques that maintain the fluridone concentration at 15-40 ppb for at least 45 days. Use split or multiple broadcast applications (or metering methods) to ensure a uniform concentration of fluridone. Use the ELISA or other analyses to ensure that the desired concentration of fluridone is maintained over time.
Static or Moving Water Canals Containing a Functioning Potable Water Intake	One may apply a concentration of greater than 20 ppb this product at least ¼ mile or greater from the functioning potable water intake. Application rates of less than 20 ppb may be made with ¼ mile of the potable water intake but use ELISA or other methods to verify that the fluridone concentration do not exceed 150 ppb at the potable water intake.

HOW TO CALCULATE APPLICATION RATES

Ponds, Lakes and Reservoirs: Use the calculation below to determine the amount in fluid ounces of this product to be applied per acre to provide the desired ppb concentration of fluridone in the treated water:

Fluid Ounces of this product required per treated acre =

(Average water depth of treatment site in feet) x (desired ppb concentration of fluridone)

x 0.0027 x 32

As an example, the calculation to determine the number of fluid ounces of this product needed to treat one acre for a herbicide concentration of 45 ppb fluridone at a site where the average water depth is 3 feet is shown as follows:

 $3 \times 45 \times 0.0027 \times 32 = 11.7$ fl. oz. per treated acre

Note: Fluid ounces can be converted to quarts by dividing the number of fluid ounces by 32. For example, 11.7 fl. oz. ÷ 32 = 0.37 quarts.

Make sure that the calculated rate does not exceed the maximum allowable rate in pints (or quarts) per treated acre for the water depth listed in the application rate tables for the sites to be treated.

Moving Water Drainage and Irrigation Canals: Calculate the amount of this product in quarts required for the proposed application through a metering system to provide the desired ppb concentration of fluridone in the treated water as follows:

- Determine the Cubic Feet per Second as follows:
 CFS (cubic feet per second) = Average flow rate (feet per second) x average canal width (ft.) x average canal depth (ft.) x 0.9
- 2. Calculate the Water Movement in Acre-Feet per Day:
 Water movement in acre-feet per day = CFS x 1.98
- Amount of this product required:
 Acre-feet per day x desired ppb x 0.0027 = Quarts of this product required per day

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

The directions on this label are believed to be reliable and must be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, all of which are beyond the control of Applied Biochemists or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. Applied Biochemists, warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use, subject to the factors noted above which are beyond the control of Applied Biochemists. Except as warranted by this label Applied Biochemists. makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose. To the extent allowed by applicable law, the exclusive remedy against Applied Biochemists, for any cause of action relating to the handling or use of this product is a claim of damage, and in no event shall damages or any other recovery of any kind against Applied Biochemists, exceed the price of the product which causes the alleged loss, damage, injury, or other claim. To the extent allowed by applicable law, Applied Biochemists, shall not be liable and any and all claims against Applied Biochemists, are waived, for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income, whether or not based on the negligence of Applied Biochemists, breach of warranty, strict liability in tort, or any other cause of action. Applied Bjochemists, and the seller offer this product, and the buyer and users accept it, subject to the foregoing conditions of sale and limitations of warranty, liability and remedies.